



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07H 19/10, 19/20, A61K 31/70, C07F 9/6561, 9/6509, 9/6571, 9/6558, A61K 31/685, C07F 9/6564, 9/36, 9/40, 9/6512, 9/58, 9/09, 9/44, G01N 33 /53</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 95/07920</b> <b>(43) International Publication Date:</b> 23 March 1995 (23.03.95)
<b>(21) International Application Number:</b> PCT/US94/10539 <b>(22) International Filing Date:</b> 16 September 1994 (16.09.94)  <b>(30) Priority Data:</b> 08/123,483      17 September 1993 (17.09.93)      US 08/193,341      8 February 1994 (08.02.94)      US  <b>(71) Applicant (for all designated States except US):</b> GILEAD SCIENCES, INC. [US/US]; 353 Lakeside Drive, Foster City, CA 94404 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> BISCHOFBERGER, Norbert, W. [AT/US]; 105 Glasgow Lane, San Carlos, CA 94070 (US). JONES, Robert, J. [US/US]; 1020 Murchinson Drive, Millbrane, CA 94030 (US). ARIMILLI, Murty, N. [IN/US]; 4789 Ridgewood Drive, Fremont, CA 94555 (US). LIN, Kuei-Ying [US/US]; 4774 Canvasback Common, Fremont, CA 94555 (US). LOUIE, Michael, S. [US/US]; 1669 McKinley Street, San Mateo, CA 94403 (US). MCGEE, Lawrence, R. [US/US]; 1 Crater Lake Way, Pacifica, CA 94044 (US). PRISBE, Ernest, J. [US/US]; 1336 Richardson Avenue, Los Altos, CA 94024 (US). LEE, William, A.		[US/US]; 749 Anderson Drive, Los Altos, CA 94024 (US). CUNDY, Kenneth, C. [US/US]; 3802 Naughton Avenue, Belmont, CA 94002 (US).  <b>(74) Agents:</b> MUENCHAU, Daryl, D. et al.; Gilead Sciences, Inc., 353 Lakeside Drive, Foster City, CA 94404 (US).  <b>(81) Designated States:</b> AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LT, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> NUCLEOTIDE ANALOGS  <b>(57) Abstract</b>  Nucleotide analogs characterized by the presence of an amidate linked amino acid or an ester linked group which is bonded to the phosphorus atom of phosphonate nucleotide analogs are disclosed. The analogs comprise a phosphoamidate or ester bond that is hydrolysed <i>in vivo</i> to yield a corresponding phosphonate nucleotide analog. Methods and intermediates for the synthesis and use are described.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

## NUCLEOTIDE ANALOGS

5

### Background of the Invention

The present invention relates to novel nucleotide analog amidates and esters, their pharmaceutically acceptable acid addition salts, a process for their production, and to their use. The nucleotides of the present invention exhibit antitumor/antineoplastic activity, a broad spectrum of antimicrobial activity and certain other desirable activities.

Compounds related to the nucleotide analogs of the present invention may be found in: U.S. Patent Numbers 5,043,339, 5,108,994 and 5,166,198; EP 206 459; EP 253 412; EP 269 947; EP 270 885; EP 319 228; EP 343 133; EP 398 231; EP 404 296; EP 465 297; EP 468 119; EP 468 866; EP 479 640; EP 481 214; EP 494 370; EP 531 597; PCT/GB91/01171; PCT/US92/01020; PCT/US92/05208; WO 91/19721; Bronson et al, Bioorg Medicinal Chem Lett (1992) 2:685-690; Bronson et al, J Med Chem, (1989) 32:1457-1463; Bronson et al, Nucleotide Analogs as Antiviral Agents, ACS Symposium Series 401, J.C. Martin, Ed., p. 72-87, American Chemical Society, Washington, DC (1989); Colla, et al, J Med Chem (1983) 26:602-604; Curley, et al, Antiviral Res (1990) 14:345-356; De Clercq, et al, Nature, (1986) 323:464-467; Farrow, et al, J Med Chem (1990) 33:1400-1406; Farquhar, et al, J. Pharm Sci (1983) 72:324-325; Freed, et al, Biochem Pharmacol (1989) 19:3193-3198; Freeman, et al, J Med Chem (1992) 35:3192-3196; Gabrielsen, B., et al, Antiviral Res Suppl I (1992) 17:149; Gumpert, et al, Proc Natl Acad Sci (1971) 2559-2563; Juodka, et al, Coll Czech Chem Commun (1974) 39:963-968; Kim, et al, Bioorg Medicinal Chem Lett (1992) 2:367-370; Kim, et al, Tet Lett (1992) 33:25-28; Kim, et al, J Med Chem (1990) 33:1207-1213; Kumar, et al, J Med Chem (1990) 33:2368-2375; McGuigan, et al, Antiviral Chem Chemother (1993) 4:97-101; McGuigan, et al, Antiviral Res (1991) 15:255-263; Rosenberg, et al, Coll Czech Chem Commun (1988) 53:2753-2777; Rosenberg, et al, Coll Czech Chem Commun (1988) 52:2792-2800; Rosenberg, et al, Coll Czech Chem Commun (1988) 52:2801-2808; Starrett, et

al, Antiviral Res (1992) 19:267-273; Yu, et al, J Med Chem (1992) 35:2958-2969; Wolff-Kugel, et al, Tet Lett (1991) 32:6341-6344.

5 A characteristic of nucleotide analogs or nucleotides having a phosphonate or a phosphate group is the presence of one or two negative charges associated with the phosphorus group at physiologic pH. The charge associated with moieties such as phosphate or phosphonate groups is believed to generally limit bioavailability by limiting cell membrane permeation via passive diffusion (Liebman, et al, J. Biol. Chem., (1955) 216:823-830; Roll, et al, J Biol Chem, (1956) 220:439-444; Srivastava, et al, 10 Bioorg Chem (1984) 12:118-129; Palu, et al, Antiviral Res (1991) 16:115-119; Sastry, et al, Mol Pharmacol (1992) 41:441-445). These compounds are often, therefore, given parenterally in order to enhance bioavailability by increasing serum or intracellular levels.

15 Other characteristics of nucleotide analogs that can limit their efficacy include unfavorable pharmacokinetic or pharmacodynamic properties, insufficient potency and/or unfavorable toxicity characteristics.


Studies were conducted to ameliorate one or more of the above-mentioned problems associated with nucleotide analog drugs. The present invention includes novel nucleotide analogs that are hydrolyzable in vivo. 20 The nucleotide analogs can have improved bioavailability, improved pharmacokinetic or pharmacodynamic properties, enhanced potency and/or improved toxicity characteristics compared to the corresponding unmodified nucleotide analog. Methods to synthesize and use the compounds and methods to obtain and use antibodies that recognize the compounds are also 25 disclosed.

### Summary of the Invention


In a principal embodiment, the objects of this invention are accomplished by a nucleotide analog amidate comprising a phosphonate radical wherein the improvement comprises an amino acid residue or 30 polypeptide radical in which an amino group of the amino acid or polypeptide is bonded to the phosphorus atom of the nucleotide analog by an amidate bond, a carboxyl group of the amino acid residue or polypeptide radical is positioned such that it is capable as the free acid of hydrolyzing the 35 phosphoroamidate bond, and the carboxyl group is blocked (such as by

moieties including esters or amides). The nucleotide analog amidates of this invention are hydrolyzed in vivo to the corresponding nucleotide analog and are thus precursors of the corresponding nucleotide analog.


In accordance with this invention the nucleotide analog amidates or a physiologically acceptable salt thereof, have the structure of formula I



wherein  $L^1$  and  $L^2$  are independently an amino acid or polypeptide residue bonded to the phosphorus atom of the nucleotide analog by an amidate bond, or  $L^1$  or  $L^2$  are an oxyester, thioester, a substituted or unsubstituted amine, or hydroxy, provided that one or both of  $L^1$  and  $L^2$  is an amino acid or polypeptide residue and any carboxyl group that is linked by less than about 5 atoms to the amidate N is esterified or amidated, the dotted lines represent facultative bonds and wherein, (i) P and Z are linked to form a compound of the formula Ib



or (ii)  $L^1$  and Z are linked to form a compound of the formula Ic


$$I_c$$


wherein

5        substituents linked to carbon atoms designated # are in the *R*, *S* or *RS*  
configuration;


$X^1$  is O or S;

Z is  $-\text{CHR}^7-\text{R}^{11}-(\text{CH}_2)_{m1}-\text{C}^\#(\text{R}^8)((\text{CH}_2)_{m2}(\text{R}^9))-(\text{CH}_2)_{m3}-\text{R}^{10}-(\text{CH}_2)_{m4}-$ ,  
 $-\text{Q}-\text{C}_6\text{H}_4-\text{CH}_2-$ ,  $-\text{CHR}^7-\text{O}-\text{CHR}^7-\text{O}-\text{CHR}^7-$ ,  $-\text{CHR}^7-(\text{CHR}^{13})_{m1}-\text{CHR}^{14}-\text{R}^{10}-$ ,


10




IV




V



VI



VII







or VIII

wherein

- R<sup>7</sup> is H or C<sub>1</sub>-C<sub>4</sub> alkyl;
- 5 R<sup>8</sup> is H or C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>2</sub>-C<sub>4</sub> alkenyl, azidomethyl or azidoethyl;
- R<sup>9</sup> is halogen (F, Cl, Br or I), H or OH;
- R<sup>10</sup> is O, CH<sub>2</sub> or a chemical bond;
- R<sup>11</sup> is O, S, CH<sub>2</sub>, CHF or CF<sub>2</sub>;
- Q is -C(R<sup>12</sup>)<sub>2</sub>-CH<sub>2</sub>-, -C(R<sup>12</sup>)<sub>2</sub>-O-, -CR<sup>12</sup>=CR<sup>12</sup>-, or -C≡C-, wherein each R<sup>12</sup>
- 10 is independently H, or halogen;
- R<sup>13</sup> is H, halogen, OH, CH<sub>3</sub>, CH<sub>2</sub>OH, or C<sub>3</sub> - C<sub>6</sub> acyloxymethyl;
- R<sup>14</sup> is H, halogen, OH, CH<sub>3</sub>, CH<sub>2</sub>OH, C<sub>3</sub> - C<sub>6</sub> acyloxymethyl, or C<sub>2</sub> - C<sub>6</sub> acyloxy;
- R<sup>25</sup> is CH<sub>2</sub>, CHF or O;
- 15 R<sup>26</sup> is CH or S, provided that when R<sup>25</sup> is CH, R<sup>26</sup> is not S;
- R<sup>27</sup> is H, OH, halogen, N<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy or, when R<sup>26</sup> is S, R<sup>27</sup> is absent;
- R<sup>27a</sup> is H, OH, halogen, N<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy;
- R<sup>28</sup> is H, OH, halogen, N<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl or C<sub>1</sub>-C<sub>4</sub> alkoxy;
- 20 R<sup>29</sup> is O, S, CH<sub>2</sub>, CHF, CF<sub>2</sub>;
- R<sup>32</sup> is O;
- m<sub>1</sub> is an integer having a value from 0 to 4;
- m<sub>2</sub> is an integer having a value from 0 to 4;
- m<sub>3</sub> is an integer having a value from 0 to 4;
- 25 m<sub>4</sub> is an integer having a value from 0 to 4;
- B is a heterocyclic base; and
- substituents linked to the carbon atom designated C<sup>#</sup> are in the R, S or RS configuration.


- 30 In a further embodiment the objects are accomplished by compounds of the formula II, IIa, IIb and IIc



5

wherein  $L^2$  is OR, SR or





### III

$n$  is an integer having a value from 1 to 5 and if  $n > 1$ , each  $-C(R^3)(R^2)-$  may be the same or different;

5  $n1$  is an integer;

substituents linked to the carbon atom designated # are in the  $R$ ,  $S$  or  $RS$  configuration;

$R$  is  $H$ ,  $C_1$ - $C_{20}$  alkyl which is unsubstituted or substituted by substituents independently selected from the group consisting of  $OH$ ,  $O$ ,  $N$  and halogen ( $F$ ,  $Cl$ ,  $Br$ ,  $I$ ),  $C_3$ - $C_{20}$  aryl which is unsubstituted or substituted by substituents independently selected from the group consisting of  $C_1$ - $C_6$  alkyl,  $C_1$ - $C_6$  alkoxy,  $C_1$ - $C_6$  haloalkyl (1 to 3 halogen atoms), cyano, nitro,  $OH$ ,  $O$ ,  $N$  and halogen or  $R$  is  $C_4$ - $C_{20}$  aryl-alkyl which is unsubstituted or substituted in the aryl moiety by substituents independently selected from the group consisting of  $C_1$ - $C_6$  alkyl,  $C_1$ - $C_6$  alkoxy,  $C_1$ - $C_6$  haloalkyl (1 to 3 halogen atoms), cyano, nitro,  $OH$ ,  $O$ ,  $N$  and halogen, or  $R$  is  $C_3$ - $C_{24}$  1-acyloxy-1-alkyl ( $C_1$ - $C_8$  alkyl), or  $R$  is  $C_6$ - $C_{24}$  1-acyloxy-1-aryl-1-alkyl ( $C_1$ - $C_6$  aryl,  $C_1$ - $C_4$  alkyl), or  $R$  is  $C_3$ - $C_{24}$  1-acyloxy-2-alkoxy-1-alkyl ( $C_1$ - $C_8$  alkyl), or  $R$  is  $C_3$ - $C_{24}$  1-acyloxy-2-haloalkyl ( $C_1$ - $C_8$  haloalkyl, 1 to 3 halogen atoms);

20  $R^1$  is  $H$  or  $C_1$ - $C_9$  alkyl which is unsubstituted or substituted by substituents independently selected from the group consisting of  $OH$ ,  $O$ ,  $N$ ,  $COOR^4$  and halogen,  $C_3$ - $C_6$  aryl which is unsubstituted or substituted by substituents independently selected from the group consisting of  $OH$ ,  $O$ ,  $N$ ,  $COOR^4$  and halogen or  $C_3$ - $C_9$  aryl-alkyl which is unsubstituted or substituted by substituents independently selected from the group consisting of  $OH$ ,  $O$ ,  $N$ ,  $COOR^4$  and halogen;

R<sup>2</sup> is H or C<sub>1</sub>-C<sub>9</sub> alkyl which is unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N, COOR<sup>4</sup> and halogen, C<sub>3</sub>-C<sub>6</sub> aryl which is unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N, COOR<sup>4</sup> and halogen or C<sub>3</sub>-C<sub>9</sub> aryl-alkyl which is unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N, COOR<sup>4</sup> and halogen;

R<sup>3</sup> is C(O)-OR<sup>4</sup>, amino, amide, guanidiny, imidazolyl, indolyl, sulfoxide, phosphoryl, C<sub>1</sub> - C<sub>3</sub> alkylamino, C<sub>1</sub> - C<sub>3</sub> alkylidiamino, C<sub>1</sub> - C<sub>6</sub> alkenylamino, hydroxy, thiol, C<sub>1</sub> - C<sub>3</sub> alkoxy, C<sub>1</sub> - C<sub>3</sub> alkthiol, (CH<sub>2</sub>)<sub>n</sub>COOR<sup>4</sup>, C<sub>1</sub>-C<sub>6</sub> alkyl which is unsubstituted or substituted with OH, halogen, SH, NH<sub>2</sub>, phenyl, hydroxyphenyl or C<sub>7</sub> - C<sub>10</sub> alkoxyphenyl; C<sub>2</sub>-C<sub>6</sub> alkenyl which is unsubstituted or substituted with OH, halogen, SH, NH<sub>2</sub>, phenyl, hydroxyphenyl or C<sub>7</sub> - C<sub>10</sub> alkoxyphenyl; C<sub>6</sub>-C<sub>12</sub> aryl which is unsubstituted or substituted with OH, halogen, SH, NH<sub>2</sub>, phenyl, hydroxyphenyl or C<sub>7</sub> - C<sub>10</sub> alkoxyphenyl; and

R<sup>4</sup> is H provided that n<sub>1</sub> greater than 1, or is C<sub>3</sub>-C<sub>9</sub> alkyl which is substituted by substituents independently selected from the group consisting of OH, O, N and halogen, C<sub>3</sub>-C<sub>6</sub> aryl which is substituted by substituents independently selected from the group consisting of OH, O, N and halogen or C<sub>3</sub>-C<sub>9</sub> aryl-alkyl which is substituted by substituents independently selected from the group consisting of OH, O, N and halogen.

The structural formula I is meant to define compounds where the phosphorus (P) atom is tetravalent (PV oxidation state) and optionally linked via the facultative bonds shown as dotted lines to either L<sup>1</sup> or Z to form a heterocyclic ring containing at least the P atom itself and a nitrogen atom of L<sup>1</sup> or an atom present, usually oxygen (O), in Z. For such compounds, L<sub>2</sub> and the facultative bond between P and Z is absent. Such heterocyclic rings will preferably be 5-, 6- or 7-membered, but are also 4-, 8-, 9-, 10-, 11- or 12-membered. Alternatively the P atom is covalently linked to L<sup>2</sup> with L<sup>1</sup> and Z optionally linked to each other to form a heterocyclic ring. The structure is not intended to include compounds where L<sup>1</sup>, L<sup>2</sup> and a heterocyclic ring containing P and Z are present in the same molecule which would exceed the valency of P. Thus, an exemplary class of compounds is represented by the

structure of formula I includes  $(L^1)(L^2)P(O)-Z-B$  (formula Id) where no heterocyclic rings are formed between any  $L^1$ ,  $L^2$ , P, Z or B moiety.

$R^2$  includes methyl, ethyl, propyl, isopropyl and benzyl.

In another embodiment, the objects of this invention are accomplished by a nucleotide analog ester comprising a phosphonate radical and an ester moiety bonded to the phosphorus atom of the nucleotide analog. The nucleotide analog esters of this invention are hydrolyzed in vivo to the corresponding nucleotide analog and are thus precursors of the corresponding nucleotide analog, or can be used as intermediates in the synthesis of the nucleotide analog amidates.

The substructure Z can have a range of atoms between the base, B, and the phosphorus atom. For example, four atoms separate the heterocyclic base and phosphorus moieties when Z is of the formula  $-CH_2-O-CH_2-CH_2-$ . In general, there will be from 2 to 16 atoms, preferably from 3 to 9 atoms, more preferably from 4 to 6 atoms that separate the heterocyclic base and the phosphorus atom. Thus, Z substructures of the formula  $-CHR^7-R^{11}-(CH_2)_{m1}-C(R^8)((CH_2)_{m2}(R^9))-(CH_2)_{m3}-R^{10}-(CH_2)_{m4}-$  may be characterized where the sum of  $m1$ ,  $m3$  and  $m4$  is in a range between 0 and 12 or preferably in a range between 1 and 6, more preferably in a range between 1 and 4.

The nucleotide analog amidate and ester compounds of the instant invention include the corresponding salts, which may be base salts of the phosphonic acid moiety or an acid addition salt of the base in addition to the zwitterionic forms and/or solvates of compounds of formula I.

Some of the compounds of the present invention can exist as optical isomers and both racemic or scalemic and diastereomeric mixtures of these isomers which may exist for certain compounds as well as the individual optical isomers which are all within the scope of the present invention. Compounds of formula IIa in the *R*, *S* or *RS* configuration at the chiral carbon, designated # herein, are examples of compounds having optical isomers. While the scalemic mixtures can be separated into their individual isomers through well-known techniques such as, for example, the separation of diastereomeric salts formed with optically active adjuncts, e.g. acids or bases followed by conversion back to the optically active substrates; in most instances, for compounds of the present invention, the preferred optical

isomer can be synthesized by means of stereospecific reactions, beginning with the appropriate stereoisomer of the desired starting material.

As indicated, the present invention also pertains to the salts, including pharmaceutically acceptable non-toxic salts of these compounds. Such salts may include those derived by combination of appropriate cations such as alkali and alkaline earth metal ions or ammonium and quaternary amino ions with the acid anion moiety of the phosphonic acid group. In addition salts may be formed from acid addition of certain organic and inorganic acids with basic centers of the purine, specifically guanine, or pyrimidine base. Finally it is to be understood that compounds of the present invention in their un-ionized as well as zwitterionic form and/or in the form of solvates are also considered part of the present invention.

In other embodiments, the foregoing nucleotide analog amidates and esters or their dihydroxy phosphonate hydrolysis products are labeled with a detectable tag such as a radioisotope (including  $^{32}\text{P}$ ,  $^{35}\text{S}$ ,  $^{14}\text{C}$ ,  $^3\text{H}$ ,  $^{125}\text{I}$ ), a fluorescent moiety, an enzyme (including peroxidase, phosphatase) or the like.

In other embodiments, the foregoing nucleotide analog amidates comprise amino acid, dipeptide or tripeptide compounds (monosubstituted or disubstituted with identical or different amino acid, dipeptide or tripeptide substituents) that are capable of entry into eukaryotic cells via amino acid or peptide transporters present in eukaryotic cells *in vivo* or *in vitro*.

Also included are immunogens for raising antibodies which are capable of binding to the nucleotide analog amidates and esters of this invention and/or their dihydroxy phosphonate hydrolysis products, as well as antibodies capable of binding to the amidate and ester compounds of this invention or to their dihydroxy phosphonate hydrolysis products.

### Chemical Structures

Structural formulas and substructures are represented as roman numerals (I, II, III, IV, V, etc) or as letters (B, Z,  $\text{L}^1$ ,  $\text{L}^2$ ,  $\text{R}^1$ ,  $\text{R}^2$ , etc). The substructures Z and  $\text{Z}^1$  represent linking groups between the heterocyclic base (B) and the phosphorus atom (P) of the phosphonate group in the nucleotide analogs described herein. Linking groups Z, such as  $-\text{CHR}^7-\text{R}^{11}-(\text{CH}_2)_{m1}-\text{C}(\text{R}^8)((\text{CH}_2)_{m2}(\text{R}^9))-(\text{CH}_2)_{m3}-\text{R}^{10}-(\text{CH}_2)_{m4}-$ , in the structure  $(\text{L}^1)(\text{L}^2)\text{P}(\text{O})-\text{Z}-\text{B}$

have the structure  $(L^1)(L^2)P(O)-CHR^7-R^{11}-(CH_2)_{m1}-C(R^8)((CH_2)_{m2}(R^9))-(CH_2)_{m3}-R^{10}-(CH_2)_{m4}-B$  (i.e. the heterocyclic base (B) is covalently linked to the unfilled valence on the right side of the structure and the phosphorus atom is linked to the unfilled valence on the left side).

5

#### Brief Description of the Drawings

Figure 1. Synthesis of formula Ib compounds where  $X^1$  is S.

Figure 2. Synthesis of formula Ia compounds.

Figure 3. Synthesis of formula IV compounds.

10 Figure 4. Synthesis of formula VII compounds.

Figure 5. Synthesis of formula VIII compounds.

Figure 6. Synthesis of formula VI compounds.

Figure 7. Synthesis of formula VI compounds.

#### 15 Detailed Description of the Invention

Amino Acid Residues. When groups  $L^1$  or  $L^2$  comprise an amino acid residue they comprise any naturally-occurring or synthetic amino acid residue, i.e., any moiety comprising at least one carboxyl and at least one amino residue directly linked by at least one carbon atom, typically a single ( $\alpha$ ) carbon atom. The nature and identity of the intervening structure located between the carboxyl and amino (amidate) groups can have a variety of structures including those described herein. All that is necessary is that the group have sufficient conformation and length to be capable of acid catalysis of the phosphoroamidate bond and release of the phosphonate when the free carboxyl is generated *in vivo*, e.g. by deesterification, deamidation or peptidolytic cleavage of the precursor. In general, the amino acids corresponding to the residues employed in the compounds of this invention are naturally occurring and have no pharmacological activity *per se*. However, optimal pharmacokinetic activity (substantially complete autocatalytic hydrolysis upon hydrolysis of the distal amide or ester bond) may be achieved by using non-naturally occurring amino acid residues. The intervening structure may be as simple as methylene (when the residue is glycyl) or substituted methylene (other  $\alpha$  amino acids). The structure ordinarily contains up to about 5 carbon or hetero atoms in the direct linkage between the carboxyl carbon and the amidate nitrogen, as for example in the

20

25

30

35

case of intervening ethylene, propylene, butylene, or pentylene groups or their substituted analogs, such as for example oxyesters in which O replaces carbon and, as appropriate, hydrogen. An example of such an intervening structure would be  $-\text{CH}-\text{O}-\text{CH}(\text{R}^3)(\text{R}^2)-$ . In general, fewer intervening atoms  
5 are employed when more rapid hydrolysis is desired, although it will be understood that larger structures are suitable if they possess sufficient flexibility or have conformations in which the carboxyl group is positioned in proximity to the amidate bond.

In general, the amino acid residue has the structure shown in formula  
10 III. Ordinarily,  $n$  is 1 or 2,  $\text{R}^2$  is H and  $\text{R}^3$  is a moiety containing one or more of the following groups: amino, carboxyl, amide, carboxyl ester, hydroxyl,  $\text{C}_6$ - $\text{C}_7$  aryl, ether,  $n$ -,  $s$ - or  $t$ -alkyl ( $\text{C}_1$  -  $\text{C}_6$ ), guanidiny, imidazolyl, indolyl, sulfhydryl, sulfoxide, and phosphoryl. The  $\text{R}^2$  and  $\text{R}^3$  substituents can have a wide variety of structures including those disclosed herein.

15 Ordinarily  $\text{R}^2$  is H and  $\text{R}^3$  is a side chain or group of a naturally occurring amino acid. With respect to the carboxyl-containing side chains it will be understood that if the C atom of the subject carboxyl is linked by 5 or less atoms to the phosphoamide N then the carboxyl optionally will be blocked, e.g. by esterification or amidation wherein the ester or amide bonds  
20 are hydrolyzable in vivo.  $\text{R}^3$  also is taken together with  $\text{R}^1$  to form a proline residue ( $\text{R}^3 = -\text{CH}_2-$ )<sub>3</sub>). Thus,  $\text{R}^3$  is generally a side group such as H,  $-\text{CH}_3$ ,  $-\text{CH}(\text{CH}_3)_2$ ,  $-\text{CH}_2-\text{CH}(\text{CH}_3)_2$ ,  $-\text{CHCH}_3-\text{CH}_2-\text{CH}_3$ ,  $-\text{CH}_2-\text{C}_6\text{H}_5$ ,  $-\text{CH}_2\text{CH}_2-\text{S}-\text{CH}_3$ ,  $-\text{CH}_2\text{OH}$ ,  $-\text{CH}(\text{OH})-\text{CH}_3$ ,  $-\text{CH}_2-\text{SH}$ ,  $-\text{CH}_2-\text{C}_6\text{H}_4\text{OH}$ ,  $-\text{CH}_2-\text{CO}-\text{NH}_2$ ,  $-\text{CH}_2-\text{CH}_2-\text{CO}-\text{NH}_2$ ,  $-\text{CH}_2-\text{COOH}$ ,  $-\text{CH}_2-\text{CH}_2-\text{COOH}$ ,  $-(\text{CH}_2)_4-\text{NH}_2$  and  $-(\text{CH}_2)_3-\text{NH}-$   
25  $\text{C}(\text{NH}_2)-\text{NH}_2$ .  $\text{R}^3$  also includes 1-guanidinoprop-3-yl, benzyl, 4-hydroxybenzyl, imidazol-4-yl, indol-3-yl, methoxyphenyl and ethoxyphenyl. The optimal  $\text{R}^3$  group is readily selected using routine assays.

When the amino acid residues contain one or more chiral centers, any of the D, L, meso, threo or erythro (as appropriate) racemates, scalemates or  
30 mixtures thereof, fall within the scope of this invention. In general, if it is desired to rely on non-enzymatic means of hydrolysis, D isomers should be used. On the other hand, L isomers may be more versatile since they can be susceptible to both non-enzymatic as well as potential targeted enzymatic hydrolysis, and are more efficiently transported by amino acid or dipeptidyl  
35 transport systems in the gastrointestinal tract.

Examples of suitable amino acid residues include the following:

Glycyl;

Aminopolycarboxylic acids, e.g., aspartic acid,  $\beta$ -hydroxyaspartic acid, glutamic acid,  $\beta$ -hydroxyglutamic acid,  $\beta$ -methyiaspartic acid,  $\beta$ -methylglutamic acid,  $\beta,\beta$ -dimethyiaspartic acid,  $\gamma$ -hydroxyglutamic acid,  $\beta,\gamma$ -dihydroxyglutamic acid,  $\beta$ -phenylglutamic acid,  $\gamma$ -methyleneglutamic acid, 3-aminoadipic acid, 2-aminopimelic acid, 2-aminosuberlic acid and 2-aminosebacic acid residues;

Amino acid amides such as glutaminy and asparaginy;

Polyamino- or polybasic-monocarboxylic acids such as arginine, lysine,  $\beta$ -aminoalanine,  $\gamma$ -aminobutyric acid, ornithine, citrulline, homoarginine, homocitrulline, 5-hydroxy-2,6-diaminohexanoic acid (commonly, hydroxylysine, including allohydroxylysine) and diaminobutyric acid residues;

Other basic amino acid residues such as histidiny;

Diaminodicarboxylic acids such as  $\alpha,\alpha'$ -diaminosuccinic acid,  $\alpha,\alpha'$ -diaminoglutaric acid,  $\alpha,\alpha'$ -diaminoadipic acid,  $\alpha,\alpha'$ -diaminopimelic acid,  $\alpha,\alpha'$ -diamino- $\beta$ -hydroxypimelic acid,  $\alpha,\alpha'$ -diaminosuberlic acid,  $\alpha,\alpha'$ -diaminoazelaic acid, and  $\alpha,\alpha'$ -aminosebacic acid residues;

Imino acids such as proline, 4- or 3-hydroxy-2-pyrrolidinecarboxylic acid (commonly, hydroxyproline, including allohydroxyproline),  $\gamma$ -methylproline, pipercolic acid, 5-hydroxypipercolic acid,  $-N([CH_2]_nCOOR^4)_2$ , wherein n and  $R^4$  are as defined above, and azetidine-2-carboxylic acid residues;

A mono- or di-alkyl (typically  $C_1 - C_8$  branched or normal) amino acid such as alanine, valine, leucine, allylglycine, butyric acid, norvaline, norleucine, heptyline,  $\alpha$ -methylserine,  $\alpha$ -amino- $\alpha$ -methyl- $\gamma$ -hydroxyvaleric acid,  $\alpha$ -amino- $\alpha$ -methyl- $\delta$ -hydroxyvaleric acid,  $\alpha$ -amino- $\alpha$ -methyl- $\epsilon$ -hydroxycaproic acid, isovaline,  $\alpha$ -methylglutamic acid,  $\alpha$ -aminoisobutyric acid,  $\alpha$ -aminodiethylacetic acid,  $\alpha$ -aminodiisopropylacetic acid,  $\alpha$ -aminodi-n-propylacetic acid,  $\alpha$ -aminodiisobutylacetic acid,  $\alpha$ -aminodi-n-butylacetic acid,  $\alpha$ -aminoethylisopropylacetic acid,  $\alpha$ -amino-n-propylacetic acid,  $\alpha$ -aminodiisoamylacetic acid,  $\alpha$ -methyiaspartic acid,  $\alpha$ -methylglutamic acid, 1-aminocyclopropane-1-carboxylic acid; isoleucine, alloseucine, tert-leucine,

$\beta$ -methyltryptophan and  $\alpha$ -amino- $\beta$ -ethyl- $\beta$ -phenylpropionic acid residues;  $\beta$ -phenylserinyl;

Aliphatic  $\alpha$ -amino- $\beta$ -hydroxy acids such as serine,  $\beta$ -hydroxyleucine,  $\beta$ -hydroxynorleucine,  $\beta$ -hydroxynorvaline, and  $\alpha$ -amino- $\beta$ -hydroxystearic acid  
5 residues;

$\alpha$ -Amino,  $\alpha$ -,  $\gamma$ -,  $\delta$ - or  $\epsilon$ -hydroxy acids such as homoserine,  $\gamma$ -hydroxynorvaline,  $\delta$ -hydroxynorvaline and epsilon-hydroxynorleucine residues; canavinyl and canalinyl;  $\gamma$ -hydroxyornithinyl;

2-hexosaminic acids such as D-glucosaminic acid or D-galactosaminic  
10 acid residues;

$\alpha$ -Amino- $\beta$ -thiols such as penicillamine,  $\beta$ -thiolnorvaline or  $\beta$ -thiolbutyrine residues;

Other sulfur containing amino acid residues including cysteine; homocystine;  $\beta$ -phenylmethionine; methionine; S-allyl-L-cysteine sulfoxide;  
15 2-thiolhistidine; cystathionine; and thiol ethers of cysteine or homocysteine;

Phenylalanine, tryptophan and ring-substituted  $\alpha$  amino acids such as the phenyl- or cyclohexylamino acids  $\alpha$ -aminophenylacetic acid,  $\alpha$ -aminocyclohexylacetic acid and  $\alpha$ -amino- $\beta$ -cyclohexylpropionic acid; phenylalanine analogues and derivatives comprising aryl, lower alkyl,  
20 hydroxy, guanidino, oxyalkylether, nitro, sulfur or halo-substituted phenyl (e.g., tyrosine, methyltyrosine and o-chloro-, p-chloro-, 3,4-dichloro, o-, m- or p-methyl-, 2,4,6-trimethyl-, 2-ethoxy-5-nitro, 2-hydroxy-5-nitro and p-nitro-phenylalanine); furyl-, thienyl-, pyridyl-, pyrimidinyl-, purine or naphthylalanines; and tryptophan analogues and derivatives including  
25 kynurenine, 3-hydroxykynurenine, 2-hydroxytryptophan and 4-carboxytryptophan residues;

$\alpha$ -Amino substituted amino acid residues including sarcosine (N-methylglycine), N-benzylglycine, N-methylalanine, N-benzylalanine, N-methylphenylalanine, N-benzylphenylalanine, N-methylvaline and N-benzylvaline; and  
30

$\alpha$ -Hydroxy and substituted  $\alpha$ -hydroxy amino acid residues including serine, threonine, allothreonine, phosphoserine and phosphothreonine residues.

Any one of the foregoing or other known amino acids are suitably  
35 employed in this invention provided that they are capable of autocatalytically



hydrolyzing the amidate bond. Thus, they must contain, or must, upon being converted (hydrolyzed) in vivo, contain a free carboxyl group. In general, the amino acids corresponding to the residues employed in the compounds of this invention are naturally occurring and have no pharmacological activity.

5 However, optimal pharmacokinetic activity may be achieved by the use of non-naturally occurring amino acid residues.

Of particular interest are hydrophobic residues such as mono- or di-alkyl or aryl amino acids, cycloalkylamino acids and the like. These residues, together with R<sup>4</sup>, contribute to cell permeability by increasing the partition  
10 coefficient of the nucleotide analog amidate. Typically, the residue does not contain a sulfhydryl or guanidino substituent.


Polypeptide Radicals. If n<sub>1</sub> is greater than 1, then the group shown in formula II, IIa, IIb or III is greater than 1, then the moiety comprises a  
15 polypeptide radical. This comprises dipeptides, short polypeptides of 3, 5 or 10 residues, or proteins having up to 100 or more residues. For the most part, dipeptides not containing aspartic or glutamic acid in the residue adjacent to the P atom, will not autocatalytically hydrolyze the amidate bond and therefore the carboxyl groups (generally 1 or 2) in the distal residue do not  
20 need to be esterified or amidated, i.e., R<sup>4</sup> can be H in these circumstances. However, if such compounds are intended to be used as precursors for the free phosphonate nucleotide analog in vivo, rather than as immunogens for example, the polypeptides ordinarily will contain a peptidolytic enzyme cleavage site at the peptide bond linking the first residue and the next residue  
25 distal to the phosphorus atom. Such cleavage sites are flanked by enzymatic recognition structures, e.g. particular residues recognized by a hydrolytic enzyme.

Peptidolytic enzymes are well known, and in particular include carboxypeptidases. Carboxypeptidases digest polypeptides by removing C-  
30 terminal residues, and are specific in many instances for particular C-terminal sequences. Such enzymes and their substrate requirements in general are well known. For example, a dipeptide having a given pair of residues and a free carboxyl terminus is covalently bonded through its  $\alpha$ -amino group to the phosphorus atom of the invention nucleotide analogs. It is expected that this  
35 peptide will be cleaved by the appropriate dipeptidase or protease, leaving the

carboxyl of the proximal amino acid residue to autocatalytically cleave the amidate bond.

Examples of suitable dipeptidyl groups (designated by their single letter code) include AA, AR, AN, AD, AC, AE, AQ, AG, AH, AI, AL, AK, AM, AF, AP, AS, AT, AW, AY, AV, RA, RR, RN, RD, RC, RE, RQ, RG, RH, RI, RL, RK, RM, RF, RP, RS, RT, RW, RY, RV, NA, NR, NN, ND, NC, NE, NQ, NG, NH, NI, NL, NK, NM, NF, NP, NS, NT, NW, NY, NV, DA, DR, DN, DD, DC, DE, DQ, DG, DH, DI, DL, DK, DM, DF, DP, DS, DT, DW, DY, DV, CA, CR, CN, CD, CC, CE, CQ, CG, CH, CI, CL, CK, CM, CF, CP, CS, CT, CW, CY, CV, EA, ER, EN, ED, EC, EE, EQ, EG, EH, EI, EL, EK, EM, EF, EP, ES, ET, EW, EY, EV, QA, QR, QN, QD, QC, QE, QQ, QG, QH, QI, QL, QK, QM, QF, QP, QS, QT, QW, QY, QV, GA, GR, GN, GD, GC, GE, GQ, GG, GH, GI, GL, GK, GM, GF, GP, GS, GT, GW, GY, GV, HA, HR, HN, HD, HC, HE, HQ, HG, HH, HI, HL, HK, HM, HF, HP, HS, HT, HW, HY, HV, IA, IR, IN, ID, IC, IE, IQ, IG, IH, II, IL, IK, IM, IF, IP, IS, IT, IW, IY, IV, LA, LR, LN, LD, LC, LE, LQ, LG, LH, LI, LL, LK, LM, LF, LP, LS, LT, LW, LY, LV, KA, KR, KN, KD, KC, KE, KQ, KG, KH, KI, KL, KK, KM, KF, KP, KS, KT, KW, KY, KV, MA, MR, MN, MD, MC, ME, MQ, MG, MH, MI, ML, MK, MM, MF, MP, MS, MT, MW, MY, MV, FA, FR, FN, FD, FC, FE, FQ, FG, FH, FI, FL, FK, FM, FF, FP, FS, FT, FW, FY, FV, PA, PR, PN, PD, PC, PE, PQ, PG, PH, PI, PL, PK, PM, PF, PP, PS, PT, PW, PY, PV, SA, SR, SN, SD, SC, SE, SQ, SG, SH, SI, SL, SK, SM, SF, SP, SS, ST, SW, SY, SV, TA, TR, TN, TD, TC, TE, TQ, TG, TH, TI, TL, TK, TM, TF, TP, TS, TT, TW, TY, TV, WA, WR, WN, WD, WC, WE, WQ, WG, WH, WI, WL, WK, WM, WF, WP, WS, WT, WW, WY, WV, YA, YR, YN, YD, YC, YE, YQ, YG, YH, YI, YL, YK, YM, YF, YP, YS, YT, YW, YY, YV, VA, VR, VN, VD, VC, VE, VQ, VG, VH, VI, VL, VK, VM, VF, VP, VS, VT, VW, VY and VV.

Exemplary dipeptidyl compounds have the structure of formula IX wherein  $R^2$  is H,  $R^3$  is the side chain of a naturally occurring amino acid,  $L^1$ ,  $R^4$ , B and Z are as defined above.




Tripeptides are also useful. The sequence -X4-pro-X5- (where X4 is any amino acid residue and X5 is an amino acid residue, a carboxyl ester of proline or hydrogen) will be cleaved by luminal carboxypeptidase to yield X4 with a free carboxyl, which in turn autocatalytically cleaves the phosphono amidate bond. X5 usually will be a benzyl ester of the carboxy group of X5. Thus, n1 is usually 1, 2 or 3, but may range up to 5, 10, 100 or more residues.

If the amino acid residue has 2 or more amine groups, e.g., a lysinyl or arginyl, or ornithinyl residue, then R<sup>3</sup> represents the group -[C(R<sup>6</sup>)<sub>2</sub>]<sub>n2</sub>N(R<sup>2</sup>)- where n2 is 0 to 6, R<sup>6</sup> is H, C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>6</sub>-C<sub>20</sub> aryl, C<sub>7</sub>-C<sub>20</sub> alkylaryl, C<sub>7</sub>-C<sub>20</sub> arylalkyl, C<sub>1</sub>-C<sub>20</sub> alkoxy, C<sub>6</sub>-C<sub>20</sub> aryloxy or hydroxyl, and R<sup>2</sup> is defined above. Such compounds will contain a plurality of phosphonate moieties. For example when both the epsilon (ε)/delta (δ) and alpha (α) amino groups of lysine or ornithine are substituted with nucleotide phosphonate moieties the amidate is believed to be capable of releasing two molecules of active drug, each expected to emerge under different pharmacokinetics and therefore further sustaining the drug release.

The number of amino acid residues, n1, in the nucleotide analog amidates of this invention can vary extensively. Where n1=1, a single amino acid is found at the designated site, and where n1>1 then a polypeptide radical is present. Typically, n1 is 1 or 2, but may range up to 3, 5, 10 or 100 or more residues.

If the residue is immediately adjacent to the phosphonate atom and its side chain contains a carboxyl group, e.g. in the case of glutamic acid or aspartic acid, then this carboxylate is substituted with R<sup>4</sup>.

The amidate group optionally is taken together with Z to form a cyclic amidate precursor. Such compounds have structure XIV.





wherein  $L^2$ ,  $R^1$ ,  $R^2$ ,  $R^3$ ,  $Z$ ,  $n_1$  and  $B$  are as defined above. Typically, in this embodiment  $R^3$  is not carboxyl,  $R^2$  is H, and  $n_1$  is 1.



5 Hydrolysis of the cyclic amidates of formulas IIa-c and IV leaves a hydroxyl-substituted substructure  $Z$  and the free carboxyl, which in turn will autolyze the amidate. Substructures  $Z$  in which the methylene backbone is substituted with hydroxymethyl are advantageous in this embodiment, particularly linkers in compounds of the formula  $-CH_2OCH(CH_2O-)CH_2-B$ .

10

Heterocyclic bases. The compounds of this invention comprise any naturally-occurring heterocycle found in nucleic acids, nucleotides or nucleosides, or analogs thereof. The radicals of such heterocyclic bases, designated herein as  $B$ , are generally the purine, pyrimidine or related

15 heterocycles shown in formulas X-XIII.





wherein  $R^{15}$  is H, OH, F, Cl, Br, I,  $OR^{16}$ , SH,  $SR^{16}$ ,  $NH_2$ , or  $NHR^{17}$ ;

$R^{16}$  is  $C_1 - C_6$  alkyl including  $CH_3$ ,  $CH_2CH_3$ ,  $CH_2CCH$  (2-propynyl),  
 5  $CH_2CHCH_2$  (2-allyl),  $C_3H_7$ ;

$R^{17}$  is  $C_1 - C_6$  alkyl including  $CH_3$ ,  $CH_2CH_3$ ,  $CH_2CCH$ ,  $CH_2CHCH_2$ ,  
 $C_3H_7$ ;

$R^{18}$  is N, CF, CCl, CBr, Cl,  $CR^{19}$  or  $CSR^{19}$ ,  $COR^{19}$ ;

$R^{19}$  is H,  $C_1 - C_9$  alkyl,  $C_2 - C_9$  alkenyl,  $C_2 - C_9$  alkynyl or  $C_7 - C_9$  aryl-  
 10 alkyl unsubstituted or substituted by OH, O, N, F, Cl, Br or I including  $CH_3$ ,  
 $CH_2CH_3$ ,  $CHCH_2$ ,  $CHCHBr$ ,  $CH_2CH_2Cl$ ,  $CH_2CH_2F$ ,  $CH_2CCH$ ,  $CH_2CHCH_2$ ,  
 $C_3H_7$ ,  $CH_2OH$ ,  $CH_2OCH_3$ ,  $CH_2OC_2H_5$ ,  $CH_2OCCH$ ,  $CH_2OCH_2CHCH_2$ ,  
 $CH_2C_3H_7$ ,  $CH_2CH_2OH$ ,  $CH_2CH_2OCH_3$ ,  $CH_2CH_2OC_2H_5$ ,  $CH_2CH_2OCCH$ ,  
 $CH_2CH_2OCH_2CHCH_2$ ,  $CH_2CH_2OC_3H_7$ ;

15  $R^{20}$  is N or CH;

$R^{21}$  is N, CH, CCN,  $CCF_3$ ,  $CC\equiv CH$  or  $CC(O)NH_2$ ;

$R^{22}$  is H, OH,  $NH_2$ , SH,  $SCH_3$ ,  $SCH_2CH_3$ ,  $SCH_2CCH$ ,  $SCH_2CHCH_2$ ,  
 $SC_3H_7$ ,  $NH(CH_3)$ ,  $N(CH_3)_2$ ,  $NH(CH_2CH_3)$ ,  $N(CH_2CH_3)_2$ ,  $NH(CH_2CCH)$ ,  
 $NH(CH_2CHCH_2)$ ,  $NH(C_3H_7)$  or halogen (F, Cl, Br or I);


20  $R^{23}$  is H, OH, F, Cl, Br, I,  $SCH_3$ ,  $SCH_2CH_3$ ,  $SCH_2CCH$ ,  $SCH_2CHCH_2$ ,  
 $SC_3H_7$ ,  $OR^{16}$ ,  $NH_2$ , or  $NHR^{17}$ ; and

$R^{24}$  is O, S or Se.

B includes both protected and unprotected forms of the heterocyclic  
 bases. Protecting groups for exocyclic amines and other groups are known  
 25 (Greene and include N-benzoyl, isobutyryl, 4,4'-dimethoxytrityl (DMT) and  
 the like. The selection of a protecting group will be apparent to the ordinary  
 artisan and will depend on the nature of the labile group and the chemistry

which the protecting group is expected to encounter, e.g., acidic, basic, oxidative, reductive or other conditions.

As used herein, B<sup>1</sup> is a protected heterocyclic base having the formula Xa, XIa, XIb, XIIa or XIIIa



- 10 wherein R<sup>18</sup>, R<sup>20</sup>, R<sup>21</sup>, R<sup>24</sup> have the meanings previously defined; R<sup>22A</sup> is R<sup>39</sup> or R<sup>22</sup> provided that R<sup>22</sup> is not NH<sub>2</sub>; R<sup>23A</sup> is R<sup>39</sup> or R<sup>23</sup> provided that R<sup>23</sup> is not NH<sub>2</sub>; R<sup>39</sup> is NHR<sup>40</sup>, NHC(O)R<sup>36</sup> or NCR<sup>41</sup>N(R<sup>38</sup>)<sub>2</sub> wherein R<sup>36</sup> is C<sub>1</sub>-C<sub>19</sub> alkyl, C<sub>1</sub>-C<sub>19</sub> alkenyl, C<sub>3</sub>-C<sub>10</sub> aryl, adamantoyl, alkylanyl, or C<sub>3</sub>-C<sub>10</sub> aryl unsubstituted or substituted with 1 or 2 atoms or groups selected from
- 15 halogen, methyl, ethyl, methoxy, ethoxy, hydroxy and cyano; R<sup>38</sup> is C<sub>1</sub>-C<sub>10</sub> alkyl, or both R<sup>38</sup> together are 1-morpholino, 1-piperidine or 1-pyrrolidine; and R<sup>41</sup> is hydrogen or CH<sub>3</sub>. For heterocyclic bases of structures XIa and XIb, if R<sup>39</sup> is present at R<sup>22A</sup> or R<sup>23A</sup>, both R<sup>39</sup> groups on the same heterocyclic base will generally be the same. Exemplary R<sup>40</sup> include methyl, ethyl, propyl,
- 20 isopropyl, butyl, isobutyl, t-butyl, pentyl, hexyl, octyl, decanyl, lauryl and hexadecyl).


Specific heterocyclic bases include hypoxanthine, inosine, thymine, uracil, xanthine, 8-aza derivatives of 2-aminopurine, 2,6-diaminopurine, 2-

amino-6-chloropurine, hypoxanthine, inosine and xanthine; 7-deaza-8-aza derivatives of adenine, guanine, 2-aminopurine, 2,6-diaminopurine, 2-amino-6-chloropurine, hypoxanthine, inosine and xanthine; 1-deaza derivatives of 2-aminopurine, 2,6-diaminopurine, 2-amino-6-chloropurine, hypoxanthine, inosine and xanthine; 7-deaza derivatives of 2-aminopurine, 2,6-diaminopurine, 2-amino-6-chloropurine, hypoxanthine, inosine and xanthine; 3-deaza derivatives of 2-aminopurine, 2,6-diaminopurine, 2-amino-6-chloropurine, hypoxanthine, inosine and xanthine; 6-azacytosine; 5-fluorocytosine; 5-chlorocytosine; 5-iodocytosine; 5-bromocytosine; 5-methylcytosine; 5-bromovinyluracil; 5-fluorouracil; 5-chlorouracil; 5-iodouracil; 5-bromouracil; 5-trifluoromethyluracil; 5-methoxymethyluracil; 5-ethynyluracil; 5-propynyluracil and the like.

Preferably, B is a 9-purinylyl residue selected from guanylyl, 3-deazaguanylyl, 1-deazaguanylyl, 8-azaguanylyl, 7-deazaguanylyl, adenyl, 3-deazaadenyl, 1-dezazadenyl, 8-azaadenyl, 7-deazaadenyl, 2,6-diaminopurinylyl, 2-aminopurinylyl, 6-chloro-2-aminopurinylyl and 6-thio-2-aminopurinylyl, or a B is a 1-pyrimidinyl residue selected from cytosinyl, 5-halocytosinyl, and 5-(C<sub>1</sub>-C<sub>3</sub>-alkyl)cytosinyl.


The invention compounds, such as those of the formulas (L<sup>1</sup>)(RO)P(O)-Z-B, are optionally esterified at the phosphorus atom by the group R defined above. Exemplary R groups include phenyl, 2- and 3-pyrrolyl, 2- and 3-thienyl, 2- and 4-imidazolyl, 2-, 4- and 5-oxazolyl, 3- and 4-isoxazolyl, 2-, 4- and 5-thiazolyl, 3-, 4- and 5-isothiazolyl, 3- and 4-pyrazolyl, 2-, 3- and 4-pyridinyl, 2-, 4- and 5-pyrimidinyl, 2-, 3- and 4-alkoxyphenyl (C<sub>1</sub>-C<sub>12</sub> alkyl including 2-, 3- and 4-methoxyphenyl and 2-, 3- and 4-ethoxyphenyl), 2-, 3- and 4-halophenyl (including 2-, 3- and 4-fluorophenyl), 2,3-, 2,4-, 2,5-, 2,6-, 3,4- and 3,5-dihalophenyl (including 2,4-difluorophenyl and 2,4-dichlorophenyl), 2-, 3- and 4-haloalkylphenyl (1 to 5 halogen atoms, C<sub>1</sub>-C<sub>12</sub> alkyl including 2-, 3- and 4-trifluoromethylphenyl and 2-, 3- and 4-trichloromethylphenyl), 2-, 3- and 4-cyanophenyl, carboalkoxyphenyl (C<sub>1</sub>-C<sub>4</sub> alkyl including 2-, 3- and 4-carboethoxyphenyl (-C<sub>6</sub>H<sub>4</sub>-C(O)-OC<sub>2</sub>H<sub>5</sub>) and 2,3-, 2,4-, 2,5-, 2,6-, 3,4- and 3,5-dicarboethoxyphenyl), 1-, 2-, 3-, and 4-pyridinyl (-C<sub>5</sub>H<sub>4</sub>N), 2-, 3- and 4-nitrophenyl, 2-, 3- and 4-haloalkylbenzyl (1 to 5 halogen atoms, C<sub>1</sub>-C<sub>12</sub> alkyl including 4-trifluoromethylbenzyl), alkylsalicylphenyl (C<sub>1</sub>-C<sub>4</sub> alkyl including 2-, 3- and 4-ethylsalicylphenyl), 2,3- and 4-acetylphenyl, 1,8-dihydroxy-

naphthyl (-O-C<sub>10</sub>H<sub>6</sub>-OH or -O-C<sub>10</sub>H<sub>6</sub>-O-), 2,2'-dihydroxybiphenyl (-O-C<sub>6</sub>H<sub>4</sub>-C<sub>6</sub>H<sub>4</sub>-O-; both oxygen atoms are linked to the phosphorus atom), alkoxy ethyl [C<sub>1</sub>-C<sub>6</sub> alkyl including -CH<sub>2</sub>-CH<sub>2</sub>-O-CH<sub>3</sub> (methoxy ethyl) and phenoxymethyl], aryloxy ethyl [C<sub>6</sub>-C<sub>9</sub> aryl (including phenoxy ethyl) or C<sub>6</sub>-C<sub>9</sub> aryl substituted by OH, NH<sub>2</sub>, halo, C<sub>1</sub>-C<sub>4</sub> alkyl or C<sub>1</sub>-C<sub>4</sub> alkyl substituted by OH or by 1 to 3 halo atoms], -C<sub>6</sub>H<sub>4</sub>-CH<sub>2</sub>-N(CH<sub>3</sub>)<sub>2</sub>, N-ethylmorpholino



( ; -(CH<sub>2</sub>)<sub>2</sub>-N[(CH<sub>2</sub>)<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>O],

adamantoyl oxymethyl, pivaloyloxy(methoxyethyl)methyl (-CH(CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>)-O-C(O)-C(CH<sub>3</sub>)<sub>3</sub>),




( ; -O-CH<sub>2</sub>-O-C(O)-C<sub>10</sub>H<sub>15</sub>),

pivaloyloxymethyl (-CH<sub>2</sub>-O-C(O)-C(CH<sub>3</sub>)<sub>3</sub>), pivaloyloxy(methoxymethyl)-methyl (-CH(CH<sub>2</sub>OCH<sub>3</sub>)-O-C(O)-C(CH<sub>3</sub>)<sub>3</sub>), pivaloyloxyisobutyl (-CH(CH(CH<sub>3</sub>)<sub>2</sub>)-O-C(O)-C(CH<sub>3</sub>)<sub>3</sub>) isobutyryloxymethyl (-CH<sub>2</sub>-O-C(O)-CH<sub>2</sub>-CH(CH<sub>3</sub>)<sub>2</sub>), cyclohexanoyl oxymethyl (-CH<sub>2</sub>-O-C(O)-C<sub>6</sub>H<sub>11</sub>), phenyl (-C<sub>6</sub>H<sub>5</sub>), benzyl (-CH<sub>2</sub>-C<sub>6</sub>H<sub>5</sub>), isopropyl (-CH(CH<sub>3</sub>)<sub>2</sub>), t-butyl (-C(CH<sub>3</sub>)<sub>3</sub>), -CH<sub>2</sub>-CH<sub>3</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CH<sub>3</sub>, -(CH<sub>2</sub>)<sub>3</sub>-CH<sub>3</sub>, -(CH<sub>2</sub>)<sub>4</sub>-CH<sub>3</sub>, -(CH<sub>2</sub>)<sub>5</sub>-CH<sub>3</sub>, -CH<sub>2</sub>-CH<sub>2</sub>F, -CH<sub>2</sub>-CH<sub>2</sub>Cl, -CH<sub>2</sub>-CF<sub>3</sub>, -CH<sub>2</sub>-CCl<sub>3</sub>, R<sup>5</sup>, NHR<sup>6A</sup> or N(R<sup>6A</sup>)<sub>2</sub> wherein R<sup>5</sup> is CH<sub>2</sub>C(O)N(R<sup>6A</sup>)<sub>2</sub>, CH<sub>2</sub>C(O)OR<sup>6A</sup>, CH<sub>2</sub>OC(O)R<sup>6A</sup>, CH(R<sup>6A</sup>)OC(O)R<sup>6A</sup>, CH<sub>2</sub>C(R<sup>6A</sup>)<sub>2</sub>CH<sub>2</sub>OH, CH<sub>2</sub>OR<sup>6A</sup>, NH-CH<sub>2</sub>-C(O)O-CH<sub>2</sub>CH<sub>3</sub>, N(CH<sub>3</sub>)-CH<sub>2</sub>-C(O)O-CH<sub>2</sub>CH<sub>3</sub>, NHR<sup>40</sup>, CH<sub>2</sub>-O-C(O)-C<sub>6</sub>H<sub>5</sub>, CH<sub>2</sub>-O-C(O)-C<sub>10</sub>H<sub>15</sub>, -CH<sub>2</sub>-O-C(O)-CH<sub>2</sub>CH<sub>3</sub>, CH<sub>2</sub>-O-C(O)-CH(CH<sub>3</sub>)<sub>2</sub>, CH<sub>2</sub>-O-C(O)-C(CH<sub>3</sub>)<sub>3</sub>, CH<sub>2</sub>-O-C(O)-CH<sub>2</sub>-C<sub>6</sub>H<sub>5</sub>, wherein R<sup>6A</sup> is C<sub>1</sub>-C<sub>20</sub> alkyl which is unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N and halogen (1 to 5 halogen atoms), C<sub>6</sub>-C<sub>20</sub> aryl which is unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N and halogen (1 to 5 halogen atoms) or C<sub>7</sub>-C<sub>20</sub> aryl-alkyl which is unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N and halogen (1 to 5 halogen atoms), provided that for compounds of formulas N(R<sup>6A</sup>)<sub>2</sub>, CH<sub>2</sub>C(O)N(R<sup>6A</sup>)<sub>2</sub>, CH<sub>2</sub>C(O)OR<sup>6A</sup>, CH<sub>2</sub>OC(O)R<sup>6A</sup>, CH(R<sup>6A</sup>)OC(O)R<sup>6A</sup> and CH<sub>2</sub>C(R<sup>6A</sup>)<sub>2</sub>CH<sub>2</sub>OH, the total




number of carbon atoms present is less than 25 (preferably the number of carbon atoms present is about 4 to about 14) and  $R^{40}$  is  $C_1$ - $C_{20}$  alkyl.

The invention compounds are optionally alkylated at the  $\alpha$ -nitrogen atom of the amino acid by the  $R^1$  group defined above. Exemplary  $R^1$  groups include H,  $CH_3$ ,  $CH_2CH_3$ , benzyl, 4-O-N-methylpiperidinyl



( $-O-CH[(CH_2)_2(CH_2)_2]N(CH_3)$ ), 3-O-N-methylpiperidinyl and the like.

The invention compounds are optionally esterified at the amino acid carboxyl moiety by the  $R^4$  group defined above. Exemplary  $R^4$  groups include H, methyl, ethyl, propyl, isopropyl, butyl, t-butyl ( $C(CH_3)_3$ ), phenyl ( $-C_6H_5$ ), benzyl ( $-CH_2-C_6H_5$ ), 1-pyridyl, 3-pyridyl, 1-pyrimidinyl, N-ethylmorpholino ( $-CH_2-CH_2-N[(CH_2)_2(CH_2)_2]O$ ), N-2-propylmorpholino ( $-CH(CH_3)-CH_2-N[(CH_2)_2(CH_2)_2]O$ ), methoxyethyl ( $-CH_2-CH_2-O-CH_3$ ), 4-N-methylpiperidyl ( $-CH[(CH_2)_2(CH_2)_2]N(CH_3)$ ), 3-N-methylpiperidyl, phenol which is 2-, 3-, or 4-substituted by  $N(R^{30})_2$  where  $R^{30}$  is independently H or  $C_1$ - $C_6$  alkyl unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N,  $COOR^4$  and halogen or  $C_6$ - $C_{12}$  aryl unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N,  $COOR^4$ ,  $N(R^7)_2$  and halogen (including 2-, 3-, and 4-N,N-dimethylaminophenol and 2-, 3-, and 4-N,N-diethylamino-phenol), 1-ethylpiperazinyl



[ $-CH_2-CH_2-NC_4H_8NH$ ], and N<sup>4</sup>-substituted 1-ethylpiperazinyl ( $-(CH_2)_2-N[(CH_2)_2(CH_2)_2]NR^2$ , where  $R^2$  is as defined above).

Additional compounds that are included in the invention are nucleotide analog dimers that are linked via an amino or carboxyl group. As used herein, dimers (or trimers) refer to the presence of two (or three) nucleoside residues that comprise a compound. Thus, a  $-L^1-P(O)(L^1)-Z-B$  or  $-P(O)(L^1)-Z-B$  radical covalently linked to a  $-L^1-P(O)(L^1)-Z-B$  or  $-P(O)(L^1)-Z-B$  radical gives  $B-Z-P(O)(L^1)-P(O)(L^1)-Z-B$ ,  $B-Z-P(O)(L^1)-L^1-P(O)(L^1)-Z-B$  or  $B-Z-P(O)(L^1)-L^1-L^1-P(O)(L^1)-Z-B$ .

Dimer nucleotide analogs are conveniently linked via amino acids, diamino acids, dicarboxylic amino acids, diamines or dicarboxylic acids such as  $\beta$ -aminoalanine, diaminobutyric acid, citrulline, homoarginine, homocitrulline, ornithine,  $\gamma$ -aminobutyric acid, arginine, histidine, asparagine, glutamine,  $\beta$ -hydroxyaspartic acid,  $\beta$ -hydroxyglutamic acid,  $\beta$ -methylaspartic acid,  $\beta$ -methylglutamic acid, 3-aminoadipic acid, 2-aminopimelic acid, 2-aminosuberic acid,  $\beta$ -amino acid analogs of lysine ( $\text{NH}_2\text{-(CH}_2\text{)}_3\text{-CH(NH}_2\text{)-CH}_2\text{-CH-C(O)OH}$ ), arginine, histidine, asparagine, glutamine and the like. Exemplary compounds include dimers linked via lysine or  $\beta$ -lysine having the formulas  $\text{B-Z-P(O)(L)-NH-(CH}_2\text{)}_4\text{-CH(C(O)OR}^4\text{)-NR}^1\text{-P(O)(L)-Z-B}$  and  $\text{B-Z-P(O)(L)-NH-(CH}_2\text{)}_3\text{-CH(CH}_2\text{C(O)OR}^4\text{)-NR}^1\text{-P(O)(L)-Z-B}$  and dimers linked via aspartic or glutamic acid having the formula  $\text{B-Z-P(O)(L)-O-C(O)-(CH}_2\text{)}_{1-2}\text{-CH(C(O)OR}^4\text{)-NR}^1\text{-P(O)(L)-Z-B}$ . L, Z and B are independently selected.

Nucleotide analogs comprising dipeptidyl or tripeptidyl L groups are also included in the compounds of the invention. Nucleotide radicals are linked through side chain groups (usually amino or carboxyl) or through amino and carboxyl groups of the amino acids. Exemplary dipeptidyl and tripeptidyl dimers and trimers include compounds of the formulas  $\text{B-Z-P(O)(L}^1\text{)-O-C(O)-(CR}^2\text{R}^3\text{)}_n\text{-NR}^1\text{-C(O)-(CR}^2\text{R}^3\text{)}_n\text{-NR}^1\text{-P(O)(L}^1\text{)-Z-B}$ ,  $\text{B-Z-P(O)(L}^1\text{)-O-C(O)-(CR}^2\text{R}^3\text{)}_n\text{-NR}^1\text{-C(O)-(CR}^2\text{R}^3\text{)}_n\text{-NR}^1\text{-O-C(O)-(CR}^2\text{R}^3\text{)}_n\text{-NR}^1\text{-P(O)(L}^1\text{)-Z-B}$ ,  $\text{B-Z-P(O)(L}^1\text{)-O-C(O)-CR}^2\text{(R}^3\text{-P(O)(L}^1\text{)-Z-B)-NR}^1\text{-C(O)-(CR}^2\text{R}^3\text{)}_n\text{-NR}^1\text{-P(O)(L}^1\text{)-Z-B}$  and  $\text{B-Z-P(O)(L}^1\text{)-O-C(O)-(CR}^2\text{R}^3\text{)}_n\text{-NR}^1\text{-C(O)-CR}^2\text{(R}^3\text{-P(O)(L}^1\text{)-Z-B)-NR}^1\text{-P(O)(L}^1\text{)-Z-B}$ . In order to provide a compound with a desired molar ratio of one Z-B compared to a second Z-B, tetramer, pentamer and higher polymer forms can also be prepared where Z and/or B are independently chosen.

As used herein, and unless modified by the immediate context: 1) the term alkyl, alkenyl and alkynyl refer to straight chain, branched and cyclic residues. Thus,  $\text{C}_1\text{-C}_4$  alkyl includes methyl, ethyl, propyl, cyclopropyl, isopropyl, n-, sec-, iso- and tert-butyl, cyclobutyl and the like while alkenyl includes ethenyl, propenyl, isopropenyl, 1-, 2- and 3-butenyl, 1- and 2-isobutenyl and the like. The term alkyl also includes cyclic N-, S- or O-heterocarbonyl (such as piperidyl and morpholino). 2) The term aryl includes N-, S- or O- heteroaryl, including phenyl, 2- and 3-pyrrolyl, 2- and 3-thienyl, 2-

and 4-imidazolyl, 2-, 4- and 5-oxazolyl, 3- and 4-isoxazolyl, 2-, 4- and 5-thiazolyl, 3-, 4- and 5-isothiazolyl, 3- and 4-pyrazolyl, 2-, 3- and 4-pyridinyl, 2-, 4- and 5-pyrimidinyl. When "O" or "N" are substituted into aryl or alkyl this means that a ring or chain methyne or methylene is replaced by O, N or NH as the case may be. The term acyl means  $R^X-C(O)-$ , acyloxy means  $R^X-C(O)-O-$ , acyloxymethyl means  $R^X-C(O)-O-CH_2-$  and thus, for example,  $C_{3-6}$  acyloxymethyl means  $R^X-C(O)-O-CH_2-$  wherein  $R^X$  is a 1 to 4 carbon alkyl or aryl group (substituted or unsubstituted).

- 10        Nucleoside Phosphonates. Table 1 lists a group of exemplary nucleotide analogs of formula I having the structure  $(L^1)(L^2)P(O)-Z-B$ . These compounds generally have  $L^1$  and  $L^2$  groups that, when amino acids, are identical, although one of the amino acid groups can be different or replaced by another hydrolyzable group such as  $-O-CH_2-O-C(O)-C(CH_3)_3$  or  $-O-C_6H_5$  as listed below.

TABLE 1

$L^1, L^2^*$	$-Z-B^{**}$
1 -NH-CH <sub>2</sub> -C(O)-OR <sup>4</sup>	1 -CH <sub>2</sub> -O-CH <sub>2</sub> -CH <sub>2</sub> -B
20 2 -NH-CH(CH <sub>3</sub> )-C(O)-OR <sup>4</sup>	2 -CH <sub>2</sub> -O-C <sup>#</sup> H(CH <sub>2</sub> -OR <sup>4</sup> )-CH <sub>2</sub> -B
3 -NH-CH(CH <sub>3</sub> ) <sub>2</sub> -C(O)-OR <sup>4</sup>	3 -CH <sub>2</sub> -O-C <sup>#</sup> H(CH <sub>3</sub> )-CH <sub>2</sub> -B
4 -NH-CH(CH(CH <sub>3</sub> ) <sub>2</sub> )-C(O)-OR <sup>4</sup>	4 -CH <sub>2</sub> -O-C <sup>#</sup> H(CH <sub>2</sub> F)-CH <sub>2</sub> -B
5 -NH-CH(CH <sub>3</sub> )(CH <sub>3</sub> ) <sub>2</sub> -C(O)-OR <sup>4</sup>	5 -CH <sub>2</sub> -O-C <sup>#</sup> H(CH=CH <sub>2</sub> )-CH <sub>2</sub> -B
6 -NH-CH <sub>2</sub> -CH <sub>2</sub> -CH <sub>2</sub> -CH-C(O)-OR <sup>4</sup>	6 -CH <sub>2</sub> -O-C <sup>#</sup> H(CH <sub>2</sub> N <sub>3</sub> )-CH <sub>2</sub> -B
25 7 -NH-CH(CH <sub>2</sub> -C <sub>6</sub> H <sub>5</sub> )-C(O)-OR <sup>4</sup>	7 ***
8 -NH-CH(CH <sub>2</sub> -C <sub>8</sub> NH <sub>6</sub> )-C(O)-OR <sup>4</sup>	8 ****
9 -NH-CH(CH <sub>2</sub> -CH <sub>2</sub> -S-CH <sub>3</sub> )-C(O)-OR <sup>4</sup>	
10 -NH-CH(CH <sub>2</sub> OH)-C(O)-OR <sup>4</sup>	
11 -NH-CH(CH(OH)(CH <sub>3</sub> ))-C(O)-OR <sup>4</sup>	
30 12 -NH-CH(-CH <sub>2</sub> SH)-C(O)-OR <sup>4</sup>	
13 -NH-CH(CH <sub>2</sub> -C <sub>6</sub> H <sub>5</sub> OH)-C(O)-OR <sup>4</sup>	
14 -NH-CH(CH <sub>2</sub> -C(O)-NH <sub>2</sub> )-C(O)-OR <sup>4</sup>	
15 -NH-CH(CH <sub>2</sub> -CH <sub>2</sub> -C(O)-NH <sub>2</sub> )-C(O)-OR <sup>4</sup>	
16 -NH-CH(CH <sub>2</sub> C(O)OR <sup>4</sup> )-C(O)-OR <sup>4</sup>	
35 17 -NH-CH(CH <sub>2</sub> CH <sub>2</sub> C(O)OR <sup>4</sup> )-C(O)-OR <sup>4</sup>	

- 18 -NH-CH(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>)-C(O)-OR<sup>4</sup>  
 19 -NH-CH(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NHC(NH)(NH<sub>2</sub>))-C(O)-OR<sup>4</sup>  
 20 -NH-CH(CH<sub>2</sub>C<sub>3</sub>N<sub>2</sub>H<sub>3</sub>)-C(O)-OR<sup>4</sup>  
 21 -NH-CH(CH<sub>3</sub>)<sub>2</sub>-CH<sub>2</sub>-C(O)-OR<sup>4</sup>  
 5 22 -NH-CH<sub>2</sub>-CH<sub>2</sub>-C(O)-OR<sup>4</sup>  
 23 -NH-CH(CH<sub>2</sub>-C<sub>6</sub>H<sub>5</sub>)-CH<sub>2</sub>-C(O)-OR<sup>4</sup>  
 24 -NH-CH(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>)-CH<sub>2</sub>-C(O)-OR<sup>4</sup>  
 25 -NH-CH(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>)-CH<sub>2</sub>-C(O)-OR<sup>4</sup>  
 26 -NH-CH(CH<sub>2</sub>CH<sub>2</sub>NHC(NH)(NH<sub>2</sub>))-CH<sub>2</sub>-C(O)-OR<sup>4</sup>  
 10 27 -NH-CH(C(O)OR<sup>4</sup>)-CH<sub>2</sub>-C(O)-OR<sup>4</sup>  
 28 -NH-CH(CH<sub>2</sub>C(O)OR<sup>4</sup>)-CH<sub>2</sub>-C(O)-OR<sup>4</sup>  
 29 -NH-CH(CH<sub>2</sub>CH<sub>2</sub>C(O)OR<sup>4</sup>)-CH<sub>2</sub>-C(O)-OR<sup>4</sup>  
 30 -N(CH<sub>3</sub>)-CH<sub>2</sub>-C(O)-OR<sup>4</sup>  
 31 -NHR<sup>6</sup>  
 15 32 -O-CH<sub>2</sub>-CH<sub>2</sub>-N[CH<sub>2</sub>]<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>O  
 33 -O-CH<sub>2</sub>-O-C(O)-C(CH<sub>3</sub>)<sub>3</sub>  
 34 -O-CH<sub>2</sub>-O-C(O)-CH(CH<sub>3</sub>)<sub>2</sub>  
 35 -O-CH<sub>2</sub>-O-C(O)-CH<sub>2</sub>C<sub>6</sub>H<sub>4</sub>-O-CH<sub>2</sub>CH<sub>3</sub>  
 36 -O-CH<sub>2</sub>-O-C(O)-C<sub>10</sub>H<sub>15</sub>  
 20 37 -O-CH<sub>2</sub>-C<sub>6</sub>H<sub>5</sub>  
 38 -O-C<sub>6</sub>H<sub>5</sub>  
 39 -O-CH<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>N(CH<sub>3</sub>)<sub>2</sub>  
 40 -OH
- 25 B
- 1 adenin-9-yl  
 2 guanin-9-yl  
 3 cytosin-1-yl  
 4 2, 6-diaminopurin-9-yl  
 30 5 2-aminopurin-9-yl  
 6 6-azacytosin-1-yl  
 7 1-deazaadenin-9-yl  
 8 3-deazaadenin-9-yl  
 9 8-azaadenin-9-yl  
 35 10 7-deaza-8-azaadenin-9-yl


\* - R<sup>4</sup> includes H, propyl, isopropyl, t-butyl, phenyl, benzyl, 1-pyridinyl, 1-pyrimidinyl, N-ethylmorpholino, methoxyethyl, 4-hydroxy-N-methylpiperidinyl, 3-hydroxy-N-methylpiperidinyl, 1-ethylpiperazinyl; atoms  
5 with unfilled valences are linked to each other.

\*\* - The carbon atom on the left of each structure is attached to the phosphorus atom; # - carbon atom having linked substituents in the *R*, *S* or *RS* configuration.


\*\*\* - Z-B substructure 7 is of formula V where R<sup>25</sup> and R<sup>29</sup> are O and B is  
10 thymine-1-yl (base 11) or one of the heterocyclic bases listed (1-10).

\*\*\*\* - Z-B substructure 8 is of formula IV where R<sup>25</sup> and R<sup>29</sup> are O, R<sup>26</sup> is S, R<sup>27</sup> is absent, R<sup>28</sup> is H and B is thymine-1-yl (base 11) or one of the heterocyclic bases listed (1-10) and includes the (+) and (-) enantiomers.

15 Compounds listed in Table 1 are designated herein by numbers assigned to L<sup>1</sup>, L<sup>2</sup>, Z and B according to the following convention, L<sup>1</sup>.L<sup>2</sup>.Z.B. Thus, compound 1.2.1.1, where R<sup>4</sup> is benzyl, represents L<sup>1</sup> structure 1 (-NH-CH<sub>2</sub>-C(O)-O-CH<sub>2</sub>-C<sub>6</sub>H<sub>5</sub>), L<sup>2</sup> structure 2 (-NH-CH(CH<sub>3</sub>)-C(O)-O-CH<sub>2</sub>-C<sub>6</sub>H<sub>5</sub>), Z structure 1 (-CH<sub>2</sub>-O-CH<sub>2</sub>-CH<sub>2</sub>-) and B structure 1 (adenine-9-yl). This  
20 compound would have the structure



which corresponds to the compound designated herein bis(alanyl benzyl ester)PMEA. Similarly, for the compound 7.7.1.1, L<sup>1</sup> structure 7 (NH-CH(CH<sub>2</sub>-C<sub>6</sub>H<sub>5</sub>)-C(O)-OR<sup>4</sup>), L<sup>2</sup> structure 7 (NH-CH(CH<sub>2</sub>-C<sub>6</sub>H<sub>5</sub>)-C(O)-OR<sup>4</sup>), Z structure 2 (-CH<sub>2</sub>-O-CH<sub>2</sub>-CH<sub>2</sub>-) and B structure 1 (adenine-9-yl) would have, when R<sup>4</sup> is  
25 methyl, the structure



and would represent the compound designated herein bis(phenylalanyl methyl ester)PMEA. Exemplary compounds include 1.1.1.1, 2.1.1.1, 3.1.1.1, 4.1.1.1, 5.1.1.1, 6.1.1.1, 7.1.1.1, 8.1.1.1, 9.1.1.1, 10.1.1.1, 11.1.1.1, 12.1.1.1, 13.1.1.1, 14.1.1.1, 15.1.1.1, 16.1.1.1, 17.1.1.1, 18.1.1.1, 19.1.1.1, 20.1.1.1, 21.1.1.1, 22.1.1.1, 23.1.1.1, 24.1.1.1, 25.1.1.1, 26.1.1.1, 27.1.1.1, 28.1.1.1, 29.1.1.1, 30.1.1.1, 31.1.1.1, 32.1.1.1, 33.1.1.1, 34.1.1.1, 35.1.1.1, 36.1.1.1, 37.1.1.1, 38.1.1.1, 39.1.1.1, 40.1.1.1, 1.2.1.1, 2.2.1.1, 3.2.1.1, 4.2.1.1, 5.2.1.1, 6.2.1.1, 7.2.1.1, 8.2.1.1, 9.2.1.1, 10.2.1.1, 11.2.1.1, 12.2.1.1, 13.2.1.1, 14.2.1.1, 15.2.1.1, 16.2.1.1, 17.2.1.1, 18.2.1.1, 19.2.1.1, 20.2.1.1, 21.2.1.1, 22.2.1.1, 23.2.1.1, 24.2.1.1, 25.2.1.1, 26.2.1.1, 27.2.1.1, 28.2.1.1, 29.2.1.1, 30.2.1.1, 31.2.1.1, 32.2.1.1, 33.2.1.1, 34.2.1.1, 35.2.1.1, 36.2.1.1, 37.2.1.1, 38.2.1.1, 39.2.1.1, 40.2.1.1, 1.3.1.1, 2.3.1.1, 3.3.1.1, 4.3.1.1, 5.3.1.1, 6.3.1.1, 7.3.1.1, 8.3.1.1, 9.3.1.1, 10.3.1.1, 11.3.1.1, 12.3.1.1, 13.3.1.1, 14.3.1.1, 15.3.1.1, 16.3.1.1, 17.3.1.1, 18.3.1.1, 19.3.1.1, 20.3.1.1, 21.3.1.1, 22.3.1.1, 23.3.1.1, 24.3.1.1, 25.3.1.1, 26.3.1.1, 27.3.1.1, 28.3.1.1, 29.3.1.1, 30.3.1.1, 31.3.1.1, 32.3.1.1, 33.3.1.1, 34.3.1.1, 35.3.1.1, 36.3.1.1, 37.3.1.1, 38.3.1.1, 39.3.1.1, 40.3.1.1, 1.4.1.1, 2.4.1.1, 3.4.1.1, 4.4.1.1, 5.4.1.1, 6.4.1.1, 7.4.1.1, 8.4.1.1, 9.4.1.1, 10.4.1.1, 11.4.1.1, 12.4.1.1, 13.4.1.1, 14.4.1.1, 15.4.1.1, 16.4.1.1, 17.4.1.1, 18.4.1.1, 19.4.1.1, 20.4.1.1, 21.4.1.1, 22.4.1.1, 23.4.1.1, 24.4.1.1, 25.4.1.1, 26.4.1.1, 27.4.1.1, 28.4.1.1, 29.4.1.1, 30.4.1.1, 31.4.1.1, 32.4.1.1, 33.4.1.1, 34.4.1.1, 35.4.1.1, 36.4.1.1, 37.4.1.1, 38.4.1.1, 39.4.1.1, 40.4.1.1, 1.5.1.1, 2.5.1.1, 3.5.1.1, 4.5.1.1, 5.5.1.1, 6.5.1.1, 7.5.1.1, 8.5.1.1, 9.5.1.1, 10.5.1.1, 11.5.1.1, 12.5.1.1, 13.5.1.1, 14.5.1.1, 15.5.1.1, 16.5.1.1, 17.5.1.1, 18.5.1.1, 19.5.1.1, 20.5.1.1, 21.5.1.1, 22.5.1.1, 23.5.1.1, 24.5.1.1, 25.5.1.1, 26.5.1.1, 27.5.1.1, 28.5.1.1, 29.5.1.1, 30.5.1.1, 31.5.1.1, 32.5.1.1, 33.5.1.1, 34.5.1.1, 35.5.1.1, 36.5.1.1, 37.5.1.1, 38.5.1.1, 39.5.1.1, 40.5.1.1, 1.6.1.1, 2.6.1.1, 3.6.1.1, 4.6.1.1, 5.6.1.1, 6.6.1.1, 7.6.1.1, 8.6.1.1, 9.6.1.1, 10.6.1.1, 11.6.1.1, 12.6.1.1, 13.6.1.1, 14.6.1.1, 15.6.1.1, 16.6.1.1, 17.6.1.1, 18.6.1.1, 19.6.1.1, 20.6.1.1, 21.6.1.1, 22.6.1.1, 23.6.1.1, 24.6.1.1, 25.6.1.1, 26.6.1.1, 27.6.1.1, 28.6.1.1, 29.6.1.1, 30.6.1.1, 31.6.1.1, 32.6.1.1, 33.6.1.1, 34.6.1.1, 35.6.1.1,

36.6.1.1, 37.6.1.1, 38.6.1.1, 39.6.1.1, 40.6.1.1, 1.7.1.1, 2.7.1.1, 3.7.1.1, 4.7.1.1, 5.7.1.1,  
 6.7.1.1, 7.7.1.1, 8.7.1.1, 9.7.1.1, 10.7.1.1, 11.7.1.1, 12.7.1.1, 13.7.1.1, 14.7.1.1, 15.7.1.1,  
 16.7.1.1, 17.7.1.1, 18.7.1.1, 19.7.1.1, 20.7.1.1, 21.7.1.1, 22.7.1.1, 23.7.1.1, 24.7.1.1,  
 25.7.1.1, 26.7.1.1, 27.7.1.1, 28.7.1.1, 29.7.1.1, 30.7.1.1, 31.7.1.1, 32.7.1.1, 33.7.1.1,  
 5 34.7.1.1, 35.7.1.1, 36.7.1.1, 37.7.1.1, 38.7.1.1, 39.7.1.1, 40.7.1.1, 1.8.1.1, 2.8.1.1, 3.8.1.1,  
 4.8.1.1, 5.8.1.1, 6.8.1.1, 7.8.1.1, 8.8.1.1, 9.8.1.1, 10.8.1.1, 11.8.1.1, 12.8.1.1, 13.8.1.1,  
 14.8.1.1, 15.8.1.1, 16.8.1.1, 17.8.1.1, 18.8.1.1, 19.8.1.1, 20.8.1.1, 21.8.1.1, 22.8.1.1,  
 23.8.1.1, 24.8.1.1, 25.8.1.1, 26.8.1.1, 27.8.1.1, 28.8.1.1, 29.8.1.1, 30.8.1.1, 31.8.1.1,  
 32.8.1.1, 33.8.1.1, 34.8.1.1, 35.8.1.1, 36.8.1.1, 37.8.1.1, 38.8.1.1, 39.8.1.1, 40.8.1.1,  
 10 1.9.1.1, 2.9.1.1, 3.9.1.1, 4.9.1.1, 5.9.1.1, 6.9.1.1, 7.9.1.1, 8.9.1.1, 9.9.1.1, 10.9.1.1,  
 11.9.1.1, 12.9.1.1, 13.9.1.1, 14.9.1.1, 15.9.1.1, 16.9.1.1, 17.9.1.1, 18.9.1.1, 19.9.1.1,  
 20.9.1.1, 21.9.1.1, 22.9.1.1, 23.9.1.1, 24.9.1.1, 25.9.1.1, 26.9.1.1, 27.9.1.1, 28.9.1.1,  
 29.9.1.1, 30.9.1.1, 31.9.1.1, 32.9.1.1, 33.9.1.1, 34.9.1.1, 35.9.1.1, 36.9.1.1, 37.9.1.1,  
 38.9.1.1, 39.9.1.1, 40.9.1.1, 1.10.1.1, 2.10.1.1, 3.10.1.1, 4.10.1.1, 5.10.1.1, 6.10.1.1,  
 15 7.10.1.1, 8.10.1.1, 9.10.1.1, 10.10.1.1, 11.10.1.1, 12.10.1.1, 13.10.1.1, 14.10.1.1,  
 15.10.1.1, 16.10.1.1, 17.10.1.1, 18.10.1.1, 19.10.1.1, 20.10.1.1, 21.10.1.1, 22.10.1.1,  
 23.10.1.1, 24.10.1.1, 25.10.1.1, 26.10.1.1, 27.10.1.1, 28.10.1.1, 29.10.1.1, 30.10.1.1,  
 31.10.1.1, 32.10.1.1, 33.10.1.1, 34.10.1.1, 35.10.1.1, 36.10.1.1, 37.10.1.1, 38.10.1.1,  
 39.10.1.1, 40.10.1.1, 1.11.1.1, 2.11.1.1, 3.11.1.1, 4.11.1.1, 5.11.1.1, 6.11.1.1, 7.11.1.1,  
 20 8.11.1.1, 9.11.1.1, 10.11.1.1, 11.11.1.1, 12.11.1.1, 13.11.1.1, 14.11.1.1, 15.11.1.1,  
 16.11.1.1, 17.11.1.1, 18.11.1.1, 19.11.1.1, 20.11.1.1, 21.11.1.1, 22.11.1.1, 23.11.1.1,  
 24.11.1.1, 25.11.1.1, 26.11.1.1, 27.11.1.1, 28.11.1.1, 29.11.1.1, 30.11.1.1, 31.11.1.1,  
 32.11.1.1, 33.11.1.1, 34.11.1.1, 35.11.1.1, 36.11.1.1, 37.11.1.1, 38.11.1.1, 39.11.1.1,  
 40.11.1.1, 1.12.1.1, 2.12.1.1, 3.12.1.1, 4.12.1.1, 5.12.1.1, 6.12.1.1, 7.12.1.1, 8.12.1.1,  
 25 9.12.1.1, 10.12.1.1, 11.12.1.1, 12.12.1.1, 13.12.1.1, 14.12.1.1, 15.12.1.1, 16.12.1.1,  
 17.12.1.1, 18.12.1.1, 19.12.1.1, 20.12.1.1, 21.12.1.1, 22.12.1.1, 23.12.1.1, 24.12.1.1,  
 25.12.1.1, 26.12.1.1, 27.12.1.1, 28.12.1.1, 29.12.1.1, 30.12.1.1, 31.12.1.1, 32.12.1.1,  
 33.12.1.1, 34.12.1.1, 35.12.1.1, 36.12.1.1, 37.12.1.1, 38.12.1.1, 39.12.1.1, 40.12.1.1,  
 1.13.1.1, 2.13.1.1, 3.13.1.1, 4.13.1.1, 5.13.1.1, 6.13.1.1, 7.13.1.1, 8.13.1.1, 9.13.1.1,  
 30 10.13.1.1, 11.13.1.1, 12.13.1.1, 13.13.1.1, 14.13.1.1, 15.13.1.1, 16.13.1.1, 17.13.1.1,  
 18.13.1.1, 19.13.1.1, 20.13.1.1, 21.13.1.1, 22.13.1.1, 23.13.1.1, 24.13.1.1, 25.13.1.1,  
 26.13.1.1, 27.13.1.1, 28.13.1.1, 29.13.1.1, 30.13.1.1, 31.13.1.1, 32.13.1.1, 33.13.1.1,  
 34.13.1.1, 35.13.1.1, 36.13.1.1, 37.13.1.1, 38.13.1.1, 39.13.1.1, 40.13.1.1, 1.14.1.1,  
 2.14.1.1, 3.14.1.1, 4.14.1.1, 5.14.1.1, 6.14.1.1, 7.14.1.1, 8.14.1.1, 9.14.1.1, 10.14.1.1,  
 35 11.14.1.1, 12.14.1.1, 13.14.1.1, 14.14.1.1, 15.14.1.1, 16.14.1.1, 17.14.1.1, 18.14.1.1,

19.14.1.1, 20.14.1.1, 21.14.1.1, 22.14.1.1, 23.14.1.1, 24.14.1.1, 25.14.1.1, 26.14.1.1,  
27.14.1.1, 28.14.1.1, 29.14.1.1, 30.14.1.1, 31.14.1.1, 32.14.1.1, 33.14.1.1, 34.14.1.1,  
35.14.1.1, 36.14.1.1, 37.14.1.1, 38.14.1.1, 39.14.1.1, 40.14.1.1, 1.15.1.1, 2.15.1.1,  
3.15.1.1, 4.15.1.1, 5.15.1.1, 6.15.1.1, 7.15.1.1, 8.15.1.1, 9.15.1.1, 10.15.1.1, 11.15.1.1,  
5 12.15.1.1, 13.15.1.1, 14.15.1.1, 15.15.1.1, 16.15.1.1, 17.15.1.1, 18.15.1.1, 19.15.1.1,  
20.15.1.1, 21.15.1.1, 22.15.1.1, 23.15.1.1, 24.15.1.1, 25.15.1.1, 26.15.1.1, 27.15.1.1,  
28.15.1.1, 29.15.1.1, 30.15.1.1, 31.15.1.1, 32.15.1.1, 33.15.1.1, 34.15.1.1, 35.15.1.1,  
36.15.1.1, 37.15.1.1, 38.15.1.1, 39.15.1.1, 40.15.1.1, 1.16.1.1, 2.16.1.1, 3.16.1.1, 4.16.1.1,  
5.16.1.1, 6.16.1.1, 7.16.1.1, 8.16.1.1, 9.16.1.1, 10.16.1.1, 11.16.1.1, 12.16.1.1, 13.16.1.1,  
10 14.16.1.1, 15.16.1.1, 16.16.1.1, 17.16.1.1, 18.16.1.1, 19.16.1.1, 20.16.1.1, 21.16.1.1,  
22.16.1.1, 23.16.1.1, 24.16.1.1, 25.16.1.1, 26.16.1.1, 27.16.1.1, 28.16.1.1, 29.16.1.1,  
30.16.1.1, 31.16.1.1, 32.16.1.1, 33.16.1.1, 34.16.1.1, 35.16.1.1, 36.16.1.1, 37.16.1.1,  
38.16.1.1, 39.16.1.1, 40.16.1.1, 1.17.1.1, 2.17.1.1, 3.17.1.1, 4.17.1.1, 5.17.1.1, 6.17.1.1,  
7.17.1.1, 8.17.1.1, 9.17.1.1, 10.17.1.1, 11.17.1.1, 12.17.1.1, 13.17.1.1, 14.17.1.1,  
15 15.17.1.1, 16.17.1.1, 17.17.1.1, 18.17.1.1, 19.17.1.1, 20.17.1.1, 21.17.1.1, 22.17.1.1,  
23.17.1.1, 24.17.1.1, 25.17.1.1, 26.17.1.1, 27.17.1.1, 28.17.1.1, 29.17.1.1, 30.17.1.1,  
31.17.1.1, 32.17.1.1, 33.17.1.1, 34.17.1.1, 35.17.1.1, 36.17.1.1, 37.17.1.1, 38.17.1.1,  
39.17.1.1, 40.17.1.1, 1.18.1.1, 2.18.1.1, 3.18.1.1, 4.18.1.1, 5.18.1.1, 6.18.1.1, 7.18.1.1,  
8.18.1.1, 9.18.1.1, 10.18.1.1, 11.18.1.1, 12.18.1.1, 13.18.1.1, 14.18.1.1, 15.18.1.1,  
20 16.18.1.1, 17.18.1.1, 18.18.1.1, 19.18.1.1, 20.18.1.1, 21.18.1.1, 22.18.1.1, 23.18.1.1,  
24.18.1.1, 25.18.1.1, 26.18.1.1, 27.18.1.1, 28.18.1.1, 29.18.1.1, 30.18.1.1, 31.18.1.1,  
32.18.1.1, 33.18.1.1, 34.18.1.1, 35.18.1.1, 36.18.1.1, 37.18.1.1, 38.18.1.1, 39.18.1.1,  
40.18.1.1, 1.19.1.1, 2.19.1.1, 3.19.1.1, 4.19.1.1, 5.19.1.1, 6.19.1.1, 7.19.1.1, 8.19.1.1,  
9.19.1.1, 10.19.1.1, 11.19.1.1, 12.19.1.1, 13.19.1.1, 14.19.1.1, 15.19.1.1, 16.19.1.1,  
25 17.19.1.1, 18.19.1.1, 19.19.1.1, 20.19.1.1, 21.19.1.1, 22.19.1.1, 23.19.1.1, 24.19.1.1,  
25.19.1.1, 26.19.1.1, 27.19.1.1, 28.19.1.1, 29.19.1.1, 30.19.1.1, 31.19.1.1, 32.19.1.1,  
33.19.1.1, 34.19.1.1, 35.19.1.1, 36.19.1.1, 37.19.1.1, 38.19.1.1, 39.19.1.1, 40.19.1.1,  
1.20.1.1, 2.20.1.1, 3.20.1.1, 4.20.1.1, 5.20.1.1, 6.20.1.1, 7.20.1.1, 8.20.1.1, 9.20.1.1,  
10.20.1.1, 11.20.1.1, 12.20.1.1, 13.20.1.1, 14.20.1.1, 15.20.1.1, 16.20.1.1, 17.20.1.1,  
30 18.20.1.1, 19.20.1.1, 20.20.1.1, 21.20.1.1, 22.20.1.1, 23.20.1.1, 24.20.1.1, 25.20.1.1,  
26.20.1.1, 27.20.1.1, 28.20.1.1, 29.20.1.1, 30.20.1.1, 31.20.1.1, 32.20.1.1, 33.20.1.1,  
34.20.1.1, 35.20.1.1, 36.20.1.1, 37.20.1.1, 38.20.1.1, 39.20.1.1, 40.20.1.1, 1.21.1.1,  
2.21.1.1, 3.21.1.1, 4.21.1.1, 5.21.1.1, 6.21.1.1, 7.21.1.1, 8.21.1.1, 9.21.1.1, 10.21.1.1,  
11.21.1.1, 12.21.1.1, 13.21.1.1, 14.21.1.1, 15.21.1.1, 16.21.1.1, 17.21.1.1, 18.21.1.1,  
35 19.21.1.1, 20.21.1.1, 21.21.1.1, 22.21.1.1, 23.21.1.1, 24.21.1.1, 25.21.1.1, 26.21.1.1,



27.21.1.1, 28.21.1.1, 29.21.1.1, 30.21.1.1, 31.21.1.1, 32.21.1.1, 33.21.1.1, 34.21.1.1,  
 35.21.1.1, 36.21.1.1, 37.21.1.1, 38.21.1.1, 39.21.1.1, 40.21.1.1, 1.22.1.1, 2.22.1.1,  
 3.22.1.1, 4.22.1.1, 5.22.1.1, 6.22.1.1, 7.22.1.1, 8.22.1.1, 9.22.1.1, 10.22.1.1, 11.22.1.1,  
 12.22.1.1, 13.22.1.1, 14.22.1.1, 15.22.1.1, 16.22.1.1, 17.22.1.1, 18.22.1.1, 19.22.1.1,  
 5 20.22.1.1, 21.22.1.1, 22.22.1.1, 23.22.1.1, 24.22.1.1, 25.22.1.1, 26.22.1.1, 27.22.1.1,  
 28.22.1.1, 29.22.1.1, 30.22.1.1, 31.22.1.1, 32.22.1.1, 33.22.1.1, 34.22.1.1, 35.22.1.1,  
 36.22.1.1, 37.22.1.1, 38.22.1.1, 39.22.1.1, 40.22.1.1, 1.23.1.1, 2.23.1.1, 3.23.1.1, 4.23.1.1,  
 5.23.1.1, 6.23.1.1, 7.23.1.1, 8.23.1.1, 9.23.1.1, 10.23.1.1, 11.23.1.1, 12.23.1.1, 13.23.1.1,  
 14.23.1.1, 15.23.1.1, 16.23.1.1, 17.23.1.1, 18.23.1.1, 19.23.1.1, 20.23.1.1, 21.23.1.1,  
 10 22.23.1.1, 23.23.1.1, 24.23.1.1, 25.23.1.1, 26.23.1.1, 27.23.1.1, 28.23.1.1, 29.23.1.1,  
 30.23.1.1, 31.23.1.1, 32.23.1.1, 33.23.1.1, 34.23.1.1, 35.23.1.1, 36.23.1.1, 37.23.1.1,  
 38.23.1.1, 39.23.1.1, 40.23.1.1, 1.24.1.1, 2.24.1.1, 3.24.1.1, 4.24.1.1, 5.24.1.1, 6.24.1.1,  
 7.24.1.1, 8.24.1.1, 9.24.1.1, 10.24.1.1, 11.24.1.1, 12.24.1.1, 13.24.1.1, 14.24.1.1,  
 15.24.1.1, 16.24.1.1, 17.24.1.1, 18.24.1.1, 19.24.1.1, 20.24.1.1, 21.24.1.1, 22.24.1.1,  
 15 23.24.1.1, 24.24.1.1, 25.24.1.1, 26.24.1.1, 27.24.1.1, 28.24.1.1, 29.24.1.1, 30.24.1.1,  
 31.24.1.1, 32.24.1.1, 33.24.1.1, 34.24.1.1, 35.24.1.1, 36.24.1.1, 37.24.1.1, 38.24.1.1,  
 39.24.1.1, 40.24.1.1, 1.25.1.1, 2.25.1.1, 3.25.1.1, 4.25.1.1, 5.25.1.1, 6.25.1.1, 7.25.1.1,  
 8.25.1.1, 9.25.1.1, 10.25.1.1, 11.25.1.1, 12.25.1.1, 13.25.1.1, 14.25.1.1, 15.25.1.1,  
 16.25.1.1, 17.25.1.1, 18.25.1.1, 19.25.1.1, 20.25.1.1, 21.25.1.1, 22.25.1.1, 23.25.1.1,  
 20 24.25.1.1, 25.25.1.1, 26.25.1.1, 27.25.1.1, 28.25.1.1, 29.25.1.1, 30.25.1.1, 31.25.1.1,  
 32.25.1.1, 33.25.1.1, 34.25.1.1, 35.25.1.1, 36.25.1.1, 37.25.1.1, 38.25.1.1, 39.25.1.1,  
 40.25.1.1, 1.26.1.1, 2.26.1.1, 3.26.1.1, 4.26.1.1, 5.26.1.1, 6.26.1.1, 7.26.1.1, 8.26.1.1,  
 9.26.1.1, 10.26.1.1, 11.26.1.1, 12.26.1.1, 13.26.1.1, 14.26.1.1, 15.26.1.1, 16.26.1.1,  
 17.26.1.1, 18.26.1.1, 19.26.1.1, 20.26.1.1, 21.26.1.1, 22.26.1.1, 23.26.1.1, 24.26.1.1,  
 25 25.26.1.1, 26.26.1.1, 27.26.1.1, 28.26.1.1, 29.26.1.1, 30.26.1.1, 31.26.1.1, 32.26.1.1,  
 33.26.1.1, 34.26.1.1, 35.26.1.1, 36.26.1.1, 37.26.1.1, 38.26.1.1, 39.26.1.1, 40.26.1.1,  
 1.27.1.1, 2.27.1.1, 3.27.1.1, 4.27.1.1, 5.27.1.1, 6.27.1.1, 7.27.1.1, 8.27.1.1, 9.27.1.1,  
 10.27.1.1, 11.27.1.1, 12.27.1.1, 13.27.1.1, 14.27.1.1, 15.27.1.1, 16.27.1.1, 17.27.1.1,  
 18.27.1.1, 19.27.1.1, 20.27.1.1, 21.27.1.1, 22.27.1.1, 23.27.1.1, 24.27.1.1, 25.27.1.1,  
 30 26.27.1.1, 27.27.1.1, 28.27.1.1, 29.27.1.1, 30.27.1.1, 31.27.1.1, 32.27.1.1, 33.27.1.1,  
 34.27.1.1, 35.27.1.1, 36.27.1.1, 37.27.1.1, 38.27.1.1, 39.27.1.1, 40.27.1.1, 1.28.1.1,  
 2.28.1.1, 3.28.1.1, 4.28.1.1, 5.28.1.1, 6.28.1.1, 7.28.1.1, 8.28.1.1, 9.28.1.1, 10.28.1.1,  
 11.28.1.1, 12.28.1.1, 13.28.1.1, 14.28.1.1, 15.28.1.1, 16.28.1.1, 17.28.1.1, 18.28.1.1,  
 19.28.1.1, 20.28.1.1, 21.28.1.1, 22.28.1.1, 23.28.1.1, 24.28.1.1, 25.28.1.1, 26.28.1.1,  
 35 27.28.1.1, 28.28.1.1, 29.28.1.1, 30.28.1.1, 31.28.1.1, 32.28.1.1, 33.28.1.1, 34.28.1.1,

35.28.1.1, 36.28.1.1, 37.28.1.1, 38.28.1.1, 39.28.1.1, 40.28.1.1, 1.29.1.1, 2.29.1.1,  
 3.29.1.1, 4.29.1.1, 5.29.1.1, 6.29.1.1, 7.29.1.1, 8.29.1.1, 9.29.1.1, 10.29.1.1, 11.29.1.1,  
 12.29.1.1, 13.29.1.1, 14.29.1.1, 15.29.1.1, 16.29.1.1, 17.29.1.1, 18.29.1.1, 19.29.1.1,  
 20.29.1.1, 21.29.1.1, 22.29.1.1, 23.29.1.1, 24.29.1.1, 25.29.1.1, 26.29.1.1, 27.29.1.1,  
 5 28.29.1.1, 29.29.1.1, 30.29.1.1, 31.29.1.1, 32.29.1.1, 33.29.1.1, 34.29.1.1, 35.29.1.1,  
 36.29.1.1, 37.29.1.1, 38.29.1.1, 39.29.1.1, 40.29.1.1, 1.1.1.2, 2.1.1.2, 3.1.1.2, 4.1.1.2,  
 5.1.1.2, 6.1.1.2, 7.1.1.2, 8.1.1.2, 9.1.1.2, 10.1.1.2, 11.1.1.2, 12.1.1.2, 13.1.1.2, 14.1.1.2,  
 15.1.1.2, 16.1.1.2, 17.1.1.2, 18.1.1.2, 19.1.1.2, 20.1.1.2, 21.1.1.2, 22.1.1.2, 23.1.1.2,  
 24.1.1.2, 25.1.1.2, 26.1.1.2, 27.1.1.2, 28.1.1.2, 29.1.1.2, 30.1.1.2, 31.1.1.2, 32.1.1.2,  
 10 33.1.1.2, 34.1.1.2, 35.1.1.2, 36.1.1.2, 37.1.1.2, 38.1.1.2, 39.1.1.2, 40.1.1.2, 1.2.1.2,  
 2.2.1.2, 3.2.1.2, 4.2.1.2, 5.2.1.2, 6.2.1.2, 7.2.1.2, 8.2.1.2, 9.2.1.2, 10.2.1.2, 11.2.1.2,  
 12.2.1.2, 13.2.1.2, 14.2.1.2, 15.2.1.2, 16.2.1.2, 17.2.1.2, 18.2.1.2, 19.2.1.2, 20.2.1.2,  
 21.2.1.2, 22.2.1.2, 23.2.1.2, 24.2.1.2, 25.2.1.2, 26.2.1.2, 27.2.1.2, 28.2.1.2, 29.2.1.2,  
 30.2.1.2, 31.2.1.2, 32.2.1.2, 33.2.1.2, 34.2.1.2, 35.2.1.2, 36.2.1.2, 37.2.1.2, 38.2.1.2,  
 15 39.2.1.2, 40.2.1.2, 1.3.1.2, 2.3.1.2, 3.3.1.2, 4.3.1.2, 5.3.1.2, 6.3.1.2, 7.3.1.2, 8.3.1.2,  
 9.3.1.2, 10.3.1.2, 11.3.1.2, 12.3.1.2, 13.3.1.2, 14.3.1.2, 15.3.1.2, 16.3.1.2, 17.3.1.2,  
 18.3.1.2, 19.3.1.2, 20.3.1.2, 21.3.1.2, 22.3.1.2, 23.3.1.2, 24.3.1.2, 25.3.1.2, 26.3.1.2,  
 27.3.1.2, 28.3.1.2, 29.3.1.2, 30.3.1.2, 31.3.1.2, 32.3.1.2, 33.3.1.2, 34.3.1.2, 35.3.1.2,  
 36.3.1.2, 37.3.1.2, 38.3.1.2, 39.3.1.2, 40.3.1.2, 1.4.1.2, 2.4.1.2, 3.4.1.2, 4.4.1.2, 5.4.1.2,  
 20 6.4.1.2, 7.4.1.2, 8.4.1.2, 9.4.1.2, 10.4.1.2, 11.4.1.2, 12.4.1.2, 13.4.1.2, 14.4.1.2, 15.4.1.2,  
 16.4.1.2, 17.4.1.2, 18.4.1.2, 19.4.1.2, 20.4.1.2, 21.4.1.2, 22.4.1.2, 23.4.1.2, 24.4.1.2,  
 25.4.1.2, 26.4.1.2, 27.4.1.2, 28.4.1.2, 29.4.1.2, 30.4.1.2, 31.4.1.2, 32.4.1.2, 33.4.1.2,  
 34.4.1.2, 35.4.1.2, 36.4.1.2, 37.4.1.2, 38.4.1.2, 39.4.1.2, 40.4.1.2, 1.5.1.2, 2.5.1.2, 3.5.1.2,  
 4.5.1.2, 5.5.1.2, 6.5.1.2, 7.5.1.2, 8.5.1.2, 9.5.1.2, 10.5.1.2, 11.5.1.2, 12.5.1.2, 13.5.1.2,  
 25 14.5.1.2, 15.5.1.2, 16.5.1.2, 17.5.1.2, 18.5.1.2, 19.5.1.2, 20.5.1.2, 21.5.1.2, 22.5.1.2,  
 23.5.1.2, 24.5.1.2, 25.5.1.2, 26.5.1.2, 27.5.1.2, 28.5.1.2, 29.5.1.2, 30.5.1.2, 31.5.1.2,  
 32.5.1.2, 33.5.1.2, 34.5.1.2, 35.5.1.2, 36.5.1.2, 37.5.1.2, 38.5.1.2, 39.5.1.2, 40.5.1.2,  
 1.6.1.2, 2.6.1.2, 3.6.1.2, 4.6.1.2, 5.6.1.2, 6.6.1.2, 7.6.1.2, 8.6.1.2, 9.6.1.2, 10.6.1.2,  
 11.6.1.2, 12.6.1.2, 13.6.1.2, 14.6.1.2, 15.6.1.2, 16.6.1.2, 17.6.1.2, 18.6.1.2, 19.6.1.2,  
 30 20.6.1.2, 21.6.1.2, 22.6.1.2, 23.6.1.2, 24.6.1.2, 25.6.1.2, 26.6.1.2, 27.6.1.2, 28.6.1.2,  
 29.6.1.2, 30.6.1.2, 31.6.1.2, 32.6.1.2, 33.6.1.2, 34.6.1.2, 35.6.1.2, 36.6.1.2, 37.6.1.2,  
 38.6.1.2, 39.6.1.2, 40.6.1.2, 1.7.1.2, 2.7.1.2, 3.7.1.2, 4.7.1.2, 5.7.1.2, 6.7.1.2, 7.7.1.2,  
 8.7.1.2, 9.7.1.2, 10.7.1.2, 11.7.1.2, 12.7.1.2, 13.7.1.2, 14.7.1.2, 15.7.1.2, 16.7.1.2,  
 17.7.1.2, 18.7.1.2, 19.7.1.2, 20.7.1.2, 21.7.1.2, 22.7.1.2, 23.7.1.2, 24.7.1.2, 25.7.1.2,  
 35 26.7.1.2, 27.7.1.2, 28.7.1.2, 29.7.1.2, 30.7.1.2, 31.7.1.2, 32.7.1.2, 33.7.1.2, 34.7.1.2,

35.7.1.2, 36.7.1.2, 37.7.1.2, 38.7.1.2, 39.7.1.2, 40.7.1.2, 1.8.1.2, 2.8.1.2, 3.8.1.2, 4.8.1.2,  
5.8.1.2, 6.8.1.2, 7.8.1.2, 8.8.1.2, 9.8.1.2, 10.8.1.2, 11.8.1.2, 12.8.1.2, 13.8.1.2, 14.8.1.2,  
15.8.1.2, 16.8.1.2, 17.8.1.2, 18.8.1.2, 19.8.1.2, 20.8.1.2, 21.8.1.2, 22.8.1.2, 23.8.1.2,  
24.8.1.2, 25.8.1.2, 26.8.1.2, 27.8.1.2, 28.8.1.2, 29.8.1.2, 30.8.1.2, 31.8.1.2, 32.8.1.2,  
5 33.8.1.2, 34.8.1.2, 35.8.1.2, 36.8.1.2, 37.8.1.2, 38.8.1.2, 39.8.1.2, 40.8.1.2, 1.9.1.2,  
2.9.1.2, 3.9.1.2, 4.9.1.2, 5.9.1.2, 6.9.1.2, 7.9.1.2, 8.9.1.2, 9.9.1.2, 10.9.1.2, 11.9.1.2,  
12.9.1.2, 13.9.1.2, 14.9.1.2, 15.9.1.2, 16.9.1.2, 17.9.1.2, 18.9.1.2, 19.9.1.2, 20.9.1.2,  
21.9.1.2, 22.9.1.2, 23.9.1.2, 24.9.1.2, 25.9.1.2, 26.9.1.2, 27.9.1.2, 28.9.1.2, 29.9.1.2,  
30.9.1.2, 31.9.1.2, 32.9.1.2, 33.9.1.2, 34.9.1.2, 35.9.1.2, 36.9.1.2, 37.9.1.2, 38.9.1.2,  
10 39.9.1.2, 40.9.1.2, 1.10.1.2, 2.10.1.2, 3.10.1.2, 4.10.1.2, 5.10.1.2, 6.10.1.2, 7.10.1.2,  
8.10.1.2, 9.10.1.2, 10.10.1.2, 11.10.1.2, 12.10.1.2, 13.10.1.2, 14.10.1.2, 15.10.1.2,  
16.10.1.2, 17.10.1.2, 18.10.1.2, 19.10.1.2, 20.10.1.2, 21.10.1.2, 22.10.1.2, 23.10.1.2,  
24.10.1.2, 25.10.1.2, 26.10.1.2, 27.10.1.2, 28.10.1.2, 29.10.1.2, 30.10.1.2, 31.10.1.2,  
32.10.1.2, 33.10.1.2, 34.10.1.2, 35.10.1.2, 36.10.1.2, 37.10.1.2, 38.10.1.2, 39.10.1.2,  
15 40.10.1.2, 1.11.1.2, 2.11.1.2, 3.11.1.2, 4.11.1.2, 5.11.1.2, 6.11.1.2, 7.11.1.2, 8.11.1.2,  
9.11.1.2, 10.11.1.2, 11.11.1.2, 12.11.1.2, 13.11.1.2, 14.11.1.2, 15.11.1.2, 16.11.1.2,  
17.11.1.2, 18.11.1.2, 19.11.1.2, 20.11.1.2, 21.11.1.2, 22.11.1.2, 23.11.1.2, 24.11.1.2,  
25.11.1.2, 26.11.1.2, 27.11.1.2, 28.11.1.2, 29.11.1.2, 30.11.1.2, 31.11.1.2, 32.11.1.2,  
33.11.1.2, 34.11.1.2, 35.11.1.2, 36.11.1.2, 37.11.1.2, 38.11.1.2, 39.11.1.2, 40.11.1.2,  
20 1.12.1.2, 2.12.1.2, 3.12.1.2, 4.12.1.2, 5.12.1.2, 6.12.1.2, 7.12.1.2, 8.12.1.2, 9.12.1.2,  
10.12.1.2, 11.12.1.2, 12.12.1.2, 13.12.1.2, 14.12.1.2, 15.12.1.2, 16.12.1.2, 17.12.1.2,  
18.12.1.2, 19.12.1.2, 20.12.1.2, 21.12.1.2, 22.12.1.2, 23.12.1.2, 24.12.1.2, 25.12.1.2,  
26.12.1.2, 27.12.1.2, 28.12.1.2, 29.12.1.2, 30.12.1.2, 31.12.1.2, 32.12.1.2, 33.12.1.2,  
34.12.1.2, 35.12.1.2, 36.12.1.2, 37.12.1.2, 38.12.1.2, 39.12.1.2, 40.12.1.2, 1.13.1.2,  
25 2.13.1.2, 3.13.1.2, 4.13.1.2, 5.13.1.2, 6.13.1.2, 7.13.1.2, 8.13.1.2, 9.13.1.2, 10.13.1.2,  
11.13.1.2, 12.13.1.2, 13.13.1.2, 14.13.1.2, 15.13.1.2, 16.13.1.2, 17.13.1.2, 18.13.1.2,  
19.13.1.2, 20.13.1.2, 21.13.1.2, 22.13.1.2, 23.13.1.2, 24.13.1.2, 25.13.1.2, 26.13.1.2,  
27.13.1.2, 28.13.1.2, 29.13.1.2, 30.13.1.2, 31.13.1.2, 32.13.1.2, 33.13.1.2, 34.13.1.2,  
35.13.1.2, 36.13.1.2, 37.13.1.2, 38.13.1.2, 39.13.1.2, 40.13.1.2, 1.14.1.2, 2.14.1.2,  
30 3.14.1.2, 4.14.1.2, 5.14.1.2, 6.14.1.2, 7.14.1.2, 8.14.1.2, 9.14.1.2, 10.14.1.2, 11.14.1.2,  
12.14.1.2, 13.14.1.2, 14.14.1.2, 15.14.1.2, 16.14.1.2, 17.14.1.2, 18.14.1.2, 19.14.1.2,  
20.14.1.2, 21.14.1.2, 22.14.1.2, 23.14.1.2, 24.14.1.2, 25.14.1.2, 26.14.1.2, 27.14.1.2,  
28.14.1.2, 29.14.1.2, 30.14.1.2, 31.14.1.2, 32.14.1.2, 33.14.1.2, 34.14.1.2, 35.14.1.2,  
36.14.1.2, 37.14.1.2, 38.14.1.2, 39.14.1.2, 40.14.1.2, 1.15.1.2, 2.15.1.2, 3.15.1.2, 4.15.1.2,  
35 5.15.1.2, 6.15.1.2, 7.15.1.2, 8.15.1.2, 9.15.1.2, 10.15.1.2, 11.15.1.2, 12.15.1.2, 13.15.1.2,

14.15.1.2, 15.15.1.2, 16.15.1.2, 17.15.1.2, 18.15.1.2, 19.15.1.2, 20.15.1.2, 21.15.1.2,  
 22.15.1.2, 23.15.1.2, 24.15.1.2, 25.15.1.2, 26.15.1.2, 27.15.1.2, 28.15.1.2, 29.15.1.2,  
 30.15.1.2, 31.15.1.2, 32.15.1.2, 33.15.1.2, 34.15.1.2, 35.15.1.2, 36.15.1.2, 37.15.1.2,  
 38.15.1.2, 39.15.1.2, 40.15.1.2, 1.16.1.2, 2.16.1.2, 3.16.1.2, 4.16.1.2, 5.16.1.2, 6.16.1.2,  
 5 7.16.1.2, 8.16.1.2, 9.16.1.2, 10.16.1.2, 11.16.1.2, 12.16.1.2, 13.16.1.2, 14.16.1.2,  
 15.16.1.2, 16.16.1.2, 17.16.1.2, 18.16.1.2, 19.16.1.2, 20.16.1.2, 21.16.1.2, 22.16.1.2,  
 23.16.1.2, 24.16.1.2, 25.16.1.2, 26.16.1.2, 27.16.1.2, 28.16.1.2, 29.16.1.2, 30.16.1.2,  
 31.16.1.2, 32.16.1.2, 33.16.1.2, 34.16.1.2, 35.16.1.2, 36.16.1.2, 37.16.1.2, 38.16.1.2,  
 39.16.1.2, 40.16.1.2, 1.17.1.2, 2.17.1.2, 3.17.1.2, 4.17.1.2, 5.17.1.2, 6.17.1.2, 7.17.1.2,  
 10 8.17.1.2, 9.17.1.2, 10.17.1.2, 11.17.1.2, 12.17.1.2, 13.17.1.2, 14.17.1.2, 15.17.1.2,  
 16.17.1.2, 17.17.1.2, 18.17.1.2, 19.17.1.2, 20.17.1.2, 21.17.1.2, 22.17.1.2, 23.17.1.2,  
 24.17.1.2, 25.17.1.2, 26.17.1.2, 27.17.1.2, 28.17.1.2, 29.17.1.2, 30.17.1.2, 31.17.1.2,  
 32.17.1.2, 33.17.1.2, 34.17.1.2, 35.17.1.2, 36.17.1.2, 37.17.1.2, 38.17.1.2, 39.17.1.2,  
 40.17.1.2, 1.18.1.2, 2.18.1.2, 3.18.1.2, 4.18.1.2, 5.18.1.2, 6.18.1.2, 7.18.1.2, 8.18.1.2,  
 15 9.18.1.2, 10.18.1.2, 11.18.1.2, 12.18.1.2, 13.18.1.2, 14.18.1.2, 15.18.1.2, 16.18.1.2,  
 17.18.1.2, 18.18.1.2, 19.18.1.2, 20.18.1.2, 21.18.1.2, 22.18.1.2, 23.18.1.2, 24.18.1.2,  
 25.18.1.2, 26.18.1.2, 27.18.1.2, 28.18.1.2, 29.18.1.2, 30.18.1.2, 31.18.1.2, 32.18.1.2,  
 33.18.1.2, 34.18.1.2, 35.18.1.2, 36.18.1.2, 37.18.1.2, 38.18.1.2, 39.18.1.2, 40.18.1.2,  
 1.19.1.2, 2.19.1.2, 3.19.1.2, 4.19.1.2, 5.19.1.2, 6.19.1.2, 7.19.1.2, 8.19.1.2, 9.19.1.2,  
 20 10.19.1.2, 11.19.1.2, 12.19.1.2, 13.19.1.2, 14.19.1.2, 15.19.1.2, 16.19.1.2, 17.19.1.2,  
 18.19.1.2, 19.19.1.2, 20.19.1.2, 21.19.1.2, 22.19.1.2, 23.19.1.2, 24.19.1.2, 25.19.1.2,  
 26.19.1.2, 27.19.1.2, 28.19.1.2, 29.19.1.2, 30.19.1.2, 31.19.1.2, 32.19.1.2, 33.19.1.2,  
 34.19.1.2, 35.19.1.2, 36.19.1.2, 37.19.1.2, 38.19.1.2, 39.19.1.2, 40.19.1.2, 1.20.1.2,  
 2.20.1.2, 3.20.1.2, 4.20.1.2, 5.20.1.2, 6.20.1.2, 7.20.1.2, 8.20.1.2, 9.20.1.2, 10.20.1.2,  
 25 11.20.1.2, 12.20.1.2, 13.20.1.2, 14.20.1.2, 15.20.1.2, 16.20.1.2, 17.20.1.2, 18.20.1.2,  
 19.20.1.2, 20.20.1.2, 21.20.1.2, 22.20.1.2, 23.20.1.2, 24.20.1.2, 25.20.1.2, 26.20.1.2,  
 27.20.1.2, 28.20.1.2, 29.20.1.2, 30.20.1.2, 31.20.1.2, 32.20.1.2, 33.20.1.2, 34.20.1.2,  
 35.20.1.2, 36.20.1.2, 37.20.1.2, 38.20.1.2, 39.20.1.2, 40.20.1.2, 1.21.1.2, 2.21.1.2,  
 3.21.1.2, 4.21.1.2, 5.21.1.2, 6.21.1.2, 7.21.1.2, 8.21.1.2, 9.21.1.2, 10.21.1.2, 11.21.1.2,  
 30 12.21.1.2, 13.21.1.2, 14.21.1.2, 15.21.1.2, 16.21.1.2, 17.21.1.2, 18.21.1.2, 19.21.1.2,  
 20.21.1.2, 21.21.1.2, 22.21.1.2, 23.21.1.2, 24.21.1.2, 25.21.1.2, 26.21.1.2, 27.21.1.2,  
 28.21.1.2, 29.21.1.2, 30.21.1.2, 31.21.1.2, 32.21.1.2, 33.21.1.2, 34.21.1.2, 35.21.1.2,  
 36.21.1.2, 37.21.1.2, 38.21.1.2, 39.21.1.2, 40.21.1.2, 1.22.1.2, 2.22.1.2, 3.22.1.2, 4.22.1.2,  
 5.22.1.2, 6.22.1.2, 7.22.1.2, 8.22.1.2, 9.22.1.2, 10.22.1.2, 11.22.1.2, 12.22.1.2, 13.22.1.2,  
 35 14.22.1.2, 15.22.1.2, 16.22.1.2, 17.22.1.2, 18.22.1.2, 19.22.1.2, 20.22.1.2, 21.22.1.2,

- 22.22.1.2, 23.22.1.2, 24.22.1.2, 25.22.1.2, 26.22.1.2, 27.22.1.2, 28.22.1.2, 29.22.1.2, 30.22.1.2, 31.22.1.2, 32.22.1.2, 33.22.1.2, 34.22.1.2, 35.22.1.2, 36.22.1.2, 37.22.1.2, 38.22.1.2, 39.22.1.2, 40.22.1.2, 1.23.1.2, 2.23.1.2, 3.23.1.2, 4.23.1.2, 5.23.1.2, 6.23.1.2, 7.23.1.2, 8.23.1.2, 9.23.1.2, 10.23.1.2, 11.23.1.2, 12.23.1.2, 13.23.1.2, 14.23.1.2,
- 5 15.23.1.2, 16.23.1.2, 17.23.1.2, 18.23.1.2, 19.23.1.2, 20.23.1.2, 21.23.1.2, 22.23.1.2, 23.23.1.2, 24.23.1.2, 25.23.1.2, 26.23.1.2, 27.23.1.2, 28.23.1.2, 29.23.1.2, 30.23.1.2, 31.23.1.2, 32.23.1.2, 33.23.1.2, 34.23.1.2, 35.23.1.2, 36.23.1.2, 37.23.1.2, 38.23.1.2, 39.23.1.2, 40.23.1.2, 1.24.1.2, 2.24.1.2, 3.24.1.2, 4.24.1.2, 5.24.1.2, 6.24.1.2, 7.24.1.2, 8.24.1.2, 9.24.1.2, 10.24.1.2, 11.24.1.2, 12.24.1.2, 13.24.1.2, 14.24.1.2, 15.24.1.2,
- 10 16.24.1.2, 17.24.1.2, 18.24.1.2, 19.24.1.2, 20.24.1.2, 21.24.1.2, 22.24.1.2, 23.24.1.2, 24.24.1.2, 25.24.1.2, 26.24.1.2, 27.24.1.2, 28.24.1.2, 29.24.1.2, 30.24.1.2, 31.24.1.2, 32.24.1.2, 33.24.1.2, 34.24.1.2, 35.24.1.2, 36.24.1.2, 37.24.1.2, 38.24.1.2, 39.24.1.2, 40.24.1.2, 1.25.1.2, 2.25.1.2, 3.25.1.2, 4.25.1.2, 5.25.1.2, 6.25.1.2, 7.25.1.2, 8.25.1.2, 9.25.1.2, 10.25.1.2, 11.25.1.2, 12.25.1.2, 13.25.1.2, 14.25.1.2, 15.25.1.2, 16.25.1.2,
- 15 17.25.1.2, 18.25.1.2, 19.25.1.2, 20.25.1.2, 21.25.1.2, 22.25.1.2, 23.25.1.2, 24.25.1.2, 25.25.1.2, 26.25.1.2, 27.25.1.2, 28.25.1.2, 29.25.1.2, 30.25.1.2, 31.25.1.2, 32.25.1.2, 33.25.1.2, 34.25.1.2, 35.25.1.2, 36.25.1.2, 37.25.1.2, 38.25.1.2, 39.25.1.2, 40.25.1.2, 1.26.1.2, 2.26.1.2, 3.26.1.2, 4.26.1.2, 5.26.1.2, 6.26.1.2, 7.26.1.2, 8.26.1.2, 9.26.1.2, 10.26.1.2, 11.26.1.2, 12.26.1.2, 13.26.1.2, 14.26.1.2, 15.26.1.2, 16.26.1.2, 17.26.1.2,
- 20 18.26.1.2, 19.26.1.2, 20.26.1.2, 21.26.1.2, 22.26.1.2, 23.26.1.2, 24.26.1.2, 25.26.1.2, 26.26.1.2, 27.26.1.2, 28.26.1.2, 29.26.1.2, 30.26.1.2, 31.26.1.2, 32.26.1.2, 33.26.1.2, 34.26.1.2, 35.26.1.2, 36.26.1.2, 37.26.1.2, 38.26.1.2, 39.26.1.2, 40.26.1.2, 1.27.1.2, 2.27.1.2, 3.27.1.2, 4.27.1.2, 5.27.1.2, 6.27.1.2, 7.27.1.2, 8.27.1.2, 9.27.1.2, 10.27.1.2, 11.27.1.2, 12.27.1.2, 13.27.1.2, 14.27.1.2, 15.27.1.2, 16.27.1.2, 17.27.1.2, 18.27.1.2,
- 25 19.27.1.2, 20.27.1.2, 21.27.1.2, 22.27.1.2, 23.27.1.2, 24.27.1.2, 25.27.1.2, 26.27.1.2, 27.27.1.2, 28.27.1.2, 29.27.1.2, 30.27.1.2, 31.27.1.2, 32.27.1.2, 33.27.1.2, 34.27.1.2, 35.27.1.2, 36.27.1.2, 37.27.1.2, 38.27.1.2, 39.27.1.2, 40.27.1.2, 1.28.1.2, 2.28.1.2, 3.28.1.2, 4.28.1.2, 5.28.1.2, 6.28.1.2, 7.28.1.2, 8.28.1.2, 9.28.1.2, 10.28.1.2, 11.28.1.2, 12.28.1.2, 13.28.1.2, 14.28.1.2, 15.28.1.2, 16.28.1.2, 17.28.1.2, 18.28.1.2, 19.28.1.2,
- 30 20.28.1.2, 21.28.1.2, 22.28.1.2, 23.28.1.2, 24.28.1.2, 25.28.1.2, 26.28.1.2, 27.28.1.2, 28.28.1.2, 29.28.1.2, 30.28.1.2, 31.28.1.2, 32.28.1.2, 33.28.1.2, 34.28.1.2, 35.28.1.2, 36.28.1.2, 37.28.1.2, 38.28.1.2, 39.28.1.2, 40.28.1.2, 1.29.1.2, 2.29.1.2, 3.29.1.2, 4.29.1.2, 5.29.1.2, 6.29.1.2, 7.29.1.2, 8.29.1.2, 9.29.1.2, 10.29.1.2, 11.29.1.2, 12.29.1.2, 13.29.1.2, 14.29.1.2, 15.29.1.2, 16.29.1.2, 17.29.1.2, 18.29.1.2, 19.29.1.2, 20.29.1.2, 21.29.1.2,
- 35 22.29.1.2, 23.29.1.2, 24.29.1.2, 25.29.1.2, 26.29.1.2, 27.29.1.2, 28.29.1.2, 29.29.1.2,

30.29.1.2, 31.29.1.2, 32.29.1.2, 33.29.1.2, 34.29.1.2, 35.29.1.2, 36.29.1.2, 37.29.1.2,  
 38.29.1.2, 39.29.1.2, 40.29.1.2, 1.1.3.1, 2.1.3.1, 3.1.3.1, 4.1.3.1, 5.1.3.1, 6.1.3.1, 7.1.3.1,  
 8.1.3.1, 9.1.3.1, 10.1.3.1, 11.1.3.1, 12.1.3.1, 13.1.3.1, 14.1.3.1, 15.1.3.1, 16.1.3.1,  
 17.1.3.1, 18.1.3.1, 19.1.3.1, 20.1.3.1, 21.1.3.1, 22.1.3.1, 23.1.3.1, 24.1.3.1, 25.1.3.1,  
 5 26.1.3.1, 27.1.3.1, 28.1.3.1, 29.1.3.1, 30.1.3.1, 31.1.3.1, 32.1.3.1, 33.1.3.1, 34.1.3.1,  
 35.1.3.1, 36.1.3.1, 37.1.3.1, 38.1.3.1, 39.1.3.1, 40.1.3.1, 1.2.3.1, 2.2.3.1, 3.2.3.1, 4.2.3.1,  
 5.2.3.1, 6.2.3.1, 7.2.3.1, 8.2.3.1, 9.2.3.1, 10.2.3.1, 11.2.3.1, 12.2.3.1, 13.2.3.1, 14.2.3.1,  
 15.2.3.1, 16.2.3.1, 17.2.3.1, 18.2.3.1, 19.2.3.1, 20.2.3.1, 21.2.3.1, 22.2.3.1, 23.2.3.1,  
 24.2.3.1, 25.2.3.1, 26.2.3.1, 27.2.3.1, 28.2.3.1, 29.2.3.1, 30.2.3.1, 31.2.3.1, 32.2.3.1,  
 10 33.2.3.1, 34.2.3.1, 35.2.3.1, 36.2.3.1, 37.2.3.1, 38.2.3.1, 39.2.3.1, 40.2.3.1, 1.3.3.1,  
 2.3.3.1, 3.3.3.1, 4.3.3.1, 5.3.3.1, 6.3.3.1, 7.3.3.1, 8.3.3.1, 9.3.3.1, 10.3.3.1, 11.3.3.1,  
 12.3.3.1, 13.3.3.1, 14.3.3.1, 15.3.3.1, 16.3.3.1, 17.3.3.1, 18.3.3.1, 19.3.3.1, 20.3.3.1,  
 21.3.3.1, 22.3.3.1, 23.3.3.1, 24.3.3.1, 25.3.3.1, 26.3.3.1, 27.3.3.1, 28.3.3.1, 29.3.3.1,  
 30.3.3.1, 31.3.3.1, 32.3.3.1, 33.3.3.1, 34.3.3.1, 35.3.3.1, 36.3.3.1, 37.3.3.1, 38.3.3.1,  
 15 39.3.3.1, 40.3.3.1, 1.4.3.1, 2.4.3.1, 3.4.3.1, 4.4.3.1, 5.4.3.1, 6.4.3.1, 7.4.3.1, 8.4.3.1,  
 9.4.3.1, 10.4.3.1, 11.4.3.1, 12.4.3.1, 13.4.3.1, 14.4.3.1, 15.4.3.1, 16.4.3.1, 17.4.3.1,  
 18.4.3.1, 19.4.3.1, 20.4.3.1, 21.4.3.1, 22.4.3.1, 23.4.3.1, 24.4.3.1, 25.4.3.1, 26.4.3.1,  
 27.4.3.1, 28.4.3.1, 29.4.3.1, 30.4.3.1, 31.4.3.1, 32.4.3.1, 33.4.3.1, 34.4.3.1, 35.4.3.1,  
 36.4.3.1, 37.4.3.1, 38.4.3.1, 39.4.3.1, 40.4.3.1, 1.5.3.1, 2.5.3.1, 3.5.3.1, 4.5.3.1, 5.5.3.1,  
 20 6.5.3.1, 7.5.3.1, 8.5.3.1, 9.5.3.1, 10.5.3.1, 11.5.3.1, 12.5.3.1, 13.5.3.1, 14.5.3.1, 15.5.3.1,  
 16.5.3.1, 17.5.3.1, 18.5.3.1, 19.5.3.1, 20.5.3.1, 21.5.3.1, 22.5.3.1, 23.5.3.1, 24.5.3.1,  
 25.5.3.1, 26.5.3.1, 27.5.3.1, 28.5.3.1, 29.5.3.1, 30.5.3.1, 31.5.3.1, 32.5.3.1, 33.5.3.1,  
 34.5.3.1, 35.5.3.1, 36.5.3.1, 37.5.3.1, 38.5.3.1, 39.5.3.1, 40.5.3.1, 1.6.3.1, 2.6.3.1, 3.6.3.1,  
 4.6.3.1, 5.6.3.1, 6.6.3.1, 7.6.3.1, 8.6.3.1, 9.6.3.1, 10.6.3.1, 11.6.3.1, 12.6.3.1, 13.6.3.1,  
 25 14.6.3.1, 15.6.3.1, 16.6.3.1, 17.6.3.1, 18.6.3.1, 19.6.3.1, 20.6.3.1, 21.6.3.1, 22.6.3.1,  
 23.6.3.1, 24.6.3.1, 25.6.3.1, 26.6.3.1, 27.6.3.1, 28.6.3.1, 29.6.3.1, 30.6.3.1, 31.6.3.1,  
 32.6.3.1, 33.6.3.1, 34.6.3.1, 35.6.3.1, 36.6.3.1, 37.6.3.1, 38.6.3.1, 39.6.3.1, 40.6.3.1,  
 1.7.3.1, 2.7.3.1, 3.7.3.1, 4.7.3.1, 5.7.3.1, 6.7.3.1, 7.7.3.1, 8.7.3.1, 9.7.3.1, 10.7.3.1,  
 11.7.3.1, 12.7.3.1, 13.7.3.1, 14.7.3.1, 15.7.3.1, 16.7.3.1, 17.7.3.1, 18.7.3.1, 19.7.3.1,  
 30 20.7.3.1, 21.7.3.1, 22.7.3.1, 23.7.3.1, 24.7.3.1, 25.7.3.1, 26.7.3.1, 27.7.3.1, 28.7.3.1,  
 29.7.3.1, 30.7.3.1, 31.7.3.1, 32.7.3.1, 33.7.3.1, 34.7.3.1, 35.7.3.1, 36.7.3.1, 37.7.3.1,  
 38.7.3.1, 39.7.3.1, 40.7.3.1, 1.8.3.1, 2.8.3.1, 3.8.3.1, 4.8.3.1, 5.8.3.1, 6.8.3.1, 7.8.3.1,  
 8.8.3.1, 9.8.3.1, 10.8.3.1, 11.8.3.1, 12.8.3.1, 13.8.3.1, 14.8.3.1, 15.8.3.1, 16.8.3.1,  
 17.8.3.1, 18.8.3.1, 19.8.3.1, 20.8.3.1, 21.8.3.1, 22.8.3.1, 23.8.3.1, 24.8.3.1, 25.8.3.1,  
 35 26.8.3.1, 27.8.3.1, 28.8.3.1, 29.8.3.1, 30.8.3.1, 31.8.3.1, 32.8.3.1, 33.8.3.1, 34.8.3.1,

35.8.3.1, 36.8.3.1, 37.8.3.1, 38.8.3.1, 39.8.3.1, 40.8.3.1, 1.9.3.1, 2.9.3.1, 3.9.3.1, 4.9.3.1,  
5.9.3.1, 6.9.3.1, 7.9.3.1, 8.9.3.1, 9.9.3.1, 10.9.3.1, 11.9.3.1, 12.9.3.1, 13.9.3.1, 14.9.3.1,  
15.9.3.1, 16.9.3.1, 17.9.3.1, 18.9.3.1, 19.9.3.1, 20.9.3.1, 21.9.3.1, 22.9.3.1, 23.9.3.1,  
24.9.3.1, 25.9.3.1, 26.9.3.1, 27.9.3.1, 28.9.3.1, 29.9.3.1, 30.9.3.1, 31.9.3.1, 32.9.3.1,  
5 33.9.3.1, 34.9.3.1, 35.9.3.1, 36.9.3.1, 37.9.3.1, 38.9.3.1, 39.9.3.1, 40.9.3.1, 1.10.3.1,  
2.10.3.1, 3.10.3.1, 4.10.3.1, 5.10.3.1, 6.10.3.1, 7.10.3.1, 8.10.3.1, 9.10.3.1, 10.10.3.1,  
11.10.3.1, 12.10.3.1, 13.10.3.1, 14.10.3.1, 15.10.3.1, 16.10.3.1, 17.10.3.1, 18.10.3.1,  
19.10.3.1, 20.10.3.1, 21.10.3.1, 22.10.3.1, 23.10.3.1, 24.10.3.1, 25.10.3.1, 26.10.3.1,  
27.10.3.1, 28.10.3.1, 29.10.3.1, 30.10.3.1, 31.10.3.1, 32.10.3.1, 33.10.3.1, 34.10.3.1,  
10 35.10.3.1, 36.10.3.1, 37.10.3.1, 38.10.3.1, 39.10.3.1, 40.10.3.1, 1.11.3.1, 2.11.3.1,  
3.11.3.1, 4.11.3.1, 5.11.3.1, 6.11.3.1, 7.11.3.1, 8.11.3.1, 9.11.3.1, 10.11.3.1, 11.11.3.1,  
12.11.3.1, 13.11.3.1, 14.11.3.1, 15.11.3.1, 16.11.3.1, 17.11.3.1, 18.11.3.1, 19.11.3.1,  
20.11.3.1, 21.11.3.1, 22.11.3.1, 23.11.3.1, 24.11.3.1, 25.11.3.1, 26.11.3.1, 27.11.3.1,  
28.11.3.1, 29.11.3.1, 30.11.3.1, 31.11.3.1, 32.11.3.1, 33.11.3.1, 34.11.3.1, 35.11.3.1,  
15 36.11.3.1, 37.11.3.1, 38.11.3.1, 39.11.3.1, 40.11.3.1, 1.12.3.1, 2.12.3.1, 3.12.3.1, 4.12.3.1,  
5.12.3.1, 6.12.3.1, 7.12.3.1, 8.12.3.1, 9.12.3.1, 10.12.3.1, 11.12.3.1, 12.12.3.1, 13.12.3.1,  
14.12.3.1, 15.12.3.1, 16.12.3.1, 17.12.3.1, 18.12.3.1, 19.12.3.1, 20.12.3.1, 21.12.3.1,  
22.12.3.1, 23.12.3.1, 24.12.3.1, 25.12.3.1, 26.12.3.1, 27.12.3.1, 28.12.3.1, 29.12.3.1,  
30.12.3.1, 31.12.3.1, 32.12.3.1, 33.12.3.1, 34.12.3.1, 35.12.3.1, 36.12.3.1, 37.12.3.1,  
20 38.12.3.1, 39.12.3.1, 40.12.3.1, 1.13.3.1, 2.13.3.1, 3.13.3.1, 4.13.3.1, 5.13.3.1, 6.13.3.1,  
7.13.3.1, 8.13.3.1, 9.13.3.1, 10.13.3.1, 11.13.3.1, 12.13.3.1, 13.13.3.1, 14.13.3.1,  
15.13.3.1, 16.13.3.1, 17.13.3.1, 18.13.3.1, 19.13.3.1, 20.13.3.1, 21.13.3.1, 22.13.3.1,  
23.13.3.1, 24.13.3.1, 25.13.3.1, 26.13.3.1, 27.13.3.1, 28.13.3.1, 29.13.3.1, 30.13.3.1,  
31.13.3.1, 32.13.3.1, 33.13.3.1, 34.13.3.1, 35.13.3.1, 36.13.3.1, 37.13.3.1, 38.13.3.1,  
25 39.13.3.1, 40.13.3.1, 1.14.3.1, 2.14.3.1, 3.14.3.1, 4.14.3.1, 5.14.3.1, 6.14.3.1, 7.14.3.1,  
8.14.3.1, 9.14.3.1, 10.14.3.1, 11.14.3.1, 12.14.3.1, 13.14.3.1, 14.14.3.1, 15.14.3.1,  
16.14.3.1, 17.14.3.1, 18.14.3.1, 19.14.3.1, 20.14.3.1, 21.14.3.1, 22.14.3.1, 23.14.3.1,  
24.14.3.1, 25.14.3.1, 26.14.3.1, 27.14.3.1, 28.14.3.1, 29.14.3.1, 30.14.3.1, 31.14.3.1,  
32.14.3.1, 33.14.3.1, 34.14.3.1, 35.14.3.1, 36.14.3.1, 37.14.3.1, 38.14.3.1, 39.14.3.1,  
30 40.14.3.1, 1.15.3.1, 2.15.3.1, 3.15.3.1, 4.15.3.1, 5.15.3.1, 6.15.3.1, 7.15.3.1, 8.15.3.1,  
9.15.3.1, 10.15.3.1, 11.15.3.1, 12.15.3.1, 13.15.3.1, 14.15.3.1, 15.15.3.1, 16.15.3.1,  
17.15.3.1, 18.15.3.1, 19.15.3.1, 20.15.3.1, 21.15.3.1, 22.15.3.1, 23.15.3.1, 24.15.3.1,  
25.15.3.1, 26.15.3.1, 27.15.3.1, 28.15.3.1, 29.15.3.1, 30.15.3.1, 31.15.3.1, 32.15.3.1,  
33.15.3.1, 34.15.3.1, 35.15.3.1, 36.15.3.1, 37.15.3.1, 38.15.3.1, 39.15.3.1, 40.15.3.1,  
35 1.16.3.1, 2.16.3.1, 3.16.3.1, 4.16.3.1, 5.16.3.1, 6.16.3.1, 7.16.3.1, 8.16.3.1, 9.16.3.1,

- 10.16.3.1, 11.16.3.1, 12.16.3.1, 13.16.3.1, 14.16.3.1, 15.16.3.1, 16.16.3.1, 17.16.3.1,  
18.16.3.1, 19.16.3.1, 20.16.3.1, 21.16.3.1, 22.16.3.1, 23.16.3.1, 24.16.3.1, 25.16.3.1,  
26.16.3.1, 27.16.3.1, 28.16.3.1, 29.16.3.1, 30.16.3.1, 31.16.3.1, 32.16.3.1, 33.16.3.1,  
34.16.3.1, 35.16.3.1, 36.16.3.1, 37.16.3.1, 38.16.3.1, 39.16.3.1, 40.16.3.1, 1.17.3.1,  
5 2.17.3.1, 3.17.3.1, 4.17.3.1, 5.17.3.1, 6.17.3.1, 7.17.3.1, 8.17.3.1, 9.17.3.1, 10.17.3.1,  
11.17.3.1, 12.17.3.1, 13.17.3.1, 14.17.3.1, 15.17.3.1, 16.17.3.1, 17.17.3.1, 18.17.3.1,  
19.17.3.1, 20.17.3.1, 21.17.3.1, 22.17.3.1, 23.17.3.1, 24.17.3.1, 25.17.3.1, 26.17.3.1,  
27.17.3.1, 28.17.3.1, 29.17.3.1, 30.17.3.1, 31.17.3.1, 32.17.3.1, 33.17.3.1, 34.17.3.1,  
35.17.3.1, 36.17.3.1, 37.17.3.1, 38.17.3.1, 39.17.3.1, 40.17.3.1, 1.18.3.1, 2.18.3.1,  
10 3.18.3.1, 4.18.3.1, 5.18.3.1, 6.18.3.1, 7.18.3.1, 8.18.3.1, 9.18.3.1, 10.18.3.1, 11.18.3.1,  
12.18.3.1, 13.18.3.1, 14.18.3.1, 15.18.3.1, 16.18.3.1, 17.18.3.1, 18.18.3.1, 19.18.3.1,  
20.18.3.1, 21.18.3.1, 22.18.3.1, 23.18.3.1, 24.18.3.1, 25.18.3.1, 26.18.3.1, 27.18.3.1,  
28.18.3.1, 29.18.3.1, 30.18.3.1, 31.18.3.1, 32.18.3.1, 33.18.3.1, 34.18.3.1, 35.18.3.1,  
36.18.3.1, 37.18.3.1, 38.18.3.1, 39.18.3.1, 40.18.3.1, 1.19.3.1, 2.19.3.1, 3.19.3.1, 4.19.3.1,  
15 5.19.3.1, 6.19.3.1, 7.19.3.1, 8.19.3.1, 9.19.3.1, 10.19.3.1, 11.19.3.1, 12.19.3.1, 13.19.3.1,  
14.19.3.1, 15.19.3.1, 16.19.3.1, 17.19.3.1, 18.19.3.1, 19.19.3.1, 20.19.3.1, 21.19.3.1,  
22.19.3.1, 23.19.3.1, 24.19.3.1, 25.19.3.1, 26.19.3.1, 27.19.3.1, 28.19.3.1, 29.19.3.1,  
30.19.3.1, 31.19.3.1, 32.19.3.1, 33.19.3.1, 34.19.3.1, 35.19.3.1, 36.19.3.1, 37.19.3.1,  
38.19.3.1, 39.19.3.1, 40.19.3.1, 1.20.3.1, 2.20.3.1, 3.20.3.1, 4.20.3.1, 5.20.3.1, 6.20.3.1,  
20 7.20.3.1, 8.20.3.1, 9.20.3.1, 10.20.3.1, 11.20.3.1, 12.20.3.1, 13.20.3.1, 14.20.3.1,  
15.20.3.1, 16.20.3.1, 17.20.3.1, 18.20.3.1, 19.20.3.1, 20.20.3.1, 21.20.3.1, 22.20.3.1,  
23.20.3.1, 24.20.3.1, 25.20.3.1, 26.20.3.1, 27.20.3.1, 28.20.3.1, 29.20.3.1, 30.20.3.1,  
31.20.3.1, 32.20.3.1, 33.20.3.1, 34.20.3.1, 35.20.3.1, 36.20.3.1, 37.20.3.1, 38.20.3.1,  
39.20.3.1, 40.20.3.1, 1.21.3.1, 2.21.3.1, 3.21.3.1, 4.21.3.1, 5.21.3.1, 6.21.3.1, 7.21.3.1,  
25 8.21.3.1, 9.21.3.1, 10.21.3.1, 11.21.3.1, 12.21.3.1, 13.21.3.1, 14.21.3.1, 15.21.3.1,  
16.21.3.1, 17.21.3.1, 18.21.3.1, 19.21.3.1, 20.21.3.1, 21.21.3.1, 22.21.3.1, 23.21.3.1,  
24.21.3.1, 25.21.3.1, 26.21.3.1, 27.21.3.1, 28.21.3.1, 29.21.3.1, 30.21.3.1, 31.21.3.1,  
32.21.3.1, 33.21.3.1, 34.21.3.1, 35.21.3.1, 36.21.3.1, 37.21.3.1, 38.21.3.1, 39.21.3.1,  
40.21.3.1, 1.22.3.1, 2.22.3.1, 3.22.3.1, 4.22.3.1, 5.22.3.1, 6.22.3.1, 7.22.3.1, 8.22.3.1,  
30 9.22.3.1, 10.22.3.1, 11.22.3.1, 12.22.3.1, 13.22.3.1, 14.22.3.1, 15.22.3.1, 16.22.3.1,  
17.22.3.1, 18.22.3.1, 19.22.3.1, 20.22.3.1, 21.22.3.1, 22.22.3.1, 23.22.3.1, 24.22.3.1,  
25.22.3.1, 26.22.3.1, 27.22.3.1, 28.22.3.1, 29.22.3.1, 30.22.3.1, 31.22.3.1, 32.22.3.1,  
33.22.3.1, 34.22.3.1, 35.22.3.1, 36.22.3.1, 37.22.3.1, 38.22.3.1, 39.22.3.1, 40.22.3.1,  
1.23.3.1, 2.23.3.1, 3.23.3.1, 4.23.3.1, 5.23.3.1, 6.23.3.1, 7.23.3.1, 8.23.3.1, 9.23.3.1,  
35 10.23.3.1, 11.23.3.1, 12.23.3.1, 13.23.3.1, 14.23.3.1, 15.23.3.1, 16.23.3.1, 17.23.3.1,



- 18.23.3.1, 19.23.3.1, 20.23.3.1, 21.23.3.1, 22.23.3.1, 23.23.3.1, 24.23.3.1, 25.23.3.1, 26.23.3.1, 27.23.3.1, 28.23.3.1, 29.23.3.1, 30.23.3.1, 31.23.3.1, 32.23.3.1, 33.23.3.1, 34.23.3.1, 35.23.3.1, 36.23.3.1, 37.23.3.1, 38.23.3.1, 39.23.3.1, 40.23.3.1, 1.24.3.1, 2.24.3.1, 3.24.3.1, 4.24.3.1, 5.24.3.1, 6.24.3.1, 7.24.3.1, 8.24.3.1, 9.24.3.1, 10.24.3.1,
- 5 11.24.3.1, 12.24.3.1, 13.24.3.1, 14.24.3.1, 15.24.3.1, 16.24.3.1, 17.24.3.1, 18.24.3.1, 19.24.3.1, 20.24.3.1, 21.24.3.1, 22.24.3.1, 23.24.3.1, 24.24.3.1, 25.24.3.1, 26.24.3.1, 27.24.3.1, 28.24.3.1, 29.24.3.1, 30.24.3.1, 31.24.3.1, 32.24.3.1, 33.24.3.1, 34.24.3.1, 35.24.3.1, 36.24.3.1, 37.24.3.1, 38.24.3.1, 39.24.3.1, 40.24.3.1, 1.25.3.1, 2.25.3.1, 3.25.3.1, 4.25.3.1, 5.25.3.1, 6.25.3.1, 7.25.3.1, 8.25.3.1, 9.25.3.1, 10.25.3.1, 11.25.3.1,
- 10 12.25.3.1, 13.25.3.1, 14.25.3.1, 15.25.3.1, 16.25.3.1, 17.25.3.1, 18.25.3.1, 19.25.3.1, 20.25.3.1, 21.25.3.1, 22.25.3.1, 23.25.3.1, 24.25.3.1, 25.25.3.1, 26.25.3.1, 27.25.3.1, 28.25.3.1, 29.25.3.1, 30.25.3.1, 31.25.3.1, 32.25.3.1, 33.25.3.1, 34.25.3.1, 35.25.3.1, 36.25.3.1, 37.25.3.1, 38.25.3.1, 39.25.3.1, 40.25.3.1, 1.26.3.1, 2.26.3.1, 3.26.3.1, 4.26.3.1, 5.26.3.1, 6.26.3.1, 7.26.3.1, 8.26.3.1, 9.26.3.1, 10.26.3.1, 11.26.3.1, 12.26.3.1, 13.26.3.1,
- 15 14.26.3.1, 15.26.3.1, 16.26.3.1, 17.26.3.1, 18.26.3.1, 19.26.3.1, 20.26.3.1, 21.26.3.1, 22.26.3.1, 23.26.3.1, 24.26.3.1, 25.26.3.1, 26.26.3.1, 27.26.3.1, 28.26.3.1, 29.26.3.1, 30.26.3.1, 31.26.3.1, 32.26.3.1, 33.26.3.1, 34.26.3.1, 35.26.3.1, 36.26.3.1, 37.26.3.1, 38.26.3.1, 39.26.3.1, 40.26.3.1, 1.27.3.1, 2.27.3.1, 3.27.3.1, 4.27.3.1, 5.27.3.1, 6.27.3.1, 7.27.3.1, 8.27.3.1, 9.27.3.1, 10.27.3.1, 11.27.3.1, 12.27.3.1, 13.27.3.1, 14.27.3.1,
- 20 15.27.3.1, 16.27.3.1, 17.27.3.1, 18.27.3.1, 19.27.3.1, 20.27.3.1, 21.27.3.1, 22.27.3.1, 23.27.3.1, 24.27.3.1, 25.27.3.1, 26.27.3.1, 27.27.3.1, 28.27.3.1, 29.27.3.1, 30.27.3.1, 31.27.3.1, 32.27.3.1, 33.27.3.1, 34.27.3.1, 35.27.3.1, 36.27.3.1, 37.27.3.1, 38.27.3.1, 39.27.3.1, 40.27.3.1, 1.28.3.1, 2.28.3.1, 3.28.3.1, 4.28.3.1, 5.28.3.1, 6.28.3.1, 7.28.3.1, 8.28.3.1, 9.28.3.1, 10.28.3.1, 11.28.3.1, 12.28.3.1, 13.28.3.1, 14.28.3.1, 15.28.3.1,
- 25 16.28.3.1, 17.28.3.1, 18.28.3.1, 19.28.3.1, 20.28.3.1, 21.28.3.1, 22.28.3.1, 23.28.3.1, 24.28.3.1, 25.28.3.1, 26.28.3.1, 27.28.3.1, 28.28.3.1, 29.28.3.1, 30.28.3.1, 31.28.3.1, 32.28.3.1, 33.28.3.1, 34.28.3.1, 35.28.3.1, 36.28.3.1, 37.28.3.1, 38.28.3.1, 39.28.3.1, 40.28.3.1, 1.29.3.1, 2.29.3.1, 3.29.3.1, 4.29.3.1, 5.29.3.1, 6.29.3.1, 7.29.3.1, 8.29.3.1, 9.29.3.1, 10.29.3.1, 11.29.3.1, 12.29.3.1, 13.29.3.1, 14.29.3.1, 15.29.3.1, 16.29.3.1,
- 30 17.29.3.1, 18.29.3.1, 19.29.3.1, 20.29.3.1, 21.29.3.1, 22.29.3.1, 23.29.3.1, 24.29.3.1, 25.29.3.1, 26.29.3.1, 27.29.3.1, 28.29.3.1, 29.29.3.1, 30.29.3.1, 31.29.3.1, 32.29.3.1, 33.29.3.1, 34.29.3.1, 35.29.3.1, 36.29.3.1, 37.29.3.1, 38.29.3.1, 39.29.3.1, 40.29.3.1, 1.1.3.2, 2.1.3.2, 3.1.3.2, 4.1.3.2, 5.1.3.2, 6.1.3.2, 7.1.3.2, 8.1.3.2, 9.1.3.2, 10.1.3.2, 11.1.3.2, 12.1.3.2, 13.1.3.2, 14.1.3.2, 15.1.3.2, 16.1.3.2, 17.1.3.2, 18.1.3.2, 19.1.3.2,
- 35 20.1.3.2, 21.1.3.2, 22.1.3.2, 23.1.3.2, 24.1.3.2, 25.1.3.2, 26.1.3.2, 27.1.3.2, 28.1.3.2,

- 29.1.3.2, 30.1.3.2, 31.1.3.2, 32.1.3.2, 33.1.3.2, 34.1.3.2, 35.1.3.2, 36.1.3.2, 37.1.3.2, 38.1.3.2, 39.1.3.2, 40.1.3.2, 1.2.3.2, 2.2.3.2, 3.2.3.2, 4.2.3.2, 5.2.3.2, 6.2.3.2, 7.2.3.2, 8.2.3.2, 9.2.3.2, 10.2.3.2, 11.2.3.2, 12.2.3.2, 13.2.3.2, 14.2.3.2, 15.2.3.2, 16.2.3.2, 17.2.3.2, 18.2.3.2, 19.2.3.2, 20.2.3.2, 21.2.3.2, 22.2.3.2, 23.2.3.2, 24.2.3.2, 25.2.3.2,
- 5 26.2.3.2, 27.2.3.2, 28.2.3.2, 29.2.3.2, 30.2.3.2, 31.2.3.2, 32.2.3.2, 33.2.3.2, 34.2.3.2, 35.2.3.2, 36.2.3.2, 37.2.3.2, 38.2.3.2, 39.2.3.2, 40.2.3.2, 1.3.3.2, 2.3.3.2, 3.3.3.2, 4.3.3.2, 5.3.3.2, 6.3.3.2, 7.3.3.2, 8.3.3.2, 9.3.3.2, 10.3.3.2, 11.3.3.2, 12.3.3.2, 13.3.3.2, 14.3.3.2, 15.3.3.2, 16.3.3.2, 17.3.3.2, 18.3.3.2, 19.3.3.2, 20.3.3.2, 21.3.3.2, 22.3.3.2, 23.3.3.2, 24.3.3.2, 25.3.3.2, 26.3.3.2, 27.3.3.2, 28.3.3.2, 29.3.3.2, 30.3.3.2, 31.3.3.2, 32.3.3.2,
- 10 33.3.3.2, 34.3.3.2, 35.3.3.2, 36.3.3.2, 37.3.3.2, 38.3.3.2, 39.3.3.2, 40.3.3.2, 1.4.3.2, 2.4.3.2, 3.4.3.2, 4.4.3.2, 5.4.3.2, 6.4.3.2, 7.4.3.2, 8.4.3.2, 9.4.3.2, 10.4.3.2, 11.4.3.2, 12.4.3.2, 13.4.3.2, 14.4.3.2, 15.4.3.2, 16.4.3.2, 17.4.3.2, 18.4.3.2, 19.4.3.2, 20.4.3.2, 21.4.3.2, 22.4.3.2, 23.4.3.2, 24.4.3.2, 25.4.3.2, 26.4.3.2, 27.4.3.2, 28.4.3.2, 29.4.3.2, 30.4.3.2, 31.4.3.2, 32.4.3.2, 33.4.3.2, 34.4.3.2, 35.4.3.2, 36.4.3.2, 37.4.3.2, 38.4.3.2,
- 15 39.4.3.2, 40.4.3.2, 1.5.3.2, 2.5.3.2, 3.5.3.2, 4.5.3.2, 5.5.3.2, 6.5.3.2, 7.5.3.2, 8.5.3.2, 9.5.3.2, 10.5.3.2, 11.5.3.2, 12.5.3.2, 13.5.3.2, 14.5.3.2, 15.5.3.2, 16.5.3.2, 17.5.3.2, 18.5.3.2, 19.5.3.2, 20.5.3.2, 21.5.3.2, 22.5.3.2, 23.5.3.2, 24.5.3.2, 25.5.3.2, 26.5.3.2, 27.5.3.2, 28.5.3.2, 29.5.3.2, 30.5.3.2, 31.5.3.2, 32.5.3.2, 33.5.3.2, 34.5.3.2, 35.5.3.2, 36.5.3.2, 37.5.3.2, 38.5.3.2, 39.5.3.2, 40.5.3.2, 1.6.3.2, 2.6.3.2, 3.6.3.2, 4.6.3.2, 5.6.3.2,
- 20 6.6.3.2, 7.6.3.2, 8.6.3.2, 9.6.3.2, 10.6.3.2, 11.6.3.2, 12.6.3.2, 13.6.3.2, 14.6.3.2, 15.6.3.2, 16.6.3.2, 17.6.3.2, 18.6.3.2, 19.6.3.2, 20.6.3.2, 21.6.3.2, 22.6.3.2, 23.6.3.2, 24.6.3.2, 25.6.3.2, 26.6.3.2, 27.6.3.2, 28.6.3.2, 29.6.3.2, 30.6.3.2, 31.6.3.2, 32.6.3.2, 33.6.3.2, 34.6.3.2, 35.6.3.2, 36.6.3.2, 37.6.3.2, 38.6.3.2, 39.6.3.2, 40.6.3.2, 1.7.3.2, 2.7.3.2, 3.7.3.2, 4.7.3.2, 5.7.3.2, 6.7.3.2, 7.7.3.2, 8.7.3.2, 9.7.3.2, 10.7.3.2, 11.7.3.2, 12.7.3.2, 13.7.3.2,
- 25 14.7.3.2, 15.7.3.2, 16.7.3.2, 17.7.3.2, 18.7.3.2, 19.7.3.2, 20.7.3.2, 21.7.3.2, 22.7.3.2, 23.7.3.2, 24.7.3.2, 25.7.3.2, 26.7.3.2, 27.7.3.2, 28.7.3.2, 29.7.3.2, 30.7.3.2, 31.7.3.2, 32.7.3.2, 33.7.3.2, 34.7.3.2, 35.7.3.2, 36.7.3.2, 37.7.3.2, 38.7.3.2, 39.7.3.2, 40.7.3.2, 1.8.3.2, 2.8.3.2, 3.8.3.2, 4.8.3.2, 5.8.3.2, 6.8.3.2, 7.8.3.2, 8.8.3.2, 9.8.3.2, 10.8.3.2, 11.8.3.2, 12.8.3.2, 13.8.3.2, 14.8.3.2, 15.8.3.2, 16.8.3.2, 17.8.3.2, 18.8.3.2, 19.8.3.2,
- 30 20.8.3.2, 21.8.3.2, 22.8.3.2, 23.8.3.2, 24.8.3.2, 25.8.3.2, 26.8.3.2, 27.8.3.2, 28.8.3.2, 29.8.3.2, 30.8.3.2, 31.8.3.2, 32.8.3.2, 33.8.3.2, 34.8.3.2, 35.8.3.2, 36.8.3.2, 37.8.3.2, 38.8.3.2, 39.8.3.2, 40.8.3.2, 1.9.3.2, 2.9.3.2, 3.9.3.2, 4.9.3.2, 5.9.3.2, 6.9.3.2, 7.9.3.2, 8.9.3.2, 9.9.3.2, 10.9.3.2, 11.9.3.2, 12.9.3.2, 13.9.3.2, 14.9.3.2, 15.9.3.2, 16.9.3.2, 17.9.3.2, 18.9.3.2, 19.9.3.2, 20.9.3.2, 21.9.3.2, 22.9.3.2, 23.9.3.2, 24.9.3.2, 25.9.3.2,
- 35 26.9.3.2, 27.9.3.2, 28.9.3.2, 29.9.3.2, 30.9.3.2, 31.9.3.2, 32.9.3.2, 33.9.3.2, 34.9.3.2,

35.9.3.2, 36.9.3.2, 37.9.3.2, 38.9.3.2, 39.9.3.2, 40.9.3.2, 1.10.3.2, 2.10.3.2, 3.10.3.2,  
4.10.3.2, 5.10.3.2, 6.10.3.2, 7.10.3.2, 8.10.3.2, 9.10.3.2, 10.10.3.2, 11.10.3.2, 12.10.3.2,  
13.10.3.2, 14.10.3.2, 15.10.3.2, 16.10.3.2, 17.10.3.2, 18.10.3.2, 19.10.3.2, 20.10.3.2,  
21.10.3.2, 22.10.3.2, 23.10.3.2, 24.10.3.2, 25.10.3.2, 26.10.3.2, 27.10.3.2, 28.10.3.2,  
5 29.10.3.2, 30.10.3.2, 31.10.3.2, 32.10.3.2, 33.10.3.2, 34.10.3.2, 35.10.3.2, 36.10.3.2,  
37.10.3.2, 38.10.3.2, 39.10.3.2, 40.10.3.2, 1.11.3.2, 2.11.3.2, 3.11.3.2, 4.11.3.2, 5.11.3.2,  
6.11.3.2, 7.11.3.2, 8.11.3.2, 9.11.3.2, 10.11.3.2, 11.11.3.2, 12.11.3.2, 13.11.3.2, 14.11.3.2,  
15.11.3.2, 16.11.3.2, 17.11.3.2, 18.11.3.2, 19.11.3.2, 20.11.3.2, 21.11.3.2, 22.11.3.2,  
23.11.3.2, 24.11.3.2, 25.11.3.2, 26.11.3.2, 27.11.3.2, 28.11.3.2, 29.11.3.2, 30.11.3.2,  
10 31.11.3.2, 32.11.3.2, 33.11.3.2, 34.11.3.2, 35.11.3.2, 36.11.3.2, 37.11.3.2, 38.11.3.2,  
39.11.3.2, 40.11.3.2, 1.12.3.2, 2.12.3.2, 3.12.3.2, 4.12.3.2, 5.12.3.2, 6.12.3.2, 7.12.3.2,  
8.12.3.2, 9.12.3.2, 10.12.3.2, 11.12.3.2, 12.12.3.2, 13.12.3.2, 14.12.3.2, 15.12.3.2,  
16.12.3.2, 17.12.3.2, 18.12.3.2, 19.12.3.2, 20.12.3.2, 21.12.3.2, 22.12.3.2, 23.12.3.2,  
24.12.3.2, 25.12.3.2, 26.12.3.2, 27.12.3.2, 28.12.3.2, 29.12.3.2, 30.12.3.2, 31.12.3.2,  
15 32.12.3.2, 33.12.3.2, 34.12.3.2, 35.12.3.2, 36.12.3.2, 37.12.3.2, 38.12.3.2, 39.12.3.2,  
40.12.3.2, 1.13.3.2, 2.13.3.2, 3.13.3.2, 4.13.3.2, 5.13.3.2, 6.13.3.2, 7.13.3.2, 8.13.3.2,  
9.13.3.2, 10.13.3.2, 11.13.3.2, 12.13.3.2, 13.13.3.2, 14.13.3.2, 15.13.3.2, 16.13.3.2,  
17.13.3.2, 18.13.3.2, 19.13.3.2, 20.13.3.2, 21.13.3.2, 22.13.3.2, 23.13.3.2, 24.13.3.2,  
25.13.3.2, 26.13.3.2, 27.13.3.2, 28.13.3.2, 29.13.3.2, 30.13.3.2, 31.13.3.2, 32.13.3.2,  
20 33.13.3.2, 34.13.3.2, 35.13.3.2, 36.13.3.2, 37.13.3.2, 38.13.3.2, 39.13.3.2, 40.13.3.2,  
1.14.3.2, 2.14.3.2, 3.14.3.2, 4.14.3.2, 5.14.3.2, 6.14.3.2, 7.14.3.2, 8.14.3.2, 9.14.3.2,  
10.14.3.2, 11.14.3.2, 12.14.3.2, 13.14.3.2, 14.14.3.2, 15.14.3.2, 16.14.3.2, 17.14.3.2,  
18.14.3.2, 19.14.3.2, 20.14.3.2, 21.14.3.2, 22.14.3.2, 23.14.3.2, 24.14.3.2, 25.14.3.2,  
26.14.3.2, 27.14.3.2, 28.14.3.2, 29.14.3.2, 30.14.3.2, 31.14.3.2, 32.14.3.2, 33.14.3.2,  
25 34.14.3.2, 35.14.3.2, 36.14.3.2, 37.14.3.2, 38.14.3.2, 39.14.3.2, 40.14.3.2, 1.15.3.2,  
2.15.3.2, 3.15.3.2, 4.15.3.2, 5.15.3.2, 6.15.3.2, 7.15.3.2, 8.15.3.2, 9.15.3.2, 10.15.3.2,  
11.15.3.2, 12.15.3.2, 13.15.3.2, 14.15.3.2, 15.15.3.2, 16.15.3.2, 17.15.3.2, 18.15.3.2,  
19.15.3.2, 20.15.3.2, 21.15.3.2, 22.15.3.2, 23.15.3.2, 24.15.3.2, 25.15.3.2, 26.15.3.2,  
27.15.3.2, 28.15.3.2, 29.15.3.2, 30.15.3.2, 31.15.3.2, 32.15.3.2, 33.15.3.2, 34.15.3.2,  
30 35.15.3.2, 36.15.3.2, 37.15.3.2, 38.15.3.2, 39.15.3.2, 40.15.3.2, 1.16.3.2, 2.16.3.2,  
3.16.3.2, 4.16.3.2, 5.16.3.2, 6.16.3.2, 7.16.3.2, 8.16.3.2, 9.16.3.2, 10.16.3.2, 11.16.3.2,  
12.16.3.2, 13.16.3.2, 14.16.3.2, 15.16.3.2, 16.16.3.2, 17.16.3.2, 18.16.3.2, 19.16.3.2,  
20.16.3.2, 21.16.3.2, 22.16.3.2, 23.16.3.2, 24.16.3.2, 25.16.3.2, 26.16.3.2, 27.16.3.2,  
28.16.3.2, 29.16.3.2, 30.16.3.2, 31.16.3.2, 32.16.3.2, 33.16.3.2, 34.16.3.2, 35.16.3.2,  
35 36.16.3.2, 37.16.3.2, 38.16.3.2, 39.16.3.2, 40.16.3.2, 1.17.3.2, 2.17.3.2, 3.17.3.2, 4.17.3.2,

5.17.3.2, 6.17.3.2, 7.17.3.2, 8.17.3.2, 9.17.3.2, 10.17.3.2, 11.17.3.2, 12.17.3.2, 13.17.3.2,  
14.17.3.2, 15.17.3.2, 16.17.3.2, 17.17.3.2, 18.17.3.2, 19.17.3.2, 20.17.3.2, 21.17.3.2,  
22.17.3.2, 23.17.3.2, 24.17.3.2, 25.17.3.2, 26.17.3.2, 27.17.3.2, 28.17.3.2, 29.17.3.2,  
30.17.3.2, 31.17.3.2, 32.17.3.2, 33.17.3.2, 34.17.3.2, 35.17.3.2, 36.17.3.2, 37.17.3.2,  
5 38.17.3.2, 39.17.3.2, 40.17.3.2, 1.18.3.2, 2.18.3.2, 3.18.3.2, 4.18.3.2, 5.18.3.2, 6.18.3.2,  
7.18.3.2, 8.18.3.2, 9.18.3.2, 10.18.3.2, 11.18.3.2, 12.18.3.2, 13.18.3.2, 14.18.3.2,  
15.18.3.2, 16.18.3.2, 17.18.3.2, 18.18.3.2, 19.18.3.2, 20.18.3.2, 21.18.3.2, 22.18.3.2,  
23.18.3.2, 24.18.3.2, 25.18.3.2, 26.18.3.2, 27.18.3.2, 28.18.3.2, 29.18.3.2, 30.18.3.2,  
31.18.3.2, 32.18.3.2, 33.18.3.2, 34.18.3.2, 35.18.3.2, 36.18.3.2, 37.18.3.2, 38.18.3.2,  
10 39.18.3.2, 40.18.3.2, 1.19.3.2, 2.19.3.2, 3.19.3.2, 4.19.3.2, 5.19.3.2, 6.19.3.2, 7.19.3.2,  
8.19.3.2, 9.19.3.2, 10.19.3.2, 11.19.3.2, 12.19.3.2, 13.19.3.2, 14.19.3.2, 15.19.3.2,  
16.19.3.2, 17.19.3.2, 18.19.3.2, 19.19.3.2, 20.19.3.2, 21.19.3.2, 22.19.3.2, 23.19.3.2,  
24.19.3.2, 25.19.3.2, 26.19.3.2, 27.19.3.2, 28.19.3.2, 29.19.3.2, 30.19.3.2, 31.19.3.2,  
32.19.3.2, 33.19.3.2, 34.19.3.2, 35.19.3.2, 36.19.3.2, 37.19.3.2, 38.19.3.2, 39.19.3.2,  
15 40.19.3.2, 1.20.3.2, 2.20.3.2, 3.20.3.2, 4.20.3.2, 5.20.3.2, 6.20.3.2, 7.20.3.2, 8.20.3.2,  
9.20.3.2, 10.20.3.2, 11.20.3.2, 12.20.3.2, 13.20.3.2, 14.20.3.2, 15.20.3.2, 16.20.3.2,  
17.20.3.2, 18.20.3.2, 19.20.3.2, 20.20.3.2, 21.20.3.2, 22.20.3.2, 23.20.3.2, 24.20.3.2,  
25.20.3.2, 26.20.3.2, 27.20.3.2, 28.20.3.2, 29.20.3.2, 30.20.3.2, 31.20.3.2, 32.20.3.2,  
33.20.3.2, 34.20.3.2, 35.20.3.2, 36.20.3.2, 37.20.3.2, 38.20.3.2, 39.20.3.2, 40.20.3.2,  
20 1.21.3.2, 2.21.3.2, 3.21.3.2, 4.21.3.2, 5.21.3.2, 6.21.3.2, 7.21.3.2, 8.21.3.2, 9.21.3.2,  
10.21.3.2, 11.21.3.2, 12.21.3.2, 13.21.3.2, 14.21.3.2, 15.21.3.2, 16.21.3.2, 17.21.3.2,  
18.21.3.2, 19.21.3.2, 20.21.3.2, 21.21.3.2, 22.21.3.2, 23.21.3.2, 24.21.3.2, 25.21.3.2,  
26.21.3.2, 27.21.3.2, 28.21.3.2, 29.21.3.2, 30.21.3.2, 31.21.3.2, 32.21.3.2, 33.21.3.2,  
34.21.3.2, 35.21.3.2, 36.21.3.2, 37.21.3.2, 38.21.3.2, 39.21.3.2, 40.21.3.2, 1.22.3.2,  
25 2.22.3.2, 3.22.3.2, 4.22.3.2, 5.22.3.2, 6.22.3.2, 7.22.3.2, 8.22.3.2, 9.22.3.2, 10.22.3.2,  
11.22.3.2, 12.22.3.2, 13.22.3.2, 14.22.3.2, 15.22.3.2, 16.22.3.2, 17.22.3.2, 18.22.3.2,  
19.22.3.2, 20.22.3.2, 21.22.3.2, 22.22.3.2, 23.22.3.2, 24.22.3.2, 25.22.3.2, 26.22.3.2,  
27.22.3.2, 28.22.3.2, 29.22.3.2, 30.22.3.2, 31.22.3.2, 32.22.3.2, 33.22.3.2, 34.22.3.2,  
35.22.3.2, 36.22.3.2, 37.22.3.2, 38.22.3.2, 39.22.3.2, 40.22.3.2, 1.23.3.2, 2.23.3.2,  
30 3.23.3.2, 4.23.3.2, 5.23.3.2, 6.23.3.2, 7.23.3.2, 8.23.3.2, 9.23.3.2, 10.23.3.2, 11.23.3.2,  
12.23.3.2, 13.23.3.2, 14.23.3.2, 15.23.3.2, 16.23.3.2, 17.23.3.2, 18.23.3.2, 19.23.3.2,  
20.23.3.2, 21.23.3.2, 22.23.3.2, 23.23.3.2, 24.23.3.2, 25.23.3.2, 26.23.3.2, 27.23.3.2,  
28.23.3.2, 29.23.3.2, 30.23.3.2, 31.23.3.2, 32.23.3.2, 33.23.3.2, 34.23.3.2, 35.23.3.2,  
36.23.3.2, 37.23.3.2, 38.23.3.2, 39.23.3.2, 40.23.3.2, 1.24.3.2, 2.24.3.2, 3.24.3.2, 4.24.3.2,  
35 5.24.3.2, 6.24.3.2, 7.24.3.2, 8.24.3.2, 9.24.3.2, 10.24.3.2, 11.24.3.2, 12.24.3.2, 13.24.3.2,

- 14.24.3.2, 15.24.3.2, 16.24.3.2, 17.24.3.2, 18.24.3.2, 19.24.3.2, 20.24.3.2, 21.24.3.2, 22.24.3.2, 23.24.3.2, 24.24.3.2, 25.24.3.2, 26.24.3.2, 27.24.3.2, 28.24.3.2, 29.24.3.2, 30.24.3.2, 31.24.3.2, 32.24.3.2, 33.24.3.2, 34.24.3.2, 35.24.3.2, 36.24.3.2, 37.24.3.2, 38.24.3.2, 39.24.3.2, 40.24.3.2, 1.25.3.2, 2.25.3.2, 3.25.3.2, 4.25.3.2, 5.25.3.2, 6.25.3.2, 7.25.3.2, 8.25.3.2, 9.25.3.2, 10.25.3.2, 11.25.3.2, 12.25.3.2, 13.25.3.2, 14.25.3.2, 15.25.3.2, 16.25.3.2, 17.25.3.2, 18.25.3.2, 19.25.3.2, 20.25.3.2, 21.25.3.2, 22.25.3.2, 23.25.3.2, 24.25.3.2, 25.25.3.2, 26.25.3.2, 27.25.3.2, 28.25.3.2, 29.25.3.2, 30.25.3.2, 31.25.3.2, 32.25.3.2, 33.25.3.2, 34.25.3.2, 35.25.3.2, 36.25.3.2, 37.25.3.2, 38.25.3.2, 39.25.3.2, 40.25.3.2, 1.26.3.2, 2.26.3.2, 3.26.3.2, 4.26.3.2, 5.26.3.2, 6.26.3.2, 7.26.3.2, 8.26.3.2, 9.26.3.2, 10.26.3.2, 11.26.3.2, 12.26.3.2, 13.26.3.2, 14.26.3.2, 15.26.3.2, 16.26.3.2, 17.26.3.2, 18.26.3.2, 19.26.3.2, 20.26.3.2, 21.26.3.2, 22.26.3.2, 23.26.3.2, 24.26.3.2, 25.26.3.2, 26.26.3.2, 27.26.3.2, 28.26.3.2, 29.26.3.2, 30.26.3.2, 31.26.3.2, 32.26.3.2, 33.26.3.2, 34.26.3.2, 35.26.3.2, 36.26.3.2, 37.26.3.2, 38.26.3.2, 39.26.3.2, 40.26.3.2, 1.27.3.2, 2.27.3.2, 3.27.3.2, 4.27.3.2, 5.27.3.2, 6.27.3.2, 7.27.3.2, 8.27.3.2, 9.27.3.2, 10.27.3.2, 11.27.3.2, 12.27.3.2, 13.27.3.2, 14.27.3.2, 15.27.3.2, 16.27.3.2, 17.27.3.2, 18.27.3.2, 19.27.3.2, 20.27.3.2, 21.27.3.2, 22.27.3.2, 23.27.3.2, 24.27.3.2, 25.27.3.2, 26.27.3.2, 27.27.3.2, 28.27.3.2, 29.27.3.2, 30.27.3.2, 31.27.3.2, 32.27.3.2, 33.27.3.2, 34.27.3.2, 35.27.3.2, 36.27.3.2, 37.27.3.2, 38.27.3.2, 39.27.3.2, 40.27.3.2, 1.28.3.2, 2.28.3.2, 3.28.3.2, 4.28.3.2, 5.28.3.2, 6.28.3.2, 7.28.3.2, 8.28.3.2, 9.28.3.2, 10.28.3.2, 11.28.3.2, 12.28.3.2, 13.28.3.2, 14.28.3.2, 15.28.3.2, 16.28.3.2, 17.28.3.2, 18.28.3.2, 19.28.3.2, 20.28.3.2, 21.28.3.2, 22.28.3.2, 23.28.3.2, 24.28.3.2, 25.28.3.2, 26.28.3.2, 27.28.3.2, 28.28.3.2, 29.28.3.2, 30.28.3.2, 31.28.3.2, 32.28.3.2, 33.28.3.2, 34.28.3.2, 35.28.3.2, 36.28.3.2, 37.28.3.2, 38.28.3.2, 39.28.3.2, 40.28.3.2, 1.29.3.2, 2.29.3.2, 3.29.3.2, 4.29.3.2, 5.29.3.2, 6.29.3.2, 7.29.3.2, 8.29.3.2, 9.29.3.2, 10.29.3.2, 11.29.3.2, 12.29.3.2, 13.29.3.2, 14.29.3.2, 15.29.3.2, 16.29.3.2, 17.29.3.2, 18.29.3.2, 19.29.3.2, 20.29.3.2, 21.29.3.2, 22.29.3.2, 23.29.3.2, 24.29.3.2, 25.29.3.2, 26.29.3.2, 27.29.3.2, 28.29.3.2, 29.29.3.2, 30.29.3.2, 31.29.3.2, 32.29.3.2, 33.29.3.2, 34.29.3.2, 35.29.3.2, 36.29.3.2, 37.29.3.2, 38.29.3.2, 39.29.3.2, 40.29.3.2, 1.1.3.4, 2.1.3.4, 3.1.3.4, 4.1.3.4, 5.1.3.4, 6.1.3.4, 7.1.3.4, 8.1.3.4, 9.1.3.4, 10.1.3.4, 11.1.3.4, 12.1.3.4, 13.1.3.4, 14.1.3.4, 15.1.3.4, 16.1.3.4, 17.1.3.4, 18.1.3.4, 19.1.3.4, 20.1.3.4, 21.1.3.4, 22.1.3.4, 23.1.3.4, 24.1.3.4, 25.1.3.4, 26.1.3.4, 27.1.3.4, 28.1.3.4, 29.1.3.4, 30.1.3.4, 31.1.3.4, 32.1.3.4, 33.1.3.4, 34.1.3.4, 35.1.3.4, 36.1.3.4, 37.1.3.4, 38.1.3.4, 39.1.3.4, 40.1.3.4, 1.2.3.4, 2.2.3.4, 3.2.3.4, 4.2.3.4, 5.2.3.4, 6.2.3.4, 7.2.3.4, 8.2.3.4, 9.2.3.4, 10.2.3.4, 11.2.3.4, 12.2.3.4, 13.2.3.4, 14.2.3.4, 15.2.3.4, 16.2.3.4, 17.2.3.4, 18.2.3.4, 19.2.3.4, 20.2.3.4, 21.2.3.4, 22.2.3.4, 23.2.3.4, 24.2.3.4, 25.2.3.4, 26.2.3.4, 27.2.3.4, 28.2.3.4,

29.2.3.4, 30.2.3.4, 31.2.3.4, 32.2.3.4, 33.2.3.4, 34.2.3.4, 35.2.3.4, 36.2.3.4, 37.2.3.4,  
38.2.3.4, 39.2.3.4, 40.2.3.4, 1.3.3.4, 2.3.3.4, 3.3.3.4, 4.3.3.4, 5.3.3.4, 6.3.3.4, 7.3.3.4,  
8.3.3.4, 9.3.3.4, 10.3.3.4, 11.3.3.4, 12.3.3.4, 13.3.3.4, 14.3.3.4, 15.3.3.4, 16.3.3.4,  
17.3.3.4, 18.3.3.4, 19.3.3.4, 20.3.3.4, 21.3.3.4, 22.3.3.4, 23.3.3.4, 24.3.3.4, 25.3.3.4,  
5 26.3.3.4, 27.3.3.4, 28.3.3.4, 29.3.3.4, 30.3.3.4, 31.3.3.4, 32.3.3.4, 33.3.3.4, 34.3.3.4,  
35.3.3.4, 36.3.3.4, 37.3.3.4, 38.3.3.4, 39.3.3.4, 40.3.3.4, 1.4.3.4, 2.4.3.4, 3.4.3.4, 4.4.3.4,  
5.4.3.4, 6.4.3.4, 7.4.3.4, 8.4.3.4, 9.4.3.4, 10.4.3.4, 11.4.3.4, 12.4.3.4, 13.4.3.4, 14.4.3.4,  
15.4.3.4, 16.4.3.4, 17.4.3.4, 18.4.3.4, 19.4.3.4, 20.4.3.4, 21.4.3.4, 22.4.3.4, 23.4.3.4,  
24.4.3.4, 25.4.3.4, 26.4.3.4, 27.4.3.4, 28.4.3.4, 29.4.3.4, 30.4.3.4, 31.4.3.4, 32.4.3.4,  
10 33.4.3.4, 34.4.3.4, 35.4.3.4, 36.4.3.4, 37.4.3.4, 38.4.3.4, 39.4.3.4, 40.4.3.4, 1.5.3.4,  
2.5.3.4, 3.5.3.4, 4.5.3.4, 5.5.3.4, 6.5.3.4, 7.5.3.4, 8.5.3.4, 9.5.3.4, 10.5.3.4, 11.5.3.4,  
12.5.3.4, 13.5.3.4, 14.5.3.4, 15.5.3.4, 16.5.3.4, 17.5.3.4, 18.5.3.4, 19.5.3.4, 20.5.3.4,  
21.5.3.4, 22.5.3.4, 23.5.3.4, 24.5.3.4, 25.5.3.4, 26.5.3.4, 27.5.3.4, 28.5.3.4, 29.5.3.4,  
30.5.3.4, 31.5.3.4, 32.5.3.4, 33.5.3.4, 34.5.3.4, 35.5.3.4, 36.5.3.4, 37.5.3.4, 38.5.3.4,  
15 39.5.3.4, 40.5.3.4, 1.6.3.4, 2.6.3.4, 3.6.3.4, 4.6.3.4, 5.6.3.4, 6.6.3.4, 7.6.3.4, 8.6.3.4,  
9.6.3.4, 10.6.3.4, 11.6.3.4, 12.6.3.4, 13.6.3.4, 14.6.3.4, 15.6.3.4, 16.6.3.4, 17.6.3.4,  
18.6.3.4, 19.6.3.4, 20.6.3.4, 21.6.3.4, 22.6.3.4, 23.6.3.4, 24.6.3.4, 25.6.3.4, 26.6.3.4,  
27.6.3.4, 28.6.3.4, 29.6.3.4, 30.6.3.4, 31.6.3.4, 32.6.3.4, 33.6.3.4, 34.6.3.4, 35.6.3.4,  
36.6.3.4, 37.6.3.4, 38.6.3.4, 39.6.3.4, 40.6.3.4, 1.7.3.4, 2.7.3.4, 3.7.3.4, 4.7.3.4, 5.7.3.4,  
20 6.7.3.4, 7.7.3.4, 8.7.3.4, 9.7.3.4, 10.7.3.4, 11.7.3.4, 12.7.3.4, 13.7.3.4, 14.7.3.4, 15.7.3.4,  
16.7.3.4, 17.7.3.4, 18.7.3.4, 19.7.3.4, 20.7.3.4, 21.7.3.4, 22.7.3.4, 23.7.3.4, 24.7.3.4,  
25.7.3.4, 26.7.3.4, 27.7.3.4, 28.7.3.4, 29.7.3.4, 30.7.3.4, 31.7.3.4, 32.7.3.4, 33.7.3.4,  
34.7.3.4, 35.7.3.4, 36.7.3.4, 37.7.3.4, 38.7.3.4, 39.7.3.4, 40.7.3.4, 1.8.3.4, 2.8.3.4, 3.8.3.4,  
4.8.3.4, 5.8.3.4, 6.8.3.4, 7.8.3.4, 8.8.3.4, 9.8.3.4, 10.8.3.4, 11.8.3.4, 12.8.3.4, 13.8.3.4,  
25 14.8.3.4, 15.8.3.4, 16.8.3.4, 17.8.3.4, 18.8.3.4, 19.8.3.4, 20.8.3.4, 21.8.3.4, 22.8.3.4,  
23.8.3.4, 24.8.3.4, 25.8.3.4, 26.8.3.4, 27.8.3.4, 28.8.3.4, 29.8.3.4, 30.8.3.4, 31.8.3.4,  
32.8.3.4, 33.8.3.4, 34.8.3.4, 35.8.3.4, 36.8.3.4, 37.8.3.4, 38.8.3.4, 39.8.3.4, 40.8.3.4,  
1.9.3.4, 2.9.3.4, 3.9.3.4, 4.9.3.4, 5.9.3.4, 6.9.3.4, 7.9.3.4, 8.9.3.4, 9.9.3.4, 10.9.3.4,  
11.9.3.4, 12.9.3.4, 13.9.3.4, 14.9.3.4, 15.9.3.4, 16.9.3.4, 17.9.3.4, 18.9.3.4, 19.9.3.4,  
30 20.9.3.4, 21.9.3.4, 22.9.3.4, 23.9.3.4, 24.9.3.4, 25.9.3.4, 26.9.3.4, 27.9.3.4, 28.9.3.4,  
29.9.3.4, 30.9.3.4, 31.9.3.4, 32.9.3.4, 33.9.3.4, 34.9.3.4, 35.9.3.4, 36.9.3.4, 37.9.3.4,  
38.9.3.4, 39.9.3.4, 40.9.3.4, 1.10.3.4, 2.10.3.4, 3.10.3.4, 4.10.3.4, 5.10.3.4, 6.10.3.4,  
7.10.3.4, 8.10.3.4, 9.10.3.4, 10.10.3.4, 11.10.3.4, 12.10.3.4, 13.10.3.4, 14.10.3.4,  
15.10.3.4, 16.10.3.4, 17.10.3.4, 18.10.3.4, 19.10.3.4, 20.10.3.4, 21.10.3.4, 22.10.3.4,  
35 23.10.3.4, 24.10.3.4, 25.10.3.4, 26.10.3.4, 27.10.3.4, 28.10.3.4, 29.10.3.4, 30.10.3.4,

31.10.3.4, 32.10.3.4, 33.10.3.4, 34.10.3.4, 35.10.3.4, 36.10.3.4, 37.10.3.4, 38.10.3.4,  
 39.10.3.4, 40.10.3.4, 1.11.3.4, 2.11.3.4, 3.11.3.4, 4.11.3.4, 5.11.3.4, 6.11.3.4, 7.11.3.4,  
 8.11.3.4, 9.11.3.4, 10.11.3.4, 11.11.3.4, 12.11.3.4, 13.11.3.4, 14.11.3.4, 15.11.3.4,  
 16.11.3.4, 17.11.3.4, 18.11.3.4, 19.11.3.4, 20.11.3.4, 21.11.3.4, 22.11.3.4, 23.11.3.4,  
 5 24.11.3.4, 25.11.3.4, 26.11.3.4, 27.11.3.4, 28.11.3.4, 29.11.3.4, 30.11.3.4, 31.11.3.4,  
 32.11.3.4, 33.11.3.4, 34.11.3.4, 35.11.3.4, 36.11.3.4, 37.11.3.4, 38.11.3.4, 39.11.3.4,  
 40.11.3.4, 1.12.3.4, 2.12.3.4, 3.12.3.4, 4.12.3.4, 5.12.3.4, 6.12.3.4, 7.12.3.4, 8.12.3.4,  
 9.12.3.4, 10.12.3.4, 11.12.3.4, 12.12.3.4, 13.12.3.4, 14.12.3.4, 15.12.3.4, 16.12.3.4,  
 17.12.3.4, 18.12.3.4, 19.12.3.4, 20.12.3.4, 21.12.3.4, 22.12.3.4, 23.12.3.4, 24.12.3.4,  
 10 25.12.3.4, 26.12.3.4, 27.12.3.4, 28.12.3.4, 29.12.3.4, 30.12.3.4, 31.12.3.4, 32.12.3.4,  
 33.12.3.4, 34.12.3.4, 35.12.3.4, 36.12.3.4, 37.12.3.4, 38.12.3.4, 39.12.3.4, 40.12.3.4,  
 1.13.3.4, 2.13.3.4, 3.13.3.4, 4.13.3.4, 5.13.3.4, 6.13.3.4, 7.13.3.4, 8.13.3.4, 9.13.3.4,  
 10.13.3.4, 11.13.3.4, 12.13.3.4, 13.13.3.4, 14.13.3.4, 15.13.3.4, 16.13.3.4, 17.13.3.4,  
 18.13.3.4, 19.13.3.4, 20.13.3.4, 21.13.3.4, 22.13.3.4, 23.13.3.4, 24.13.3.4, 25.13.3.4,  
 15 26.13.3.4, 27.13.3.4, 28.13.3.4, 29.13.3.4, 30.13.3.4, 31.13.3.4, 32.13.3.4, 33.13.3.4,  
 34.13.3.4, 35.13.3.4, 36.13.3.4, 37.13.3.4, 38.13.3.4, 39.13.3.4, 40.13.3.4, 1.14.3.4,  
 2.14.3.4, 3.14.3.4, 4.14.3.4, 5.14.3.4, 6.14.3.4, 7.14.3.4, 8.14.3.4, 9.14.3.4, 10.14.3.4,  
 11.14.3.4, 12.14.3.4, 13.14.3.4, 14.14.3.4, 15.14.3.4, 16.14.3.4, 17.14.3.4, 18.14.3.4,  
 19.14.3.4, 20.14.3.4, 21.14.3.4, 22.14.3.4, 23.14.3.4, 24.14.3.4, 25.14.3.4, 26.14.3.4,  
 20 27.14.3.4, 28.14.3.4, 29.14.3.4, 30.14.3.4, 31.14.3.4, 32.14.3.4, 33.14.3.4, 34.14.3.4,  
 35.14.3.4, 36.14.3.4, 37.14.3.4, 38.14.3.4, 39.14.3.4, 40.14.3.4, 1.15.3.4, 2.15.3.4,  
 3.15.3.4, 4.15.3.4, 5.15.3.4, 6.15.3.4, 7.15.3.4, 8.15.3.4, 9.15.3.4, 10.15.3.4, 11.15.3.4,  
 12.15.3.4, 13.15.3.4, 14.15.3.4, 15.15.3.4, 16.15.3.4, 17.15.3.4, 18.15.3.4, 19.15.3.4,  
 20.15.3.4, 21.15.3.4, 22.15.3.4, 23.15.3.4, 24.15.3.4, 25.15.3.4, 26.15.3.4, 27.15.3.4,  
 25 28.15.3.4, 29.15.3.4, 30.15.3.4, 31.15.3.4, 32.15.3.4, 33.15.3.4, 34.15.3.4, 35.15.3.4,  
 36.15.3.4, 37.15.3.4, 38.15.3.4, 39.15.3.4, 40.15.3.4, 1.16.3.4, 2.16.3.4, 3.16.3.4, 4.16.3.4,  
 5.16.3.4, 6.16.3.4, 7.16.3.4, 8.16.3.4, 9.16.3.4, 10.16.3.4, 11.16.3.4, 12.16.3.4, 13.16.3.4,  
 14.16.3.4, 15.16.3.4, 16.16.3.4, 17.16.3.4, 18.16.3.4, 19.16.3.4, 20.16.3.4, 21.16.3.4,  
 22.16.3.4, 23.16.3.4, 24.16.3.4, 25.16.3.4, 26.16.3.4, 27.16.3.4, 28.16.3.4, 29.16.3.4,  
 30 30.16.3.4, 31.16.3.4, 32.16.3.4, 33.16.3.4, 34.16.3.4, 35.16.3.4, 36.16.3.4, 37.16.3.4,  
 38.16.3.4, 39.16.3.4, 40.16.3.4, 1.17.3.4, 2.17.3.4, 3.17.3.4, 4.17.3.4, 5.17.3.4, 6.17.3.4,  
 7.17.3.4, 8.17.3.4, 9.17.3.4, 10.17.3.4, 11.17.3.4, 12.17.3.4, 13.17.3.4, 14.17.3.4,  
 15.17.3.4, 16.17.3.4, 17.17.3.4, 18.17.3.4, 19.17.3.4, 20.17.3.4, 21.17.3.4, 22.17.3.4,  
 23.17.3.4, 24.17.3.4, 25.17.3.4, 26.17.3.4, 27.17.3.4, 28.17.3.4, 29.17.3.4, 30.17.3.4,  
 35 31.17.3.4, 32.17.3.4, 33.17.3.4, 34.17.3.4, 35.17.3.4, 36.17.3.4, 37.17.3.4, 38.17.3.4,

39.17.3.4, 40.17.3.4, 1.18.3.4, 2.18.3.4, 3.18.3.4, 4.18.3.4, 5.18.3.4, 6.18.3.4, 7.18.3.4,  
 8.18.3.4, 9.18.3.4, 10.18.3.4, 11.18.3.4, 12.18.3.4, 13.18.3.4, 14.18.3.4, 15.18.3.4,  
 16.18.3.4, 17.18.3.4, 18.18.3.4, 19.18.3.4, 20.18.3.4, 21.18.3.4, 22.18.3.4, 23.18.3.4,  
 24.18.3.4, 25.18.3.4, 26.18.3.4, 27.18.3.4, 28.18.3.4, 29.18.3.4, 30.18.3.4, 31.18.3.4,  
 5 32.18.3.4, 33.18.3.4, 34.18.3.4, 35.18.3.4, 36.18.3.4, 37.18.3.4, 38.18.3.4, 39.18.3.4,  
 40.18.3.4, 1.19.3.4, 2.19.3.4, 3.19.3.4, 4.19.3.4, 5.19.3.4, 6.19.3.4, 7.19.3.4, 8.19.3.4,  
 9.19.3.4, 10.19.3.4, 11.19.3.4, 12.19.3.4, 13.19.3.4, 14.19.3.4, 15.19.3.4, 16.19.3.4,  
 17.19.3.4, 18.19.3.4, 19.19.3.4, 20.19.3.4, 21.19.3.4, 22.19.3.4, 23.19.3.4, 24.19.3.4,  
 25.19.3.4, 26.19.3.4, 27.19.3.4, 28.19.3.4, 29.19.3.4, 30.19.3.4, 31.19.3.4, 32.19.3.4,  
 10 33.19.3.4, 34.19.3.4, 35.19.3.4, 36.19.3.4, 37.19.3.4, 38.19.3.4, 39.19.3.4, 40.19.3.4,  
 1.20.3.4, 2.20.3.4, 3.20.3.4, 4.20.3.4, 5.20.3.4, 6.20.3.4, 7.20.3.4, 8.20.3.4, 9.20.3.4,  
 10.20.3.4, 11.20.3.4, 12.20.3.4, 13.20.3.4, 14.20.3.4, 15.20.3.4, 16.20.3.4, 17.20.3.4,  
 18.20.3.4, 19.20.3.4, 20.20.3.4, 21.20.3.4, 22.20.3.4, 23.20.3.4, 24.20.3.4, 25.20.3.4,  
 26.20.3.4, 27.20.3.4, 28.20.3.4, 29.20.3.4, 30.20.3.4, 31.20.3.4, 32.20.3.4, 33.20.3.4,  
 15 34.20.3.4, 35.20.3.4, 36.20.3.4, 37.20.3.4, 38.20.3.4, 39.20.3.4, 40.20.3.4, 1.21.3.4,  
 2.21.3.4, 3.21.3.4, 4.21.3.4, 5.21.3.4, 6.21.3.4, 7.21.3.4, 8.21.3.4, 9.21.3.4, 10.21.3.4,  
 11.21.3.4, 12.21.3.4, 13.21.3.4, 14.21.3.4, 15.21.3.4, 16.21.3.4, 17.21.3.4, 18.21.3.4,  
 19.21.3.4, 20.21.3.4, 21.21.3.4, 22.21.3.4, 23.21.3.4, 24.21.3.4, 25.21.3.4, 26.21.3.4,  
 27.21.3.4, 28.21.3.4, 29.21.3.4, 30.21.3.4, 31.21.3.4, 32.21.3.4, 33.21.3.4, 34.21.3.4,  
 20 35.21.3.4, 36.21.3.4, 37.21.3.4, 38.21.3.4, 39.21.3.4, 40.21.3.4, 1.22.3.4, 2.22.3.4,  
 3.22.3.4, 4.22.3.4, 5.22.3.4, 6.22.3.4, 7.22.3.4, 8.22.3.4, 9.22.3.4, 10.22.3.4, 11.22.3.4,  
 12.22.3.4, 13.22.3.4, 14.22.3.4, 15.22.3.4, 16.22.3.4, 17.22.3.4, 18.22.3.4, 19.22.3.4,  
 20.22.3.4, 21.22.3.4, 22.22.3.4, 23.22.3.4, 24.22.3.4, 25.22.3.4, 26.22.3.4, 27.22.3.4,  
 28.22.3.4, 29.22.3.4, 30.22.3.4, 31.22.3.4, 32.22.3.4, 33.22.3.4, 34.22.3.4, 35.22.3.4,  
 25 36.22.3.4, 37.22.3.4, 38.22.3.4, 39.22.3.4, 40.22.3.4, 1.23.3.4, 2.23.3.4, 3.23.3.4, 4.23.3.4,  
 5.23.3.4, 6.23.3.4, 7.23.3.4, 8.23.3.4, 9.23.3.4, 10.23.3.4, 11.23.3.4, 12.23.3.4, 13.23.3.4,  
 14.23.3.4, 15.23.3.4, 16.23.3.4, 17.23.3.4, 18.23.3.4, 19.23.3.4, 20.23.3.4, 21.23.3.4,  
 22.23.3.4, 23.23.3.4, 24.23.3.4, 25.23.3.4, 26.23.3.4, 27.23.3.4, 28.23.3.4, 29.23.3.4,  
 30.23.3.4, 31.23.3.4, 32.23.3.4, 33.23.3.4, 34.23.3.4, 35.23.3.4, 36.23.3.4, 37.23.3.4,  
 30 38.23.3.4, 39.23.3.4, 40.23.3.4, 1.24.3.4, 2.24.3.4, 3.24.3.4, 4.24.3.4, 5.24.3.4, 6.24.3.4,  
 7.24.3.4, 8.24.3.4, 9.24.3.4, 10.24.3.4, 11.24.3.4, 12.24.3.4, 13.24.3.4, 14.24.3.4,  
 15.24.3.4, 16.24.3.4, 17.24.3.4, 18.24.3.4, 19.24.3.4, 20.24.3.4, 21.24.3.4, 22.24.3.4,  
 23.24.3.4, 24.24.3.4, 25.24.3.4, 26.24.3.4, 27.24.3.4, 28.24.3.4, 29.24.3.4, 30.24.3.4,  
 31.24.3.4, 32.24.3.4, 33.24.3.4, 34.24.3.4, 35.24.3.4, 36.24.3.4, 37.24.3.4, 38.24.3.4,  
 35 39.24.3.4, 40.24.3.4, 1.25.3.4, 2.25.3.4, 3.25.3.4, 4.25.3.4, 5.25.3.4, 6.25.3.4, 7.25.3.4,



8.25.3.4, 9.25.3.4, 10.25.3.4, 11.25.3.4, 12.25.3.4, 13.25.3.4, 14.25.3.4, 15.25.3.4,  
 16.25.3.4, 17.25.3.4, 18.25.3.4, 19.25.3.4, 20.25.3.4, 21.25.3.4, 22.25.3.4, 23.25.3.4,  
 24.25.3.4, 25.25.3.4, 26.25.3.4, 27.25.3.4, 28.25.3.4, 29.25.3.4, 30.25.3.4, 31.25.3.4,  
 32.25.3.4, 33.25.3.4, 34.25.3.4, 35.25.3.4, 36.25.3.4, 37.25.3.4, 38.25.3.4, 39.25.3.4,  
 5 40.25.3.4, 1.26.3.4, 2.26.3.4, 3.26.3.4, 4.26.3.4, 5.26.3.4, 6.26.3.4, 7.26.3.4, 8.26.3.4,  
 9.26.3.4, 10.26.3.4, 11.26.3.4, 12.26.3.4, 13.26.3.4, 14.26.3.4, 15.26.3.4, 16.26.3.4,  
 17.26.3.4, 18.26.3.4, 19.26.3.4, 20.26.3.4, 21.26.3.4, 22.26.3.4, 23.26.3.4, 24.26.3.4,  
 25.26.3.4, 26.26.3.4, 27.26.3.4, 28.26.3.4, 29.26.3.4, 30.26.3.4, 31.26.3.4, 32.26.3.4,  
 33.26.3.4, 34.26.3.4, 35.26.3.4, 36.26.3.4, 37.26.3.4, 38.26.3.4, 39.26.3.4, 40.26.3.4,  
 10 1.27.3.4, 2.27.3.4, 3.27.3.4, 4.27.3.4, 5.27.3.4, 6.27.3.4, 7.27.3.4, 8.27.3.4, 9.27.3.4,  
 10.27.3.4, 11.27.3.4, 12.27.3.4, 13.27.3.4, 14.27.3.4, 15.27.3.4, 16.27.3.4, 17.27.3.4,  
 18.27.3.4, 19.27.3.4, 20.27.3.4, 21.27.3.4, 22.27.3.4, 23.27.3.4, 24.27.3.4, 25.27.3.4,  
 26.27.3.4, 27.27.3.4, 28.27.3.4, 29.27.3.4, 30.27.3.4, 31.27.3.4, 32.27.3.4, 33.27.3.4,  
 34.27.3.4, 35.27.3.4, 36.27.3.4, 37.27.3.4, 38.27.3.4, 39.27.3.4, 40.27.3.4, 1.28.3.4,  
 15 2.28.3.4, 3.28.3.4, 4.28.3.4, 5.28.3.4, 6.28.3.4, 7.28.3.4, 8.28.3.4, 9.28.3.4, 10.28.3.4,  
 11.28.3.4, 12.28.3.4, 13.28.3.4, 14.28.3.4, 15.28.3.4, 16.28.3.4, 17.28.3.4, 18.28.3.4,  
 19.28.3.4, 20.28.3.4, 21.28.3.4, 22.28.3.4, 23.28.3.4, 24.28.3.4, 25.28.3.4, 26.28.3.4,  
 27.28.3.4, 28.28.3.4, 29.28.3.4, 30.28.3.4, 31.28.3.4, 32.28.3.4, 33.28.3.4, 34.28.3.4,  
 35.28.3.4, 36.28.3.4, 37.28.3.4, 38.28.3.4, 39.28.3.4, 40.28.3.4, 1.29.3.4, 2.29.3.4,  
 20 3.29.3.4, 4.29.3.4, 5.29.3.4, 6.29.3.4, 7.29.3.4, 8.29.3.4, 9.29.3.4, 10.29.3.4, 11.29.3.4,  
 12.29.3.4, 13.29.3.4, 14.29.3.4, 15.29.3.4, 16.29.3.4, 17.29.3.4, 18.29.3.4, 19.29.3.4,  
 20.29.3.4, 21.29.3.4, 22.29.3.4, 23.29.3.4, 24.29.3.4, 25.29.3.4, 26.29.3.4, 27.29.3.4,  
 28.29.3.4, 29.29.3.4, 30.29.3.4, 31.29.3.4, 32.29.3.4, 33.29.3.4, 34.29.3.4, 35.29.3.4,  
 36.29.3.4, 37.29.3.4, 38.29.3.4, 39.29.3.4, 40.29.3.4, 1.1.3.5, 2.1.3.5, 3.1.3.5, 4.1.3.5,  
 25 5.1.3.5, 6.1.3.5, 7.1.3.5, 8.1.3.5, 9.1.3.5, 10.1.3.5, 11.1.3.5, 12.1.3.5, 13.1.3.5, 14.1.3.5,  
 15.1.3.5, 16.1.3.5, 17.1.3.5, 18.1.3.5, 19.1.3.5, 20.1.3.5, 21.1.3.5, 22.1.3.5, 23.1.3.5,  
 24.1.3.5, 25.1.3.5, 26.1.3.5, 27.1.3.5, 28.1.3.5, 29.1.3.5, 30.1.3.5, 31.1.3.5, 32.1.3.5,  
 33.1.3.5, 34.1.3.5, 35.1.3.5, 36.1.3.5, 37.1.3.5, 38.1.3.5, 39.1.3.5, 40.1.3.5, 1.2.3.5,  
 2.2.3.5, 3.2.3.5, 4.2.3.5, 5.2.3.5, 6.2.3.5, 7.2.3.5, 8.2.3.5, 9.2.3.5, 10.2.3.5, 11.2.3.5,  
 30 12.2.3.5, 13.2.3.5, 14.2.3.5, 15.2.3.5, 16.2.3.5, 17.2.3.5, 18.2.3.5, 19.2.3.5, 20.2.3.5,  
 21.2.3.5, 22.2.3.5, 23.2.3.5, 24.2.3.5, 25.2.3.5, 26.2.3.5, 27.2.3.5, 28.2.3.5, 29.2.3.5,  
 30.2.3.5, 31.2.3.5, 32.2.3.5, 33.2.3.5, 34.2.3.5, 35.2.3.5, 36.2.3.5, 37.2.3.5, 38.2.3.5,  
 39.2.3.5, 40.2.3.5, 1.3.3.5, 2.3.3.5, 3.3.3.5, 4.3.3.5, 5.3.3.5, 6.3.3.5, 7.3.3.5, 8.3.3.5,  
 9.3.3.5, 10.3.3.5, 11.3.3.5, 12.3.3.5, 13.3.3.5, 14.3.3.5, 15.3.3.5, 16.3.3.5, 17.3.3.5,  
 35 18.3.3.5, 19.3.3.5, 20.3.3.5, 21.3.3.5, 22.3.3.5, 23.3.3.5, 24.3.3.5, 25.3.3.5, 26.3.3.5,

27.3.3.5, 28.3.3.5, 29.3.3.5, 30.3.3.5, 31.3.3.5, 32.3.3.5, 33.3.3.5, 34.3.3.5, 35.3.3.5,  
 36.3.3.5, 37.3.3.5, 38.3.3.5, 39.3.3.5, 40.3.3.5, 1.4.3.5, 2.4.3.5, 3.4.3.5, 4.4.3.5, 5.4.3.5,  
 6.4.3.5, 7.4.3.5, 8.4.3.5, 9.4.3.5, 10.4.3.5, 11.4.3.5, 12.4.3.5, 13.4.3.5, 14.4.3.5, 15.4.3.5,  
 16.4.3.5, 17.4.3.5, 18.4.3.5, 19.4.3.5, 20.4.3.5, 21.4.3.5, 22.4.3.5, 23.4.3.5, 24.4.3.5,  
 5 25.4.3.5, 26.4.3.5, 27.4.3.5, 28.4.3.5, 29.4.3.5, 30.4.3.5, 31.4.3.5, 32.4.3.5, 33.4.3.5,  
 34.4.3.5, 35.4.3.5, 36.4.3.5, 37.4.3.5, 38.4.3.5, 39.4.3.5, 40.4.3.5, 1.5.3.5, 2.5.3.5, 3.5.3.5,  
 4.5.3.5, 5.5.3.5, 6.5.3.5, 7.5.3.5; 8.5.3.5, 9.5.3.5, 10.5.3.5, 11.5.3.5, 12.5.3.5, 13.5.3.5,  
 14.5.3.5, 15.5.3.5, 16.5.3.5, 17.5.3.5, 18.5.3.5, 19.5.3.5, 20.5.3.5, 21.5.3.5, 22.5.3.5,  
 23.5.3.5, 24.5.3.5, 25.5.3.5, 26.5.3.5, 27.5.3.5, 28.5.3.5, 29.5.3.5, 30.5.3.5, 31.5.3.5,  
 10 32.5.3.5, 33.5.3.5, 34.5.3.5, 35.5.3.5, 36.5.3.5, 37.5.3.5, 38.5.3.5, 39.5.3.5, 40.5.3.5,  
 1.6.3.5, 2.6.3.5, 3.6.3.5, 4.6.3.5, 5.6.3.5, 6.6.3.5, 7.6.3.5, 8.6.3.5, 9.6.3.5, 10.6.3.5,  
 11.6.3.5, 12.6.3.5, 13.6.3.5, 14.6.3.5, 15.6.3.5, 16.6.3.5, 17.6.3.5, 18.6.3.5, 19.6.3.5,  
 20.6.3.5, 21.6.3.5, 22.6.3.5, 23.6.3.5, 24.6.3.5, 25.6.3.5, 26.6.3.5, 27.6.3.5, 28.6.3.5,  
 29.6.3.5, 30.6.3.5, 31.6.3.5, 32.6.3.5, 33.6.3.5, 34.6.3.5, 35.6.3.5, 36.6.3.5, 37.6.3.5,  
 15 38.6.3.5, 39.6.3.5, 40.6.3.5, 1.7.3.5, 2.7.3.5, 3.7.3.5, 4.7.3.5, 5.7.3.5, 6.7.3.5, 7.7.3.5,  
 8.7.3.5, 9.7.3.5, 10.7.3.5, 11.7.3.5, 12.7.3.5, 13.7.3.5, 14.7.3.5, 15.7.3.5, 16.7.3.5,  
 17.7.3.5, 18.7.3.5, 19.7.3.5, 20.7.3.5, 21.7.3.5, 22.7.3.5, 23.7.3.5, 24.7.3.5, 25.7.3.5,  
 26.7.3.5, 27.7.3.5, 28.7.3.5, 29.7.3.5, 30.7.3.5, 31.7.3.5, 32.7.3.5, 33.7.3.5, 34.7.3.5,  
 35.7.3.5, 36.7.3.5, 37.7.3.5, 38.7.3.5, 39.7.3.5, 40.7.3.5, 1.8.3.5, 2.8.3.5, 3.8.3.5, 4.8.3.5,  
 20 5.8.3.5, 6.8.3.5, 7.8.3.5, 8.8.3.5, 9.8.3.5, 10.8.3.5, 11.8.3.5, 12.8.3.5, 13.8.3.5, 14.8.3.5,  
 15.8.3.5, 16.8.3.5, 17.8.3.5, 18.8.3.5, 19.8.3.5, 20.8.3.5, 21.8.3.5, 22.8.3.5, 23.8.3.5,  
 24.8.3.5, 25.8.3.5, 26.8.3.5, 27.8.3.5, 28.8.3.5, 29.8.3.5, 30.8.3.5, 31.8.3.5, 32.8.3.5,  
 33.8.3.5, 34.8.3.5, 35.8.3.5, 36.8.3.5, 37.8.3.5, 38.8.3.5, 39.8.3.5, 40.8.3.5, 1.9.3.5,  
 2.9.3.5, 3.9.3.5, 4.9.3.5, 5.9.3.5, 6.9.3.5, 7.9.3.5, 8.9.3.5, 9.9.3.5, 10.9.3.5, 11.9.3.5,  
 25 12.9.3.5, 13.9.3.5, 14.9.3.5, 15.9.3.5, 16.9.3.5, 17.9.3.5, 18.9.3.5, 19.9.3.5, 20.9.3.5,  
 21.9.3.5, 22.9.3.5, 23.9.3.5, 24.9.3.5, 25.9.3.5, 26.9.3.5, 27.9.3.5, 28.9.3.5, 29.9.3.5,  
 30.9.3.5, 31.9.3.5, 32.9.3.5, 33.9.3.5, 34.9.3.5, 35.9.3.5, 36.9.3.5, 37.9.3.5, 38.9.3.5,  
 39.9.3.5, 40.9.3.5, 1.10.3.5, 2.10.3.5, 3.10.3.5, 4.10.3.5, 5.10.3.5, 6.10.3.5, 7.10.3.5,  
 8.10.3.5, 9.10.3.5, 10.10.3.5, 11.10.3.5, 12.10.3.5, 13.10.3.5, 14.10.3.5, 15.10.3.5,  
 30 16.10.3.5, 17.10.3.5, 18.10.3.5, 19.10.3.5, 20.10.3.5, 21.10.3.5, 22.10.3.5, 23.10.3.5,  
 24.10.3.5, 25.10.3.5, 26.10.3.5, 27.10.3.5, 28.10.3.5, 29.10.3.5, 30.10.3.5, 31.10.3.5,  
 32.10.3.5, 33.10.3.5, 34.10.3.5, 35.10.3.5, 36.10.3.5, 37.10.3.5, 38.10.3.5, 39.10.3.5,  
 40.10.3.5, 1.11.3.5, 2.11.3.5, 3.11.3.5, 4.11.3.5, 5.11.3.5, 6.11.3.5, 7.11.3.5, 8.11.3.5,  
 9.11.3.5, 10.11.3.5, 11.11.3.5, 12.11.3.5, 13.11.3.5, 14.11.3.5, 15.11.3.5, 16.11.3.5,  
 35 17.11.3.5, 18.11.3.5, 19.11.3.5, 20.11.3.5, 21.11.3.5, 22.11.3.5, 23.11.3.5, 24.11.3.5,

25.11.3.5, 26.11.3.5, 27.11.3.5, 28.11.3.5, 29.11.3.5, 30.11.3.5, 31.11.3.5, 32.11.3.5,  
33.11.3.5, 34.11.3.5, 35.11.3.5, 36.11.3.5, 37.11.3.5, 38.11.3.5, 39.11.3.5, 40.11.3.5,  
1.12.3.5, 2.12.3.5, 3.12.3.5, 4.12.3.5, 5.12.3.5, 6.12.3.5, 7.12.3.5, 8.12.3.5, 9.12.3.5,  
10.12.3.5, 11.12.3.5, 12.12.3.5, 13.12.3.5, 14.12.3.5, 15.12.3.5, 16.12.3.5, 17.12.3.5,  
5 18.12.3.5, 19.12.3.5, 20.12.3.5, 21.12.3.5, 22.12.3.5, 23.12.3.5, 24.12.3.5, 25.12.3.5,  
26.12.3.5, 27.12.3.5, 28.12.3.5, 29.12.3.5, 30.12.3.5, 31.12.3.5, 32.12.3.5, 33.12.3.5,  
34.12.3.5, 35.12.3.5, 36.12.3.5, 37.12.3.5, 38.12.3.5, 39.12.3.5, 40.12.3.5, 1.13.3.5,  
2.13.3.5, 3.13.3.5, 4.13.3.5, 5.13.3.5, 6.13.3.5, 7.13.3.5, 8.13.3.5, 9.13.3.5, 10.13.3.5,  
11.13.3.5, 12.13.3.5, 13.13.3.5, 14.13.3.5, 15.13.3.5, 16.13.3.5, 17.13.3.5, 18.13.3.5,  
10 19.13.3.5, 20.13.3.5, 21.13.3.5, 22.13.3.5, 23.13.3.5, 24.13.3.5, 25.13.3.5, 26.13.3.5,  
27.13.3.5, 28.13.3.5, 29.13.3.5, 30.13.3.5, 31.13.3.5, 32.13.3.5, 33.13.3.5, 34.13.3.5,  
35.13.3.5, 36.13.3.5, 37.13.3.5, 38.13.3.5, 39.13.3.5, 40.13.3.5, 1.14.3.5, 2.14.3.5,  
3.14.3.5, 4.14.3.5, 5.14.3.5, 6.14.3.5, 7.14.3.5, 8.14.3.5, 9.14.3.5, 10.14.3.5, 11.14.3.5,  
12.14.3.5, 13.14.3.5, 14.14.3.5, 15.14.3.5, 16.14.3.5, 17.14.3.5, 18.14.3.5, 19.14.3.5,  
15 20.14.3.5, 21.14.3.5, 22.14.3.5, 23.14.3.5, 24.14.3.5, 25.14.3.5, 26.14.3.5, 27.14.3.5,  
28.14.3.5, 29.14.3.5, 30.14.3.5, 31.14.3.5, 32.14.3.5, 33.14.3.5, 34.14.3.5, 35.14.3.5,  
36.14.3.5, 37.14.3.5, 38.14.3.5, 39.14.3.5, 40.14.3.5, 1.15.3.5, 2.15.3.5, 3.15.3.5, 4.15.3.5,  
5.15.3.5, 6.15.3.5, 7.15.3.5, 8.15.3.5, 9.15.3.5, 10.15.3.5, 11.15.3.5, 12.15.3.5, 13.15.3.5,  
14.15.3.5, 15.15.3.5, 16.15.3.5, 17.15.3.5, 18.15.3.5, 19.15.3.5, 20.15.3.5, 21.15.3.5,  
20 22.15.3.5, 23.15.3.5, 24.15.3.5, 25.15.3.5, 26.15.3.5, 27.15.3.5, 28.15.3.5, 29.15.3.5,  
30.15.3.5, 31.15.3.5, 32.15.3.5, 33.15.3.5, 34.15.3.5, 35.15.3.5, 36.15.3.5, 37.15.3.5,  
38.15.3.5, 39.15.3.5, 40.15.3.5, 1.16.3.5, 2.16.3.5, 3.16.3.5, 4.16.3.5, 5.16.3.5, 6.16.3.5,  
7.16.3.5, 8.16.3.5, 9.16.3.5, 10.16.3.5, 11.16.3.5, 12.16.3.5, 13.16.3.5, 14.16.3.5,  
15.16.3.5, 16.16.3.5, 17.16.3.5, 18.16.3.5, 19.16.3.5, 20.16.3.5, 21.16.3.5, 22.16.3.5,  
25 23.16.3.5, 24.16.3.5, 25.16.3.5, 26.16.3.5, 27.16.3.5, 28.16.3.5, 29.16.3.5, 30.16.3.5,  
31.16.3.5, 32.16.3.5, 33.16.3.5, 34.16.3.5, 35.16.3.5, 36.16.3.5, 37.16.3.5, 38.16.3.5,  
39.16.3.5, 40.16.3.5, 1.17.3.5, 2.17.3.5, 3.17.3.5, 4.17.3.5, 5.17.3.5, 6.17.3.5, 7.17.3.5,  
8.17.3.5, 9.17.3.5, 10.17.3.5, 11.17.3.5, 12.17.3.5, 13.17.3.5, 14.17.3.5, 15.17.3.5,  
16.17.3.5, 17.17.3.5, 18.17.3.5, 19.17.3.5, 20.17.3.5, 21.17.3.5, 22.17.3.5, 23.17.3.5,  
30 24.17.3.5, 25.17.3.5, 26.17.3.5, 27.17.3.5, 28.17.3.5, 29.17.3.5, 30.17.3.5, 31.17.3.5,  
32.17.3.5, 33.17.3.5, 34.17.3.5, 35.17.3.5, 36.17.3.5, 37.17.3.5, 38.17.3.5, 39.17.3.5,  
40.17.3.5, 1.18.3.5, 2.18.3.5, 3.18.3.5, 4.18.3.5, 5.18.3.5, 6.18.3.5, 7.18.3.5, 8.18.3.5,  
9.18.3.5, 10.18.3.5, 11.18.3.5, 12.18.3.5, 13.18.3.5, 14.18.3.5, 15.18.3.5, 16.18.3.5,  
17.18.3.5, 18.18.3.5, 19.18.3.5, 20.18.3.5, 21.18.3.5, 22.18.3.5, 23.18.3.5, 24.18.3.5,  
35 25.18.3.5, 26.18.3.5, 27.18.3.5, 28.18.3.5, 29.18.3.5, 30.18.3.5, 31.18.3.5, 32.18.3.5,

33.18.3.5, 34.18.3.5, 35.18.3.5, 36.18.3.5, 37.18.3.5, 38.18.3.5, 39.18.3.5, 40.18.3.5,  
1.19.3.5, 2.19.3.5, 3.19.3.5, 4.19.3.5, 5.19.3.5, 6.19.3.5, 7.19.3.5, 8.19.3.5, 9.19.3.5,  
10.19.3.5, 11.19.3.5, 12.19.3.5, 13.19.3.5, 14.19.3.5, 15.19.3.5, 16.19.3.5, 17.19.3.5,  
18.19.3.5, 19.19.3.5, 20.19.3.5, 21.19.3.5, 22.19.3.5, 23.19.3.5, 24.19.3.5, 25.19.3.5,  
5 26.19.3.5, 27.19.3.5, 28.19.3.5, 29.19.3.5, 30.19.3.5, 31.19.3.5, 32.19.3.5, 33.19.3.5,  
34.19.3.5, 35.19.3.5, 36.19.3.5, 37.19.3.5, 38.19.3.5, 39.19.3.5, 40.19.3.5, 1.20.3.5,  
2.20.3.5, 3.20.3.5, 4.20.3.5, 5.20.3.5, 6.20.3.5, 7.20.3.5, 8.20.3.5, 9.20.3.5, 10.20.3.5,  
11.20.3.5, 12.20.3.5, 13.20.3.5, 14.20.3.5, 15.20.3.5, 16.20.3.5, 17.20.3.5, 18.20.3.5,  
19.20.3.5, 20.20.3.5, 21.20.3.5, 22.20.3.5, 23.20.3.5, 24.20.3.5, 25.20.3.5, 26.20.3.5,  
10 27.20.3.5, 28.20.3.5, 29.20.3.5, 30.20.3.5, 31.20.3.5, 32.20.3.5, 33.20.3.5, 34.20.3.5,  
35.20.3.5, 36.20.3.5, 37.20.3.5, 38.20.3.5, 39.20.3.5, 40.20.3.5, 1.21.3.5, 2.21.3.5,  
3.21.3.5, 4.21.3.5, 5.21.3.5, 6.21.3.5, 7.21.3.5, 8.21.3.5, 9.21.3.5, 10.21.3.5, 11.21.3.5,  
12.21.3.5, 13.21.3.5, 14.21.3.5, 15.21.3.5, 16.21.3.5, 17.21.3.5, 18.21.3.5, 19.21.3.5,  
20.21.3.5, 21.21.3.5, 22.21.3.5, 23.21.3.5, 24.21.3.5, 25.21.3.5, 26.21.3.5, 27.21.3.5,  
15 28.21.3.5, 29.21.3.5, 30.21.3.5, 31.21.3.5, 32.21.3.5, 33.21.3.5, 34.21.3.5, 35.21.3.5,  
36.21.3.5, 37.21.3.5, 38.21.3.5, 39.21.3.5, 40.21.3.5, 1.22.3.5, 2.22.3.5, 3.22.3.5, 4.22.3.5,  
5.22.3.5, 6.22.3.5, 7.22.3.5, 8.22.3.5, 9.22.3.5, 10.22.3.5, 11.22.3.5, 12.22.3.5, 13.22.3.5,  
14.22.3.5, 15.22.3.5, 16.22.3.5, 17.22.3.5, 18.22.3.5, 19.22.3.5, 20.22.3.5, 21.22.3.5,  
22.22.3.5, 23.22.3.5, 24.22.3.5, 25.22.3.5, 26.22.3.5, 27.22.3.5, 28.22.3.5, 29.22.3.5,  
20 30.22.3.5, 31.22.3.5, 32.22.3.5, 33.22.3.5, 34.22.3.5, 35.22.3.5, 36.22.3.5, 37.22.3.5,  
38.22.3.5, 39.22.3.5, 40.22.3.5, 1.23.3.5, 2.23.3.5, 3.23.3.5, 4.23.3.5, 5.23.3.5, 6.23.3.5,  
7.23.3.5, 8.23.3.5, 9.23.3.5, 10.23.3.5, 11.23.3.5, 12.23.3.5, 13.23.3.5, 14.23.3.5,  
15.23.3.5, 16.23.3.5, 17.23.3.5, 18.23.3.5, 19.23.3.5, 20.23.3.5, 21.23.3.5, 22.23.3.5,  
23.23.3.5, 24.23.3.5, 25.23.3.5, 26.23.3.5, 27.23.3.5, 28.23.3.5, 29.23.3.5, 30.23.3.5,  
25 31.23.3.5, 32.23.3.5, 33.23.3.5, 34.23.3.5, 35.23.3.5, 36.23.3.5, 37.23.3.5, 38.23.3.5,  
39.23.3.5, 40.23.3.5, 1.24.3.5, 2.24.3.5, 3.24.3.5, 4.24.3.5, 5.24.3.5, 6.24.3.5, 7.24.3.5,  
8.24.3.5, 9.24.3.5, 10.24.3.5, 11.24.3.5, 12.24.3.5, 13.24.3.5, 14.24.3.5, 15.24.3.5,  
16.24.3.5, 17.24.3.5, 18.24.3.5, 19.24.3.5, 20.24.3.5, 21.24.3.5, 22.24.3.5, 23.24.3.5,  
24.24.3.5, 25.24.3.5, 26.24.3.5, 27.24.3.5, 28.24.3.5, 29.24.3.5, 30.24.3.5, 31.24.3.5,  
30 32.24.3.5, 33.24.3.5, 34.24.3.5, 35.24.3.5, 36.24.3.5, 37.24.3.5, 38.24.3.5, 39.24.3.5,  
40.24.3.5, 1.25.3.5, 2.25.3.5, 3.25.3.5, 4.25.3.5, 5.25.3.5, 6.25.3.5, 7.25.3.5, 8.25.3.5,  
9.25.3.5, 10.25.3.5, 11.25.3.5, 12.25.3.5, 13.25.3.5, 14.25.3.5, 15.25.3.5, 16.25.3.5,  
17.25.3.5, 18.25.3.5, 19.25.3.5, 20.25.3.5, 21.25.3.5, 22.25.3.5, 23.25.3.5, 24.25.3.5,  
25.25.3.5, 26.25.3.5, 27.25.3.5, 28.25.3.5, 29.25.3.5, 30.25.3.5, 31.25.3.5, 32.25.3.5,  
35 33.25.3.5, 34.25.3.5, 35.25.3.5, 36.25.3.5, 37.25.3.5, 38.25.3.5, 39.25.3.5, 40.25.3.5,

1.26.3.5, 2.26.3.5, 3.26.3.5, 4.26.3.5, 5.26.3.5, 6.26.3.5, 7.26.3.5, 8.26.3.5, 9.26.3.5,  
 10.26.3.5, 11.26.3.5, 12.26.3.5, 13.26.3.5, 14.26.3.5, 15.26.3.5, 16.26.3.5, 17.26.3.5,  
 18.26.3.5, 19.26.3.5, 20.26.3.5, 21.26.3.5, 22.26.3.5, 23.26.3.5, 24.26.3.5, 25.26.3.5,  
 26.26.3.5, 27.26.3.5, 28.26.3.5, 29.26.3.5, 30.26.3.5, 31.26.3.5, 32.26.3.5, 33.26.3.5,  
 5 34.26.3.5, 35.26.3.5, 36.26.3.5, 37.26.3.5, 38.26.3.5, 39.26.3.5, 40.26.3.5, 1.27.3.5,  
 2.27.3.5, 3.27.3.5, 4.27.3.5, 5.27.3.5, 6.27.3.5, 7.27.3.5, 8.27.3.5, 9.27.3.5, 10.27.3.5,  
 11.27.3.5, 12.27.3.5, 13.27.3.5, 14.27.3.5, 15.27.3.5, 16.27.3.5, 17.27.3.5, 18.27.3.5,  
 19.27.3.5, 20.27.3.5, 21.27.3.5, 22.27.3.5, 23.27.3.5, 24.27.3.5, 25.27.3.5, 26.27.3.5,  
 27.27.3.5, 28.27.3.5, 29.27.3.5, 30.27.3.5, 31.27.3.5, 32.27.3.5, 33.27.3.5, 34.27.3.5,  
 10 35.27.3.5, 36.27.3.5, 37.27.3.5, 38.27.3.5, 39.27.3.5, 40.27.3.5, 1.28.3.5, 2.28.3.5,  
 3.28.3.5, 4.28.3.5, 5.28.3.5, 6.28.3.5, 7.28.3.5, 8.28.3.5, 9.28.3.5, 10.28.3.5, 11.28.3.5,  
 12.28.3.5, 13.28.3.5, 14.28.3.5, 15.28.3.5, 16.28.3.5, 17.28.3.5, 18.28.3.5, 19.28.3.5,  
 20.28.3.5, 21.28.3.5, 22.28.3.5, 23.28.3.5, 24.28.3.5, 25.28.3.5, 26.28.3.5, 27.28.3.5,  
 28.28.3.5, 29.28.3.5, 30.28.3.5, 31.28.3.5, 32.28.3.5, 33.28.3.5, 34.28.3.5, 35.28.3.5,  
 15 36.28.3.5, 37.28.3.5, 38.28.3.5, 39.28.3.5, 40.28.3.5, 1.29.3.5, 2.29.3.5, 3.29.3.5, 4.29.3.5,  
 5.29.3.5, 6.29.3.5, 7.29.3.5, 8.29.3.5, 9.29.3.5, 10.29.3.5, 11.29.3.5, 12.29.3.5, 13.29.3.5,  
 14.29.3.5, 15.29.3.5, 16.29.3.5, 17.29.3.5, 18.29.3.5, 19.29.3.5, 20.29.3.5, 21.29.3.5,  
 22.29.3.5, 23.29.3.5, 24.29.3.5, 25.29.3.5, 26.29.3.5, 27.29.3.5, 28.29.3.5, 29.29.3.5,  
 30.29.3.5, 31.29.3.5, 32.29.3.5, 33.29.3.5, 34.29.3.5, 35.29.3.5, 36.29.3.5, 37.29.3.5,  
 20 38.29.3.5, 39.29.3.5 and 40.29.3.5.

Table 2 lists a group of cyclic nucleotide analogs of structure I wherein  
 Z forms a heterocyclic ring containing the phosphorus atom of the  
 25 phosphonate group and two oxygen atoms as shown. Hydrolysis of the L<sup>1</sup>  
 group linked to the phosphorus atom and subsequent ring hydrolysis results  
 in formation of an HPMP nucleoside such as HPMPC (1-(2-  
 phosphonomethoxy-3-hydroxypropyl)-cytosine).

30



TABLE 2

---

	L <sup>1*</sup>
1	-NH-CH <sub>2</sub> -C(O)-OR <sup>4</sup>
2	-NH-CH(CH <sub>3</sub> )-C(O)-OR <sup>4</sup>
35 3	-NH-CH(CH <sub>3</sub> ) <sub>2</sub> -C(O)-OR <sup>4</sup>

- 4 -NH-CH(CH(CH<sub>3</sub>)<sub>2</sub>)-C(O)-OR<sup>4</sup>  
 5 -NH-CH(CH<sub>3</sub>)(CH<sub>3</sub>)<sub>2</sub>-C(O)-OR<sup>4</sup>  
 6 -N-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH-C(O)-OR<sup>4</sup>  
 7 -NH-CH(CH<sub>2</sub>-C<sub>6</sub>H<sub>5</sub>)-C(O)-OR<sup>4</sup>  
 5 8 -NH-CH(CH<sub>2</sub>-C<sub>8</sub>NH<sub>6</sub>)-C(O)-OR<sup>4</sup>  
 9 -NH-CH(CH<sub>2</sub>-CH<sub>2</sub>-S-CH<sub>3</sub>)-C(O)-OR<sup>4</sup>  
 10 -NH-CH(CH<sub>2</sub>OH)-C(O)-OR<sup>4</sup>  
 11 -NH-CH(CH(OH)(CH<sub>3</sub>)-C(O)-OR<sup>4</sup>  
 12 -NH-CH(-CH<sub>2</sub>SH)-C(O)-OR<sup>4</sup>  
 10 13 -NH-CH(CH<sub>2</sub>-C<sub>6</sub>H<sub>5</sub>OH)-C(O)-OR<sup>4</sup>  
 14 -NH-CH(CH<sub>2</sub>-C(O)-NH<sub>2</sub>)-C(O)-OR<sup>4</sup>  
 15 -NH-CH(CH<sub>2</sub>-CH<sub>2</sub>-C(O)-NH<sub>2</sub>)-C(O)-OR<sup>4</sup>  
 16 -NH-CH(CH<sub>2</sub>C(O)OR<sup>4</sup>)-C(O)-OR<sup>4</sup>  
 17 -NH-CH(CH<sub>2</sub>CH<sub>2</sub>C(O)OR<sup>4</sup>)-C(O)-OR<sup>4</sup>  
 15 18 -NH-CH(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>)-C(O)-OR<sup>4</sup>  
 19 -NH-CH(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NHC(NH)(NH<sub>2</sub>))-C(O)-OR<sup>4</sup>  
 20 -NH-CH(CH<sub>2</sub>C<sub>3</sub>N<sub>2</sub>H<sub>3</sub>)-C(O)-OR<sup>4</sup>  
 21 -NH-CH(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>)-CH<sub>2</sub>-C(O)-OR<sup>4</sup>  
 22 -NH-CH(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>)-CH<sub>2</sub>-C(O)-OR<sup>4</sup>  
 20 23 -NH-CH(CH<sub>2</sub>CH<sub>2</sub>NHC(NH)(NH<sub>2</sub>))-CH<sub>2</sub>-C(O)-OR<sup>4</sup>  
 24 -NH-CH(C(O)OR<sup>4</sup>)-CH<sub>2</sub>-C(O)-OR<sup>4</sup>  
 25 -NH-CH(CH<sub>2</sub>C(O)OR<sup>4</sup>)-CH<sub>2</sub>-C(O)-OR<sup>4</sup>  
 26 -NH-CH(CH<sub>2</sub>CH<sub>2</sub>C(O)OR<sup>4</sup>)-CH<sub>2</sub>-C(O)-OR<sup>4</sup>

25 Z-B\*\*



B

1. adenin-9-yl  
 2. guanin-9-yl  
 30 3. cytosin-1-yl  
 4. 2, 6-diaminopurin-9-yl

- 5. 2-aminopurin-9-yl
  - 6. 6-azacytosin-1-yl
  - 7. 1-deazaadenin-9-yl
  - 8. 3-deazaadenin-9-yl
  - 9. 8-azaadenin-9-yl
  - 10. 7-deaza-8-azaadenin-9-yl
- 

5 \* - See Table 1 footnote.

\*\* - See Table 1 footnote.

# - See Table 1 footnote.

Compounds listed in Table 2 are designated herein by numbers assigned to L<sup>1</sup>, Z and B according to the following convention, L.Z.B. Thus, compounds 1.1.3 and 1.2.3 represent, when R<sub>4</sub> is H, glycyl cyclic HPMPC and alanyl cyclic HPMPC. Exemplary compounds include 1.1.1, 1.1.2, 1.1.3, 1.1.4, 1.1.5, 1.1.6, 1.1.7, 1.1.8, 1.1.9, 1.1.10, 2.1.1, 2.1.2, 2.1.3, 2.1.4, 2.1.5, 2.1.6, 2.1.7, 2.1.8, 2.1.9, 2.1.10, 3.1.1, 3.1.2, 3.1.3, 3.1.4, 3.1.5, 3.1.6, 3.1.7, 3.1.8, 3.1.9, 3.1.10, 4.1.1, 4.1.2, 4.1.3, 4.1.4, 4.1.5, 4.1.6, 4.1.7, 4.1.8, 4.1.9, 4.1.10, 5.1.1, 5.1.2, 5.1.3, 5.1.4, 5.1.5, 5.1.6, 5.1.7, 5.1.8, 5.1.9, 5.1.10, 6.1.1, 6.1.2, 6.1.3, 6.1.4, 6.1.5, 6.1.6, 6.1.7, 6.1.8, 6.1.9, 6.1.10, 7.1.1, 7.1.2, 7.1.3, 7.1.4, 7.1.5, 7.1.6, 7.1.7, 7.1.8, 7.1.9, 7.1.10, 8.1.1, 8.1.2, 8.1.3, 8.1.4, 8.1.5, 8.1.6, 8.1.7, 8.1.8, 8.1.9, 8.1.10, 9.1.1, 9.1.2, 9.1.3, 9.1.4, 9.1.5, 9.1.6, 9.1.7, 9.1.8, 9.1.9, 9.1.10, 10.1.1, 10.1.2, 10.1.3, 10.1.4, 10.1.5, 10.1.6, 10.1.7, 10.1.8, 10.1.9, 10.1.10, 11.1.1, 11.1.2, 11.1.3, 11.1.4, 11.1.5, 11.1.6, 11.1.7, 11.1.8, 11.1.9, 11.1.10, 12.1.1, 12.1.2, 12.1.3, 12.1.4, 12.1.5, 12.1.6, 12.1.7, 12.1.8, 12.1.9, 12.1.10, 13.1.1, 13.1.2, 13.1.3, 13.1.4, 13.1.5, 13.1.6, 13.1.7, 13.1.8, 13.1.9, 13.1.10, 14.1.1, 14.1.2, 14.1.3, 14.1.4, 14.1.5, 14.1.6, 14.1.7, 14.1.8, 14.1.9, 14.1.10, 15.1.1, 15.1.2, 15.1.3, 15.1.4, 15.1.5, 15.1.6, 15.1.7, 15.1.8, 15.1.9, 15.1.10, 16.1.1, 16.1.2, 16.1.3, 16.1.4, 16.1.5, 16.1.6, 16.1.7, 16.1.8, 16.1.9, 16.1.10, 17.1.1, 17.1.2, 17.1.3, 17.1.4, 17.1.5, 17.1.6, 17.1.7, 17.1.8, 17.1.9, 17.1.10, 18.1.1, 18.1.2, 18.1.3, 18.1.4, 18.1.5, 18.1.6, 18.1.7, 18.1.8, 18.1.9, 18.1.10, 19.1.1, 19.1.2, 19.1.3, 19.1.4, 19.1.5, 19.1.6, 19.1.7, 19.1.8, 19.1.9, 19.1.10, 20.1.1, 20.1.2, 20.1.3, 20.1.4, 20.1.5, 20.1.6, 20.1.7, 20.1.8, 20.1.9, 20.1.10, 21.1.1, 21.1.2, 21.1.3, 21.1.4, 21.1.5, 21.1.6, 21.1.7, 21.1.8, 21.1.9, 21.1.10, 22.1.1, 22.1.2, 22.1.3, 22.1.4, 22.1.5, 22.1.6, 22.1.7, 22.1.8, 22.1.9, 22.1.10, 23.1.1, 23.1.2, 23.1.3, 23.1.4, 23.1.5, 23.1.6, 23.1.7, 23.1.8, 23.1.9, 23.1.10, 24.1.1, 24.1.2, 24.1.3, 24.1.4, 24.1.5, 24.1.6, 24.1.7, 24.1.8, 24.1.9, 24.1.10, 25.1.1, 25.1.2, 25.1.3, 25.1.4, 25.1.5, 25.1.6, 25.1.7, 25.1.8, 25.1.9, 25.1.10, 26.1.1, 26.1.2, 26.1.3, 26.1.4, 26.1.5, 26.1.6, 26.1.7, 26.1.8, 26.1.9, 26.1.10, 27.1.1, 27.1.2, 27.1.3, 27.1.4, 27.1.5, 27.1.6, 27.1.7, 27.1.8, 27.1.9, 27.1.10, 28.1.1, 28.1.2, 28.1.3, 28.1.4, 28.1.5, 28.1.6, 28.1.7,

28.1.8, 28.1.9, 28.1.10, 1.2.1, 1.2.2, 1.2.3, 1.2.4, 1.2.5, 1.2.6, 1.2.7, 1.2.8, 1.2.9, 1.2.10, 2.2.1, 2.2.2, 2.2.3, 2.2.4, 2.2.5, 2.2.6, 2.2.7, 2.2.8, 2.2.9, 2.2.10, 3.2.1, 3.2.2, 3.2.3, 3.2.4, 3.2.5, 3.2.6, 3.2.7, 3.2.8, 3.2.9, 3.2.10, 4.2.1, 4.2.2, 4.2.3, 4.2.4, 4.2.5, 4.2.6, 4.2.7, 4.2.8, 4.2.9, 4.2.10, 5.2.1, 5.2.2, 5.2.3, 5.2.4, 5.2.5, 5.2.6, 5.2.7, 5.2.8, 5.2.9, 5.2.10, 6.2.1, 6.2.2, 6.2.3, 6.2.4, 6.2.5, 6.2.6, 6.2.7, 6.2.8, 6.2.9, 6.2.10, 7.2.1, 7.2.2, 7.2.3, 7.2.4, 7.2.5, 7.2.6, 7.2.7, 7.2.8, 7.2.9, 7.2.10, 8.2.1, 8.2.2, 8.2.3, 8.2.4, 8.2.5, 8.2.6, 8.2.7, 8.2.8, 8.2.9, 8.2.10, 9.2.1, 9.2.2, 9.2.3, 9.2.4, 9.2.5, 9.2.6, 9.2.7, 9.2.8, 9.2.9, 9.2.10, 10.2.1, 10.2.2, 10.2.3, 10.2.4, 10.2.5, 10.2.6, 10.2.7, 10.2.8, 10.2.9, 10.2.10, 11.2.1, 11.2.2, 11.2.3, 11.2.4, 11.2.5, 11.2.6, 11.2.7, 11.2.8, 11.2.9, 11.2.10, 12.2.1, 12.2.2, 12.2.3, 12.2.4, 12.2.5, 12.2.6, 12.2.7, 12.2.8, 12.2.9, 12.2.10, 13.2.1, 13.2.2, 13.2.3, 13.2.4, 13.2.5, 13.2.6, 13.2.7, 13.2.8, 13.2.9, 13.2.10, 14.2.1, 14.2.2, 14.2.3, 14.2.4, 14.2.5, 14.2.6, 14.2.7, 14.2.8, 14.2.9, 14.2.10, 15.2.1, 15.2.2, 15.2.3, 15.2.4, 15.2.5, 15.2.6, 15.2.7, 15.2.8, 15.2.9, 15.2.10, 16.2.1, 16.2.2, 16.2.3, 16.2.4, 16.2.5, 16.2.6, 16.2.7, 16.2.8, 16.2.9, 16.2.10, 17.2.1, 17.2.2, 17.2.3, 17.2.4, 17.2.5, 17.2.6, 17.2.7, 17.2.8, 17.2.9, 17.2.10, 18.2.1, 18.2.2, 18.2.3, 18.2.4, 18.2.5, 18.2.6, 18.2.7, 18.2.8, 18.2.9, 18.2.10, 19.2.1, 19.2.2, 19.2.3, 19.2.4, 19.2.5, 19.2.6, 19.2.7, 19.2.8, 19.2.9, 19.2.10, 20.2.1, 20.2.2, 20.2.3, 20.2.4, 20.2.5, 20.2.6, 20.2.7, 20.2.8, 20.2.9, 20.2.10, 21.2.1, 21.2.2, 21.2.3, 21.2.4, 21.2.5, 21.2.6, 21.2.7, 21.2.8, 21.2.9, 21.2.10, 22.2.1, 22.2.2, 22.2.3, 22.2.4, 22.2.5, 22.2.6, 22.2.7, 22.2.8, 22.2.9, 22.2.10, 23.2.1, 23.2.2, 23.2.3, 23.2.4, 23.2.5, 23.2.6, 23.2.7, 23.2.8, 23.2.9, 23.2.10, 24.2.1, 24.2.2, 24.2.3, 24.2.4, 24.2.5, 24.2.6, 24.2.7, 24.2.8, 24.2.9, 24.2.10, 25.2.1, 25.2.2, 25.2.3, 25.2.4, 25.2.5, 25.2.6, 25.2.7, 25.2.8, 25.2.9, 25.2.10, 26.2.1, 26.2.2, 26.2.3, 26.2.4, 26.2.5, 26.2.6, 26.2.7, 26.2.8, 26.2.9 and 26.2.10.

Table 3 lists a group of cyclic nucleotide analog amidates of structure I wherein L<sup>1</sup> forms a heterocyclic ring containing the phosphorus atom of the phosphonate group. Hydrolysis of the heterocyclic ring linked through the phosphorus atom results in formation of a phosphonate nucleotide analog such as HPMPC, PMEA, PMEG or PMPDAP depending on the Z group that is present.

TABLE 3

L <sup>1</sup>	Z-B <sup>**</sup>
1 -NH-CH <sub>2</sub> -C(O)-O-CH <sub>2</sub> -O-	1 -CH <sub>2</sub> -O-CH <sub>2</sub> -CH <sub>2</sub> -B
2 -NH-CH(CH <sub>3</sub> )-C(O)-O-CH <sub>2</sub> -O-	2 -CH <sub>2</sub> -O-C <sup>#</sup> H(CH <sub>2</sub> -OR <sup>4</sup> )-CH <sub>2</sub> -B



- |    |   |   |
|----|---|---|
|    | 3 -NH-CH(CH <sub>3</sub> ) <sub>2</sub> -C(O)-O-CH <sub>2</sub> -O-   | 3 -CH <sub>2</sub> -O-C#H(CH <sub>3</sub> )-CH <sub>2</sub> -B                |
|    | 4 -NH-CH(CH(CH <sub>3</sub> ) <sub>2</sub> )-C(O)-O-CH <sub>2</sub> -O-   | 4 -CH <sub>2</sub> -O-C#H(CH <sub>2</sub> F)-CH <sub>2</sub> -B               |
|    | 5 -NH-CH(CH <sub>3</sub> )(CH <sub>3</sub> ) <sub>2</sub> -C(O)-O-CH <sub>2</sub> -O-   | 5 -CH <sub>2</sub> -O-C#H(CH=CH <sub>2</sub> )-CH <sub>2</sub> -B             |
|    | 6 -NH-CH <sub>2</sub> -CH <sub>2</sub> -CH <sub>2</sub> -CH-C(O)-O-CH <sub>2</sub> -O-  | 6 -CH <sub>2</sub> -O-C#H(CH <sub>2</sub> N <sub>3</sub> )-CH <sub>2</sub> -B |
| 5  | 7 -NH-CH(CH <sub>2</sub> -C <sub>6</sub> H <sub>5</sub> )-C(O)-O-CH <sub>2</sub> -O-  |   |
|    | 8 -NH-CH(CH <sub>2</sub> -C <sub>8</sub> NH <sub>6</sub> )-C(O)-O-CH <sub>2</sub> -O-   |   |
|    | 9 -NH-CH(CH <sub>2</sub> -CH <sub>2</sub> -S-CH <sub>3</sub> )-C(O)-O-  |   |
|    | 10 -NH-CH(CH <sub>2</sub> OH)-C(O)-O-CH <sub>2</sub> -O-  |   |
|    | 11 -NH-CH(CH(OH)(CH <sub>3</sub> ))-C(O)-O-CH <sub>2</sub> -O-  |   |
| 10 | 12 -NH-CH(-CH <sub>2</sub> SH)-C(O)-O-CH <sub>2</sub> -O-   |   |
|    | 13 -NH-CH(CH <sub>2</sub> -C <sub>6</sub> H <sub>5</sub> OH)-C(O)-O-CH <sub>2</sub> -O-   |   |
|    | 14 -NH-CH(CH <sub>2</sub> -C(O)-NH <sub>2</sub> )-C(O)-O-CH <sub>2</sub> -O-  |   |
|    | 15 -NH-CH(CH <sub>2</sub> -CH <sub>2</sub> -C(O)-NH <sub>2</sub> )-C(O)-O-CH <sub>2</sub> -O-   |   |
|    | 16 -NH-CH(CH <sub>2</sub> C(O)OR <sup>4</sup> )-C(O)-O-CH <sub>2</sub> -O-  |   |
| 15 | 17 -NH-CH(CH <sub>2</sub> CH <sub>2</sub> C(O)OR <sup>4</sup> )-C(O)-O-CH <sub>2</sub> -O-  |   |
|    | 18 -NH-CH(CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> NH <sub>2</sub> )-C(O)-O-CH <sub>2</sub> -O-                  |   |
|    | 19 -NH-CH(CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> NHC(NH)(NH <sub>2</sub> ))-C(O)-O-CH <sub>2</sub> -O-                         |   |
|    | 20 -NH-CH(CH <sub>2</sub> C <sub>3</sub> N <sub>2</sub> H <sub>3</sub> )-C(O)-O-CH <sub>2</sub> -O-                                     |   |
|    | 21 -NH-CH(CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> NH <sub>2</sub> )-CH <sub>2</sub> -C(O)-O-CH <sub>2</sub> -O-                 |   |
| 20 | 22 -NH-CH(CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> NH <sub>2</sub> )-CH <sub>2</sub> -C(O)-O-CH <sub>2</sub> -O- |   |
|    | 23 -NH-CH(CH <sub>2</sub> CH <sub>2</sub> NHC(NH)(NH <sub>2</sub> ))-CH <sub>2</sub> -C(O)-O-CH <sub>2</sub> -O-                        |   |
|    | 24 -NH-CH(C(O)OR <sup>4</sup> )-CH <sub>2</sub> -C(O)-O-CH <sub>2</sub> -O-   |   |
|    | 25 -NH-CH(CH <sub>2</sub> C(O)OR <sup>4</sup> )-CH <sub>2</sub> -C(O)-O-  |   |
|    | 26 -NH-CH(CH <sub>2</sub> CH <sub>2</sub> C(O)OR <sup>4</sup> )-CH <sub>2</sub> -C(O)-O-CH <sub>2</sub> -O-                             |   |
| 25 | 27 -NH-CH <sub>2</sub> -C(O)-O-CH(C(O)OR <sup>4</sup> )-N-  |   |
|    | 28 -NH-CH(CH <sub>3</sub> )-C(O)-O-CH(C(O)OR <sup>4</sup> )-N-  |   |

**B**

- |    |                          |
|----|--------------------------|
|    | 1 adenin-9-yl            |
| 30 | 2 guanin-9-yl            |
|    | 3 cytosin-1-yl           |
|    | 4 2, 6-diaminopurin-9-yl |
|    | 5 2-aminopurin-9-yl      |
|    | 6 6-azacytosin-1-yl      |
| 35 | 7 1-deazaadenin-9-yl     |

- 8 3-deazaadenin-9-yl
- 9 8-azaadenin-9-yl
- 10 7-deaza-8-azaadenin-9-yl

5 \* - See Table 1 footnote; the terminal nitrogen and oxygen or nitrogen atoms are both linked to the phosphorus atom of the phosphonate group.

\*\* - See Table 1 footnote.

# - See Table 1 footnote.

- 10 Compounds listed in Table 3 are designated herein by numbers assigned to L<sup>1</sup>, Z and B according to the following convention, L<sup>1</sup>.Z.B. Thus, compounds 1.1.1 and 2.3.4 represent compounds designated cyclic glycynylPMEA and cyclic alanylPMPDAP. Exemplary compounds include 1.1.1, 1.1.2, 1.1.3, 1.1.4, 1.1.5, 1.1.6, 1.1.7, 1.1.8, 1.1.9, 1.1.10, 2.1.1, 2.1.2, 2.1.3, 2.1.4, 2.1.5, 2.1.6, 2.1.7, 2.1.8, 2.1.9, 15 2.1.10, 3.1.1, 3.1.2, 3.1.3, 3.1.4, 3.1.5, 3.1.6, 3.1.7, 3.1.8, 3.1.9, 3.1.10, 4.1.1, 4.1.2, 4.1.3, 4.1.4, 4.1.5, 4.1.6, 4.1.7, 4.1.8, 4.1.9, 4.1.10, 5.1.1, 5.1.2, 5.1.3, 5.1.4, 5.1.5, 5.1.6, 5.1.7, 5.1.8, 5.1.9, 5.1.10, 6.1.1, 6.1.2, 6.1.3, 6.1.4, 6.1.5, 6.1.6, 6.1.7, 6.1.8, 6.1.9, 6.1.10, 7.1.1, 7.1.2, 7.1.3, 7.1.4, 7.1.5, 7.1.6, 7.1.7, 7.1.8, 7.1.9, 7.1.10, 8.1.1, 8.1.2, 8.1.3, 8.1.4, 8.1.5, 8.1.6, 8.1.7, 8.1.8, 8.1.9, 8.1.10, 9.1.1, 9.1.2, 9.1.3, 9.1.4, 9.1.5, 9.1.6, 9.1.7, 9.1.8, 9.1.9, 20 9.1.10, 10.1.1, 10.1.2, 10.1.3, 10.1.4, 10.1.5, 10.1.6, 10.1.7, 10.1.8, 10.1.9, 10.1.10, 11.1.1, 11.1.2, 11.1.3, 11.1.4, 11.1.5, 11.1.6, 11.1.7, 11.1.8, 11.1.9, 11.1.10, 12.1.1, 12.1.2, 12.1.3, 12.1.4, 12.1.5, 12.1.6, 12.1.7, 12.1.8, 12.1.9, 12.1.10, 13.1.1, 13.1.2, 13.1.3, 13.1.4, 13.1.5, 13.1.6, 13.1.7, 13.1.8, 13.1.9, 13.1.10, 14.1.1, 14.1.2, 14.1.3, 14.1.4, 14.1.5, 14.1.6, 14.1.7, 14.1.8, 14.1.9, 14.1.10, 15.1.1, 15.1.2, 15.1.3, 15.1.4, 25 15.1.5, 15.1.6, 15.1.7, 15.1.8, 15.1.9, 15.1.10, 16.1.1, 16.1.2, 16.1.3, 16.1.4, 16.1.5, 16.1.6, 16.1.7, 16.1.8, 16.1.9, 16.1.10, 17.1.1, 17.1.2, 17.1.3, 17.1.4, 17.1.5, 17.1.6, 17.1.7, 17.1.8, 17.1.9, 17.1.10, 18.1.1, 18.1.2, 18.1.3, 18.1.4, 18.1.5, 18.1.6, 18.1.7, 18.1.8, 18.1.9, 18.1.10, 19.1.1, 19.1.2, 19.1.3, 19.1.4, 19.1.5, 19.1.6, 19.1.7, 19.1.8, 19.1.9, 19.1.10, 20.1.1, 20.1.2, 20.1.3, 20.1.4, 20.1.5, 20.1.6, 20.1.7, 20.1.8, 20.1.9, 30 20.1.10, 21.1.1, 21.1.2, 21.1.3, 21.1.4, 21.1.5, 21.1.6, 21.1.7, 21.1.8, 21.1.9, 21.1.10, 22.1.1, 22.1.2, 22.1.3, 22.1.4, 22.1.5, 22.1.6, 22.1.7, 22.1.8, 22.1.9, 22.1.10, 23.1.1, 23.1.2, 23.1.3, 23.1.4, 23.1.5, 23.1.6, 23.1.7, 23.1.8, 23.1.9, 23.1.10, 24.1.1, 24.1.2, 24.1.3, 24.1.4, 24.1.5, 24.1.6, 24.1.7, 24.1.8, 24.1.9, 24.1.10, 25.1.1, 25.1.2, 25.1.3, 25.1.4, 25.1.5, 25.1.6, 25.1.7, 25.1.8, 25.1.9, 25.1.10, 26.1.1, 26.1.2, 26.1.3, 26.1.4, 35 26.1.5, 26.1.6, 26.1.7, 26.1.8, 26.1.9, 26.1.10, 27.1.1, 27.1.2, 27.1.3, 27.1.4, 27.1.5,

27.1.6, 27.1.7, 27.1.8, 27.1.9, 27.1.10, 28.1.1, 28.1.2, 28.1.3, 28.1.4, 28.1.5, 28.1.6,  
 28.1.7, 28.1.8, 28.1.9, 28.1.10, 1.2.1, 1.2.2, 1.2.3, 1.2.4, 1.2.5, 1.2.6, 1.2.7, 1.2.8, 1.2.9,  
 1.2.10, 2.2.1, 2.2.2, 2.2.3, 2.2.4, 2.2.5, 2.2.6, 2.2.7, 2.2.8, 2.2.9, 2.2.10, 3.2.1, 3.2.2, 3.2.3,  
 3.2.4, 3.2.5, 3.2.6, 3.2.7, 3.2.8, 3.2.9, 3.2.10, 4.2.1, 4.2.2, 4.2.3, 4.2.4, 4.2.5, 4.2.6, 4.2.7,  
 5 4.2.8, 4.2.9, 4.2.10, 5.2.1, 5.2.2, 5.2.3, 5.2.4, 5.2.5, 5.2.6, 5.2.7, 5.2.8, 5.2.9, 5.2.10, 6.2.1,  
 6.2.2, 6.2.3, 6.2.4, 6.2.5, 6.2.6, 6.2.7, 6.2.8, 6.2.9, 6.2.10, 7.2.1, 7.2.2, 7.2.3, 7.2.4, 7.2.5,  
 7.2.6, 7.2.7, 7.2.8, 7.2.9, 7.2.10, 8.2.1, 8.2.2, 8.2.3, 8.2.4, 8.2.5, 8.2.6, 8.2.7, 8.2.8, 8.2.9,  
 8.2.10, 9.2.1, 9.2.2, 9.2.3, 9.2.4, 9.2.5, 9.2.6, 9.2.7, 9.2.8, 9.2.9, 9.2.10, 10.2.1, 10.2.2,  
 10.2.3, 10.2.4, 10.2.5, 10.2.6, 10.2.7, 10.2.8, 10.2.9, 10.2.10, 11.2.1, 11.2.2, 11.2.3,  
 10 11.2.4, 11.2.5, 11.2.6, 11.2.7, 11.2.8, 11.2.9, 11.2.10, 12.2.1, 12.2.2, 12.2.3, 12.2.4,  
 12.2.5, 12.2.6, 12.2.7, 12.2.8, 12.2.9, 12.2.10, 13.2.1, 13.2.2, 13.2.3, 13.2.4, 13.2.5,  
 13.2.6, 13.2.7, 13.2.8, 13.2.9, 13.2.10, 14.2.1, 14.2.2, 14.2.3, 14.2.4, 14.2.5, 14.2.6,  
 14.2.7, 14.2.8, 14.2.9, 14.2.10, 15.2.1, 15.2.2, 15.2.3, 15.2.4, 15.2.5, 15.2.6, 15.2.7,  
 15.2.8, 15.2.9, 15.2.10, 16.2.1, 16.2.2, 16.2.3, 16.2.4, 16.2.5, 16.2.6, 16.2.7, 16.2.8,  
 15 16.2.9, 16.2.10, 17.2.1, 17.2.2, 17.2.3, 17.2.4, 17.2.5, 17.2.6, 17.2.7, 17.2.8, 17.2.9,  
 17.2.10, 18.2.1, 18.2.2, 18.2.3, 18.2.4, 18.2.5, 18.2.6, 18.2.7, 18.2.8, 18.2.9, 18.2.10,  
 19.2.1, 19.2.2, 19.2.3, 19.2.4, 19.2.5, 19.2.6, 19.2.7, 19.2.8, 19.2.9, 19.2.10, 20.2.1,  
 20.2.2, 20.2.3, 20.2.4, 20.2.5, 20.2.6, 20.2.7, 20.2.8, 20.2.9, 20.2.10, 21.2.1, 21.2.2,  
 21.2.3, 21.2.4, 21.2.5, 21.2.6, 21.2.7, 21.2.8, 21.2.9, 21.2.10, 22.2.1, 22.2.2, 22.2.3,  
 20 22.2.4, 22.2.5, 22.2.6, 22.2.7, 22.2.8, 22.2.9, 22.2.10, 23.2.1, 23.2.2, 23.2.3, 23.2.4,  
 23.2.5, 23.2.6, 23.2.7, 23.2.8, 23.2.9, 23.2.10, 24.2.1, 24.2.2, 24.2.3, 24.2.4, 24.2.5,  
 24.2.6, 24.2.7, 24.2.8, 24.2.9, 24.2.10, 25.2.1, 25.2.2, 25.2.3, 25.2.4, 25.2.5, 25.2.6,  
 25.2.7, 25.2.8, 25.2.9, 25.2.10, 26.2.1, 26.2.2, 26.2.3, 26.2.4, 26.2.5, 26.2.6, 26.2.7,  
 26.2.8, 26.2.9, 26.2.10, 27.2.1, 27.2.2, 27.2.3, 27.2.4, 27.2.5, 27.2.6, 27.2.7, 27.2.8,  
 25 27.2.9, 27.2.10, 28.2.1, 28.2.2, 28.2.3, 28.2.4, 28.2.5, 28.2.6, 28.2.7, 28.2.8, 28.2.9,  
 28.2.10, 1.3.1, 1.3.2, 1.3.3, 1.3.4, 1.3.5, 1.3.6, 1.3.7, 1.3.8, 1.3.9, 1.3.10, 2.3.1, 2.3.2,  
 2.3.3, 2.3.4, 2.3.5, 2.3.6, 2.3.7, 2.3.8, 2.3.9, 2.3.10, 3.3.1, 3.3.2, 3.3.3, 3.3.4, 3.3.5, 3.3.6,  
 3.3.7, 3.3.8, 3.3.9, 3.3.10, 4.3.1, 4.3.2, 4.3.3, 4.3.4, 4.3.5, 4.3.6, 4.3.7, 4.3.8, 4.3.9, 4.3.10,  
 5.3.1, 5.3.2, 5.3.3, 5.3.4, 5.3.5, 5.3.6, 5.3.7, 5.3.8, 5.3.9, 5.3.10, 6.3.1, 6.3.2, 6.3.3, 6.3.4,  
 30 6.3.5, 6.3.6, 6.3.7, 6.3.8, 6.3.9, 6.3.10, 7.3.1, 7.3.2, 7.3.3, 7.3.4, 7.3.5, 7.3.6, 7.3.7, 7.3.8,  
 7.3.9, 7.3.10, 8.3.1, 8.3.2, 8.3.3, 8.3.4, 8.3.5, 8.3.6, 8.3.7, 8.3.8, 8.3.9, 8.3.10, 9.3.1, 9.3.2,  
 9.3.3, 9.3.4, 9.3.5, 9.3.6, 9.3.7, 9.3.8, 9.3.9, 9.3.10, 10.3.1, 10.3.2, 10.3.3, 10.3.4, 10.3.5,  
 10.3.6, 10.3.7, 10.3.8, 10.3.9, 10.3.10, 11.3.1, 11.3.2, 11.3.3, 11.3.4, 11.3.5, 11.3.6,  
 11.3.7, 11.3.8, 11.3.9, 11.3.10, 12.3.1, 12.3.2, 12.3.3, 12.3.4, 12.3.5, 12.3.6, 12.3.7,  
 35 12.3.8, 12.3.9, 12.3.10, 13.3.1, 13.3.2, 13.3.3, 13.3.4, 13.3.5, 13.3.6, 13.3.7, 13.3.8,

13.3.9, 13.3.10, 14.3.1, 14.3.2, 14.3.3, 14.3.4, 14.3.5, 14.3.6, 14.3.7, 14.3.8, 14.3.9, 14.3.10, 15.3.1, 15.3.2, 15.3.3, 15.3.4, 15.3.5, 15.3.6, 15.3.7, 15.3.8, 15.3.9, 15.3.10, 16.3.1, 16.3.2, 16.3.3, 16.3.4, 16.3.5, 16.3.6, 16.3.7, 16.3.8, 16.3.9, 16.3.10, 17.3.1, 17.3.2, 17.3.3, 17.3.4, 17.3.5, 17.3.6, 17.3.7, 17.3.8, 17.3.9, 17.3.10, 18.3.1, 18.3.2, 18.3.3, 18.3.4, 18.3.5, 18.3.6, 18.3.7, 18.3.8, 18.3.9, 18.3.10, 19.3.1, 19.3.2, 19.3.3, 19.3.4, 19.3.5, 19.3.6, 19.3.7, 19.3.8, 19.3.9, 19.3.10, 20.3.1, 20.3.2, 20.3.3, 20.3.4, 20.3.5, 20.3.6, 20.3.7, 20.3.8, 20.3.9, 20.3.10, 21.3.1, 21.3.2, 21.3.3, 21.3.4, 21.3.5, 21.3.6, 21.3.7, 21.3.8, 21.3.9, 21.3.10, 22.3.1, 22.3.2, 22.3.3, 22.3.4, 22.3.5, 22.3.6, 22.3.7, 22.3.8, 22.3.9, 22.3.10, 23.3.1, 23.3.2, 23.3.3, 23.3.4, 23.3.5, 23.3.6, 23.3.7, 23.3.8, 23.3.9, 23.3.10, 24.3.1, 24.3.2, 24.3.3, 24.3.4, 24.3.5, 24.3.6, 24.3.7, 24.3.8, 24.3.9, 24.3.10, 25.3.1, 25.3.2, 25.3.3, 25.3.4, 25.3.5, 25.3.6, 25.3.7, 25.3.8, 25.3.9, 25.3.10, 26.3.1, 26.3.2, 26.3.3, 26.3.4, 26.3.5, 26.3.6, 26.3.7, 26.3.8, 26.3.9, 26.3.10, 27.3.1, 27.3.2, 27.3.3, 27.3.4, 27.3.5, 27.3.6, 27.3.7, 27.3.8, 27.3.9, 27.3.10, 28.3.1, 28.3.2, 28.3.3, 28.3.4, 28.3.5, 28.3.6, 28.3.7, 28.3.8, 28.3.9 and 28.3.10.

15

Table 4 lists a group of cyclic nucleotide analogs of structure I wherein a heterocyclic ring comprising  $L^1$  and the phosphorus atom of the phosphonate group along with part of the Z-B substructure  $-O-CH_2-C^{\#}H(CH_2-)-CH_2-B$ . The unbonded O atom in the Z substructure is linked to  $L^1$  through the  $\alpha$  carboxyl group of the amino acid while the  $CH_2$  moiety on the right side is linked to the P atom and the  $CH_2$  moiety linked to the chiral carbon is linked to B (i.e.,  $-L^1-O-CH_2-C^{\#}H(CH_2-B)-O-CH_2-P(O)(L^2)-$  with  $-P(O)(L^2)-$  and  $-L^1-$  linked together). Hydrolysis of the compound results in formation of an HPMP nucleoside phosphonate. A related group of compounds comprises a heterocyclic ring linked through a side chain or other carboxyl group instead of through the carboxyl group linked to the  $\alpha$  carbon atom. Hydrolysis of these compounds also result in formation of an HPMP nucleoside phosphonate.

30

TABLE 4

---

 $L^1*-Z(B)-P(O)(L^2)$ 
1  $-NH-CH_2-C(O)-O-CH_2-C^{\#}H(CH_2-B)-O-CH_2-P(O)(L^2)-$ 2  $-NH-CH(CH_3)-C(O)-O-CH_2-C^{\#}H(CH_2-B)-O-CH_2-P(O)(L^2)-$ 35 3  $-NH-CH(CH_3)_2-C(O)-O-CH_2-C^{\#}H(CH_2-B)-O-CH_2-P(O)(L^2)-$

- 4 -NH-CH(CH(CH<sub>3</sub>)<sub>2</sub>)-C(O)-O-CH<sub>2</sub>-C#H(CH<sub>2</sub>-B)-O-CH<sub>2</sub>-P(O)(L<sup>2</sup>)-  
 5 -NH-CH(CH<sub>3</sub>)(CH<sub>3</sub>)<sub>2</sub>-C(O)-O-CH<sub>2</sub>-C#H(CH<sub>2</sub>-B)-O-CH<sub>2</sub>-P(O)(L<sup>2</sup>)-  
 6 -NH-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH-C(O)-O-CH<sub>2</sub>-C#H(CH<sub>2</sub>-B)-O-CH<sub>2</sub>-P(O)(L<sup>2</sup>)-  
 7 -NH-CH(CH<sub>2</sub>-C<sub>6</sub>H<sub>5</sub>)-C(O)-O-CH<sub>2</sub>-C#H(CH<sub>2</sub>-B)-O-CH<sub>2</sub>-P(O)(L<sup>2</sup>)-  
 5 8 -NH-CH(CH<sub>2</sub>-C<sub>8</sub>NH<sub>6</sub>)-C(O)-O-CH<sub>2</sub>-C#H(CH<sub>2</sub>-B)-O-CH<sub>2</sub>-P(O)(L<sup>2</sup>)-  
 9 -NH-CH(CH<sub>2</sub>-CH<sub>2</sub>-S-CH<sub>3</sub>)-C(O)-O-CH<sub>2</sub>-C#H(CH<sub>2</sub>-B)-O-CH<sub>2</sub>-P(O)(L<sup>2</sup>)-  
 10 -NH-CH(CH<sub>2</sub>OH)-C(O)-O-CH<sub>2</sub>-O-CH<sub>2</sub>-C#H(CH<sub>2</sub>-B)-O-CH<sub>2</sub>-P(O)(L<sup>2</sup>)-  
 11 -NH-CH(CH(OH)(CH<sub>3</sub>))-C(O)-O-CH<sub>2</sub>-C#H(CH<sub>2</sub>-B)-O-CH<sub>2</sub>-P(O)(L<sup>2</sup>)-  
 12 -NH-CH(-CH<sub>2</sub>SH)-C(O)-O-CH<sub>2</sub>-O-CH<sub>2</sub>-C#H(CH<sub>2</sub>-B)-O-CH<sub>2</sub>-P(O)(L<sup>2</sup>)-  
 10 13 -NH-CH(CH<sub>2</sub>-C<sub>6</sub>H<sub>5</sub>OH)-C(O)-O-CH<sub>2</sub>-C#H(CH<sub>2</sub>-B)-O-CH<sub>2</sub>-P(O)(L<sup>2</sup>)-  
 14 -NH-CH(CH<sub>2</sub>-C(O)-NH<sub>2</sub>)-C(O)-O-CH<sub>2</sub>-C#H(CH<sub>2</sub>-B)-O-CH<sub>2</sub>-P(O)(L<sup>2</sup>)-  
 15 -NH-CH(CH<sub>2</sub>-CH<sub>2</sub>-C(O)-NH<sub>2</sub>)-C(O)-O-CH<sub>2</sub>-C#H(CH<sub>2</sub>-B)-O-CH<sub>2</sub>-P(O)(L<sup>2</sup>)-  
 16 -NH-CH(CH<sub>2</sub>C(O)OR<sup>4</sup>)-C(O)-O-CH<sub>2</sub>-C#H(CH<sub>2</sub>-B)-O-CH<sub>2</sub>-P(O)(L<sup>2</sup>)-  
 17 -NH-CH(CH<sub>2</sub>CH<sub>2</sub>C(O)OR<sup>4</sup>)-C(O)-O-CH<sub>2</sub>-C#H(CH<sub>2</sub>-B)-O-CH<sub>2</sub>-P(O)(L<sup>2</sup>)-  
 15 18 -NH-CH(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>)-C(O)-O-CH<sub>2</sub>-C#H(CH<sub>2</sub>-B)-O-CH<sub>2</sub>-P(O)(L<sup>2</sup>)-  
 19 -NH-CH(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NHC(NH)(NH<sub>2</sub>))-C(O)-O-CH<sub>2</sub>-C#H(CH<sub>2</sub>-B)-O-CH<sub>2</sub>-  
 P(O)(L<sup>2</sup>)-  
 20 -NH-CH(CH<sub>2</sub>C<sub>3</sub>N<sub>2</sub>H<sub>3</sub>)-C(O)-O-CH<sub>2</sub>-C#H(CH<sub>2</sub>-B)-O-CH<sub>2</sub>-P(O)(L<sup>2</sup>)-  
 21 -NH-CH(CH<sub>3</sub>)-CH<sub>2</sub>-C(O)-O-CH<sub>2</sub>-C#H(CH<sub>2</sub>-B)-O-CH<sub>2</sub>-P(O)(L<sup>2</sup>)-  
 20 22 -NH-CH(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>)-CH<sub>2</sub>-C(O)-O-CH<sub>2</sub>-C#H(CH<sub>2</sub>-B)-O-CH<sub>2</sub>-P(O)(L<sup>2</sup>)-

**L<sup>2</sup>**

- 1 -NH-CH<sub>2</sub>-C(O)-OR<sup>4</sup>  
 2 -NH-CH(CH<sub>3</sub>)-C(O)-OR<sup>4</sup>  
 25 3 -O-CH<sub>2</sub>-O-C(O)-C(CH<sub>3</sub>)<sub>3</sub>  
 4 -O-CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>  
 5 -O-C<sub>6</sub>H<sub>5</sub>  
 6 -O-CH(CH<sub>3</sub>)<sub>2</sub>  
 7 -NH-CH(CH<sub>2</sub>C<sub>6</sub>H<sub>4</sub>)-C(O)-OR<sup>4</sup>  
 30 8 -OH

**B**

1. adenin-9-yl  
 2. guanin-9-yl  
 35 3. cytosin-1-yl

- |                           |                              |
|---------------------------|------------------------------|
| 4. 2, 6-diaminopurin-9-yl |                              |
| 5. 2-aminopurin-9-yl      | 8. 3-deazaadenin-9-yl        |
| 6. 6-azacytosin-1-yl      | 9. 8-azaadenin-9-yl          |
| 7. 1-deazaadenin-9-yl     | 10. 7-deaza-8-azaadenin-9-yl |
- 

- 5 \* - See Table 1 footnote; the terminal nitrogen and phosphorus atoms are linked to each other.

- Compounds listed in Table 4 are designated herein by numbers assigned to L<sup>1</sup>, L<sup>2</sup>, and B according to the following convention, L<sup>1</sup>.L<sup>2</sup>.B. All Z correspond to the esterified HPMP substructure moiety. Thus, compounds 1.1.3 and 2.4.3 represent compounds designated "glycyl cyclic glycinyl HPMP" and "benzyl cyclic alanyl HPMP" esters. Exemplary compounds include
- 1.1.1, 1.1.2, 1.1.3, 1.1.4, 1.1.5, 1.1.6, 1.1.7, 1.1.8, 1.1.9, 1.1.10, 2.1.1, 2.1.2, 2.1.3, 2.1.4, 2.1.5, 2.1.6, 2.1.7, 2.1.8, 2.1.9, 2.1.10, 3.1.1, 3.1.2, 3.1.3, 3.1.4, 3.1.5, 3.1.6, 3.1.7, 3.1.8, 3.1.9, 3.1.10, 4.1.1, 4.1.2, 4.1.3, 4.1.4, 4.1.5, 4.1.6, 4.1.7, 4.1.8, 4.1.9, 4.1.10, 5.1.1, 5.1.2, 5.1.3, 5.1.4, 5.1.5, 5.1.6, 5.1.7, 5.1.8, 5.1.9, 5.1.10, 6.1.1, 6.1.2, 6.1.3, 6.1.4, 6.1.5, 6.1.6, 6.1.7, 6.1.8, 6.1.9, 6.1.10, 7.1.1, 7.1.2, 7.1.3, 7.1.4, 7.1.5, 7.1.6, 7.1.7, 7.1.8, 7.1.9, 7.1.10, 8.1.1, 8.1.2, 8.1.3, 8.1.4, 8.1.5, 8.1.6, 8.1.7, 8.1.8, 8.1.9, 8.1.10, 9.1.1, 9.1.2, 9.1.3, 9.1.4, 9.1.5, 9.1.6, 9.1.7, 9.1.8, 9.1.9, 9.1.10, 10.1.1, 10.1.2, 10.1.3, 10.1.4, 10.1.5, 10.1.6, 10.1.7, 10.1.8, 10.1.9, 10.1.10, 11.1.1, 11.1.2, 11.1.3, 11.1.4, 11.1.5, 11.1.6, 11.1.7, 11.1.8, 11.1.9, 11.1.10, 12.1.1, 12.1.2, 12.1.3, 12.1.4, 12.1.5, 12.1.6, 12.1.7, 12.1.8, 12.1.9, 12.1.10, 13.1.1, 13.1.2, 13.1.3, 13.1.4, 13.1.5, 13.1.6, 13.1.7, 13.1.8, 13.1.9, 13.1.10, 14.1.1, 14.1.2, 14.1.3, 14.1.4, 14.1.5, 14.1.6, 14.1.7, 14.1.8, 14.1.9, 14.1.10, 15.1.1, 15.1.2, 15.1.3, 15.1.4, 15.1.5, 15.1.6, 15.1.7, 15.1.8, 15.1.9, 15.1.10, 16.1.1, 16.1.2, 16.1.3, 16.1.4, 16.1.5, 16.1.6, 16.1.7, 16.1.8, 16.1.9, 16.1.10, 17.1.1, 17.1.2, 17.1.3, 17.1.4, 17.1.5, 17.1.6, 17.1.7, 17.1.8, 17.1.9, 17.1.10, 18.1.1, 18.1.2, 18.1.3, 18.1.4, 18.1.5, 18.1.6, 18.1.7, 18.1.8, 18.1.9, 18.1.10, 19.1.1, 19.1.2, 19.1.3, 19.1.4, 19.1.5, 19.1.6, 19.1.7, 19.1.8, 19.1.9, 19.1.10, 20.1.1, 20.1.2, 20.1.3, 20.1.4, 20.1.5, 20.1.6, 20.1.7, 20.1.8, 20.1.9, 20.1.10, 21.1.1, 21.1.2, 21.1.3, 21.1.4, 21.1.5, 21.1.6, 21.1.7, 21.1.8, 21.1.9, 21.1.10, 22.1.1, 22.1.2, 22.1.3, 22.1.4, 22.1.5, 22.1.6, 22.1.7, 22.1.8, 22.1.9, 22.1.10, 1.2.1, 1.2.2, 1.2.3, 1.2.4, 1.2.5, 1.2.6, 1.2.7, 1.2.8, 1.2.9, 1.2.10, 2.2.1, 2.2.2, 2.2.3, 2.2.4, 2.2.5, 2.2.6, 2.2.7, 2.2.8, 2.2.9, 2.2.10, 3.2.1, 3.2.2, 3.2.3, 3.2.4, 3.2.5, 3.2.6, 3.2.7, 3.2.8, 3.2.9, 3.2.10, 4.2.1, 4.2.2, 4.2.3, 4.2.4, 4.2.5, 4.2.6, 4.2.7, 4.2.8, 4.2.9, 4.2.10, 5.2.1, 5.2.2, 5.2.3, 5.2.4, 5.2.5, 5.2.6, 5.2.7, 5.2.8, 5.2.9, 5.2.10, 6.2.1, 6.2.2, 6.2.3, 6.2.4, 6.2.5, 6.2.6, 6.2.7, 6.2.8, 6.2.9, 6.2.10, 7.2.1, 7.2.2, 7.2.3, 7.2.4, 7.2.5, 7.2.6,

7.2.7, 7.2.8, 7.2.9, 7.2.10, 8.2.1, 8.2.2, 8.2.3, 8.2.4, 8.2.5, 8.2.6, 8.2.7, 8.2.8, 8.2.9, 8.2.10,  
 9.2.1, 9.2.2, 9.2.3, 9.2.4, 9.2.5, 9.2.6, 9.2.7, 9.2.8, 9.2.9, 9.2.10, 10.2.1, 10.2.2, 10.2.3,  
 10.2.4, 10.2.5, 10.2.6, 10.2.7, 10.2.8, 10.2.9, 10.2.10, 11.2.1, 11.2.2, 11.2.3, 11.2.4,  
 11.2.5, 11.2.6, 11.2.7, 11.2.8, 11.2.9, 11.2.10, 12.2.1, 12.2.2, 12.2.3, 12.2.4, 12.2.5,  
 5 12.2.6, 12.2.7, 12.2.8, 12.2.9, 12.2.10, 13.2.1, 13.2.2, 13.2.3, 13.2.4, 13.2.5, 13.2.6,  
 13.2.7, 13.2.8, 13.2.9, 13.2.10, 14.2.1, 14.2.2, 14.2.3, 14.2.4, 14.2.5, 14.2.6, 14.2.7,  
 14.2.8, 14.2.9, 14.2.10, 15.2.1, 15.2.2, 15.2.3, 15.2.4, 15.2.5, 15.2.6, 15.2.7, 15.2.8,  
 15.2.9, 15.2.10, 16.2.1, 16.2.2, 16.2.3, 16.2.4, 16.2.5, 16.2.6, 16.2.7, 16.2.8, 16.2.9,  
 16.2.10, 17.2.1, 17.2.2, 17.2.3, 17.2.4, 17.2.5, 17.2.6, 17.2.7, 17.2.8, 17.2.9, 17.2.10,  
 10 18.2.1, 18.2.2, 18.2.3, 18.2.4, 18.2.5, 18.2.6, 18.2.7, 18.2.8, 18.2.9, 18.2.10, 19.2.1,  
 19.2.2, 19.2.3, 19.2.4, 19.2.5, 19.2.6, 19.2.7, 19.2.8, 19.2.9, 19.2.10, 20.2.1, 20.2.2,  
 20.2.3, 20.2.4, 20.2.5, 20.2.6, 20.2.7, 20.2.8, 20.2.9, 20.2.10, 21.2.1, 21.2.2, 21.2.3,  
 21.2.4, 21.2.5, 21.2.6, 21.2.7, 21.2.8, 21.2.9, 21.2.10, 22.2.1, 22.2.2, 22.2.3, 22.2.4,  
 22.2.5, 22.2.6, 22.2.7, 22.2.8, 22.2.9, 22.2.10, 1.3.1, 1.3.2, 1.3.3, 1.3.4, 1.3.5, 1.3.6, 1.3.7,  
 15 1.3.8, 1.3.9, 1.3.10, 2.3.1, 2.3.2, 2.3.3, 2.3.4, 2.3.5, 2.3.6, 2.3.7, 2.3.8, 2.3.9, 2.3.10, 3.3.1,  
 3.3.2, 3.3.3, 3.3.4, 3.3.5, 3.3.6, 3.3.7, 3.3.8, 3.3.9, 3.3.10, 4.3.1, 4.3.2, 4.3.3, 4.3.4, 4.3.5,  
 4.3.6, 4.3.7, 4.3.8, 4.3.9, 4.3.10, 5.3.1, 5.3.2, 5.3.3, 5.3.4, 5.3.5, 5.3.6, 5.3.7, 5.3.8, 5.3.9,  
 5.3.10, 6.3.1, 6.3.2, 6.3.3, 6.3.4, 6.3.5, 6.3.6, 6.3.7, 6.3.8, 6.3.9, 6.3.10, 7.3.1, 7.3.2, 7.3.3,  
 7.3.4, 7.3.5, 7.3.6, 7.3.7, 7.3.8, 7.3.9, 7.3.10, 8.3.1, 8.3.2, 8.3.3, 8.3.4, 8.3.5, 8.3.6, 8.3.7,  
 20 8.3.8, 8.3.9, 8.3.10, 9.3.1, 9.3.2, 9.3.3, 9.3.4, 9.3.5, 9.3.6, 9.3.7, 9.3.8, 9.3.9, 9.3.10,  
 10.3.1, 10.3.2, 10.3.3, 10.3.4, 10.3.5, 10.3.6, 10.3.7, 10.3.8, 10.3.9, 10.3.10, 11.3.1,  
 11.3.2, 11.3.3, 11.3.4, 11.3.5, 11.3.6, 11.3.7, 11.3.8, 11.3.9, 11.3.10, 12.3.1, 12.3.2,  
 12.3.3, 12.3.4, 12.3.5, 12.3.6, 12.3.7, 12.3.8, 12.3.9, 12.3.10, 13.3.1, 13.3.2, 13.3.3,  
 13.3.4, 13.3.5, 13.3.6, 13.3.7, 13.3.8, 13.3.9, 13.3.10, 14.3.1, 14.3.2, 14.3.3, 14.3.4,  
 25 14.3.5, 14.3.6, 14.3.7, 14.3.8, 14.3.9, 14.3.10, 15.3.1, 15.3.2, 15.3.3, 15.3.4, 15.3.5,  
 15.3.6, 15.3.7, 15.3.8, 15.3.9, 15.3.10, 16.3.1, 16.3.2, 16.3.3, 16.3.4, 16.3.5, 16.3.6,  
 16.3.7, 16.3.8, 16.3.9, 16.3.10, 17.3.1, 17.3.2, 17.3.3, 17.3.4, 17.3.5, 17.3.6, 17.3.7,  
 17.3.8, 17.3.9, 17.3.10, 18.3.1, 18.3.2, 18.3.3, 18.3.4, 18.3.5, 18.3.6, 18.3.7, 18.3.8,  
 18.3.9, 18.3.10, 19.3.1, 19.3.2, 19.3.3, 19.3.4, 19.3.5, 19.3.6, 19.3.7, 19.3.8, 19.3.9,  
 30 19.3.10, 20.3.1, 20.3.2, 20.3.3, 20.3.4, 20.3.5, 20.3.6, 20.3.7, 20.3.8, 20.3.9, 20.3.10,  
 21.3.1, 21.3.2, 21.3.3, 21.3.4, 21.3.5, 21.3.6, 21.3.7, 21.3.8, 21.3.9, 21.3.10, 22.3.1,  
 22.3.2, 22.3.3, 22.3.4, 22.3.5, 22.3.6, 22.3.7, 22.3.8, 22.3.9, 22.3.10, 1.4.1, 1.4.2, 1.4.3,  
 1.4.4, 1.4.5, 1.4.6, 1.4.7, 1.4.8, 1.4.9, 1.4.10, 2.4.1, 2.4.2, 2.4.3, 2.4.4, 2.4.5, 2.4.6, 2.4.7,  
 2.4.8, 2.4.9, 2.4.10, 3.4.1, 3.4.2, 3.4.3, 3.4.4, 3.4.5, 3.4.6, 3.4.7, 3.4.8, 3.4.9, 3.4.10, 4.4.1,  
 35 4.4.2, 4.4.3, 4.4.4, 4.4.5, 4.4.6, 4.4.7, 4.4.8, 4.4.9, 4.4.10, 5.4.1, 5.4.2, 5.4.3, 5.4.4, 5.4.5,

5.4.6, 5.4.7, 5.4.8, 5.4.9, 5.4.10, 6.4.1, 6.4.2, 6.4.3, 6.4.4, 6.4.5, 6.4.6, 6.4.7, 6.4.8, 6.4.9,  
 6.4.10, 7.4.1, 7.4.2, 7.4.3, 7.4.4, 7.4.5, 7.4.6, 7.4.7, 7.4.8, 7.4.9, 7.4.10, 8.4.1, 8.4.2, 8.4.3,  
 8.4.4, 8.4.5, 8.4.6, 8.4.7, 8.4.8, 8.4.9, 8.4.10, 9.4.1, 9.4.2, 9.4.3, 9.4.4, 9.4.5, 9.4.6, 9.4.7,  
 9.4.8, 9.4.9, 9.4.10, 10.4.1, 10.4.2, 10.4.3, 10.4.4, 10.4.5, 10.4.6, 10.4.7, 10.4.8, 10.4.9,  
 5 10.4.10, 11.4.1, 11.4.2, 11.4.3, 11.4.4, 11.4.5, 11.4.6, 11.4.7, 11.4.8, 11.4.9, 11.4.10,  
 12.4.1, 12.4.2, 12.4.3, 12.4.4, 12.4.5, 12.4.6, 12.4.7, 12.4.8, 12.4.9, 12.4.10, 13.4.1,  
 13.4.2, 13.4.3, 13.4.4, 13.4.5, 13.4.6, 13.4.7, 13.4.8, 13.4.9, 13.4.10, 14.4.1, 14.4.2,  
 14.4.3, 14.4.4, 14.4.5, 14.4.6, 14.4.7, 14.4.8, 14.4.9, 14.4.10, 15.4.1, 15.4.2, 15.4.3,  
 15.4.4, 15.4.5, 15.4.6, 15.4.7, 15.4.8, 15.4.9, 15.4.10, 16.4.1, 16.4.2, 16.4.3, 16.4.4,  
 10 16.4.5, 16.4.6, 16.4.7, 16.4.8, 16.4.9, 16.4.10, 17.4.1, 17.4.2, 17.4.3, 17.4.4, 17.4.5,  
 17.4.6, 17.4.7, 17.4.8, 17.4.9, 17.4.10, 18.4.1, 18.4.2, 18.4.3, 18.4.4, 18.4.5, 18.4.6,  
 18.4.7, 18.4.8, 18.4.9, 18.4.10, 19.4.1, 19.4.2, 19.4.3, 19.4.4, 19.4.5, 19.4.6, 19.4.7,  
 19.4.8, 19.4.9, 19.4.10, 20.4.1, 20.4.2, 20.4.3, 20.4.4, 20.4.5, 20.4.6, 20.4.7, 20.4.8,  
 20.4.9, 20.4.10, 21.4.1, 21.4.2, 21.4.3, 21.4.4, 21.4.5, 21.4.6, 21.4.7, 21.4.8, 21.4.9,  
 15 21.4.10, 22.4.1, 22.4.2, 22.4.3, 22.4.4, 22.4.5, 22.4.6, 22.4.7, 22.4.8, 22.4.9, 22.4.10,  
 1.5.1, 1.5.2, 1.5.3, 1.5.4, 1.5.5, 1.5.6, 1.5.7, 1.5.8, 1.5.9, 1.5.10, 2.5.1, 2.5.2, 2.5.3, 2.5.4,  
 2.5.5, 2.5.6, 2.5.7, 2.5.8, 2.5.9, 2.5.10, 3.5.1, 3.5.2, 3.5.3, 3.5.4, 3.5.5, 3.5.6, 3.5.7, 3.5.8,  
 3.5.9, 3.5.10, 4.5.1, 4.5.2, 4.5.3, 4.5.4, 4.5.5, 4.5.6, 4.5.7, 4.5.8, 4.5.9, 4.5.10, 5.5.1, 5.5.2,  
 5.5.3, 5.5.4, 5.5.5, 5.5.6, 5.5.7, 5.5.8, 5.5.9, 5.5.10, 6.5.1, 6.5.2, 6.5.3, 6.5.4, 6.5.5, 6.5.6,  
 20 6.5.7, 6.5.8, 6.5.9, 6.5.10, 7.5.1, 7.5.2, 7.5.3, 7.5.4, 7.5.5, 7.5.6, 7.5.7, 7.5.8, 7.5.9, 7.5.10,  
 8.5.1, 8.5.2, 8.5.3, 8.5.4, 8.5.5, 8.5.6, 8.5.7, 8.5.8, 8.5.9, 8.5.10, 9.5.1, 9.5.2, 9.5.3, 9.5.4,  
 9.5.5, 9.5.6, 9.5.7, 9.5.8, 9.5.9, 9.5.10, 10.5.1, 10.5.2, 10.5.3, 10.5.4, 10.5.5, 10.5.6,  
 10.5.7, 10.5.8, 10.5.9, 10.5.10, 11.5.1, 11.5.2, 11.5.3, 11.5.4, 11.5.5, 11.5.6, 11.5.7,  
 11.5.8, 11.5.9, 11.5.10, 12.5.1, 12.5.2, 12.5.3, 12.5.4, 12.5.5, 12.5.6, 12.5.7, 12.5.8,  
 25 12.5.9, 12.5.10, 13.5.1, 13.5.2, 13.5.3, 13.5.4, 13.5.5, 13.5.6, 13.5.7, 13.5.8, 13.5.9,  
 13.5.10, 14.5.1, 14.5.2, 14.5.3, 14.5.4, 14.5.5, 14.5.6, 14.5.7, 14.5.8, 14.5.9, 14.5.10,  
 15.5.1, 15.5.2, 15.5.3, 15.5.4, 15.5.5, 15.5.6, 15.5.7, 15.5.8, 15.5.9, 15.5.10, 16.5.1,  
 16.5.2, 16.5.3, 16.5.4, 16.5.5, 16.5.6, 16.5.7, 16.5.8, 16.5.9, 16.5.10, 17.5.1, 17.5.2,  
 17.5.3, 17.5.4, 17.5.5, 17.5.6, 17.5.7, 17.5.8, 17.5.9, 17.5.10, 18.5.1, 18.5.2, 18.5.3,  
 30 18.5.4, 18.5.5, 18.5.6, 18.5.7, 18.5.8, 18.5.9, 18.5.10, 19.5.1, 19.5.2, 19.5.3, 19.5.4,  
 19.5.5, 19.5.6, 19.5.7, 19.5.8, 19.5.9, 19.5.10, 20.5.1, 20.5.2, 20.5.3, 20.5.4, 20.5.5,  
 20.5.6, 20.5.7, 20.5.8, 20.5.9, 20.5.10, 21.5.1, 21.5.2, 21.5.3, 21.5.4, 21.5.5, 21.5.6,  
 21.5.7, 21.5.8, 21.5.9, 21.5.10, 22.5.1, 22.5.2, 22.5.3, 22.5.4, 22.5.5, 22.5.6, 22.5.7,  
 22.5.8, 22.5.9, 22.5.10, 1.6.1, 1.6.2, 1.6.3, 1.6.4, 1.6.5, 1.6.6, 1.6.7, 1.6.8, 1.6.9, 1.6.10,  
 35 2.6.1, 2.6.2, 2.6.3, 2.6.4, 2.6.5, 2.6.6, 2.6.7, 2.6.8, 2.6.9, 2.6.10, 3.6.1, 3.6.2, 3.6.3, 3.6.4,



3.6.5, 3.6.6, 3.6.7, 3.6.8, 3.6.9, 3.6.10, 4.6.1, 4.6.2, 4.6.3, 4.6.4, 4.6.5, 4.6.6, 4.6.7, 4.6.8,  
 4.6.9, 4.6.10, 5.6.1, 5.6.2, 5.6.3, 5.6.4, 5.6.5, 5.6.6, 5.6.7, 5.6.8, 5.6.9, 5.6.10, 6.6.1, 6.6.2,  
 6.6.3, 6.6.4, 6.6.5, 6.6.6, 6.6.7, 6.6.8, 6.6.9, 6.6.10, 7.6.1, 7.6.2, 7.6.3, 7.6.4, 7.6.5, 7.6.6,  
 7.6.7, 7.6.8, 7.6.9, 7.6.10, 8.6.1, 8.6.2, 8.6.3, 8.6.4, 8.6.5, 8.6.6, 8.6.7, 8.6.8, 8.6.9, 8.6.10,  
 5 9.6.1, 9.6.2, 9.6.3, 9.6.4, 9.6.5, 9.6.6, 9.6.7, 9.6.8, 9.6.9, 9.6.10, 10.6.1, 10.6.2, 10.6.3,  
 10.6.4, 10.6.5, 10.6.6, 10.6.7, 10.6.8, 10.6.9, 10.6.10, 11.6.1, 11.6.2, 11.6.3, 11.6.4,  
 11.6.5, 11.6.6, 11.6.7, 11.6.8, 11.6.9, 11.6.10, 12.6.1, 12.6.2, 12.6.3, 12.6.4, 12.6.5,  
 12.6.6, 12.6.7, 12.6.8, 12.6.9, 12.6.10, 13.6.1, 13.6.2, 13.6.3, 13.6.4, 13.6.5, 13.6.6,  
 13.6.7, 13.6.8, 13.6.9, 13.6.10, 14.6.1, 14.6.2, 14.6.3, 14.6.4, 14.6.5, 14.6.6, 14.6.7,  
 10 14.6.8, 14.6.9, 14.6.10, 15.6.1, 15.6.2, 15.6.3, 15.6.4, 15.6.5, 15.6.6, 15.6.7, 15.6.8,  
 15.6.9, 15.6.10, 16.6.1, 16.6.2, 16.6.3, 16.6.4, 16.6.5, 16.6.6, 16.6.7, 16.6.8, 16.6.9,  
 16.6.10, 17.6.1, 17.6.2, 17.6.3, 17.6.4, 17.6.5, 17.6.6, 17.6.7, 17.6.8, 17.6.9, 17.6.10,  
 18.6.1, 18.6.2, 18.6.3, 18.6.4, 18.6.5, 18.6.6, 18.6.7, 18.6.8, 18.6.9, 18.6.10, 19.6.1,  
 19.6.2, 19.6.3, 19.6.4, 19.6.5, 19.6.6, 19.6.7, 19.6.8, 19.6.9, 19.6.10, 20.6.1, 20.6.2,  
 15 20.6.3, 20.6.4, 20.6.5, 20.6.6, 20.6.7, 20.6.8, 20.6.9, 20.6.10, 21.6.1, 21.6.2, 21.6.3,  
 21.6.4, 21.6.5, 21.6.6, 21.6.7, 21.6.8, 21.6.9, 21.6.10, 22.6.1, 22.6.2, 22.6.3, 22.6.4,  
 22.6.5, 22.6.6, 22.6.7, 22.6.8, 22.6.9, 22.6.10, 1.7.1, 1.7.2, 1.7.3, 1.7.4, 1.7.5, 1.7.6, 1.7.7,  
 1.7.8, 1.7.9, 1.7.10, 2.7.1, 2.7.2, 2.7.3, 2.7.4, 2.7.5, 2.7.6, 2.7.7, 2.7.8, 2.7.9, 2.7.10, 3.7.1,  
 3.7.2, 3.7.3, 3.7.4, 3.7.5, 3.7.6, 3.7.7, 3.7.8, 3.7.9, 3.7.10, 4.7.1, 4.7.2, 4.7.3, 4.7.4, 4.7.5,  
 20 4.7.6, 4.7.7, 4.7.8, 4.7.9, 4.7.10, 5.7.1, 5.7.2, 5.7.3, 5.7.4, 5.7.5, 5.7.6, 5.7.7, 5.7.8, 5.7.9,  
 5.7.10, 6.7.1, 6.7.2, 6.7.3, 6.7.4, 6.7.5, 6.7.6, 6.7.7, 6.7.8, 6.7.9, 6.7.10, 7.7.1, 7.7.2, 7.7.3,  
 7.7.4, 7.7.5, 7.7.6, 7.7.7, 7.7.8, 7.7.9, 7.7.10, 8.7.1, 8.7.2, 8.7.3, 8.7.4, 8.7.5, 8.7.6, 8.7.7,  
 8.7.8, 8.7.9, 8.7.10, 9.7.1, 9.7.2, 9.7.3, 9.7.4, 9.7.5, 9.7.6, 9.7.7, 9.7.8, 9.7.9, 9.7.10,  
 10.7.1, 10.7.2, 10.7.3, 10.7.4, 10.7.5, 10.7.6, 10.7.7, 10.7.8, 10.7.9, 10.7.10, 11.7.1,  
 25 11.7.2, 11.7.3, 11.7.4, 11.7.5, 11.7.6, 11.7.7, 11.7.8, 11.7.9, 11.7.10, 12.7.1, 12.7.2,  
 12.7.3, 12.7.4, 12.7.5, 12.7.6, 12.7.7, 12.7.8, 12.7.9, 12.7.10, 13.7.1, 13.7.2, 13.7.3,  
 13.7.4, 13.7.5, 13.7.6, 13.7.7, 13.7.8, 13.7.9, 13.7.10, 14.7.1, 14.7.2, 14.7.3, 14.7.4,  
 14.7.5, 14.7.6, 14.7.7, 14.7.8, 14.7.9, 14.7.10, 15.7.1, 15.7.2, 15.7.3, 15.7.4, 15.7.5,  
 15.7.6, 15.7.7, 15.7.8, 15.7.9, 15.7.10, 16.7.1, 16.7.2, 16.7.3, 16.7.4, 16.7.5, 16.7.6,  
 30 16.7.7, 16.7.8, 16.7.9, 16.7.10, 17.7.1, 17.7.2, 17.7.3, 17.7.4, 17.7.5, 17.7.6, 17.7.7,  
 17.7.8, 17.7.9, 17.7.10, 18.7.1, 18.7.2, 18.7.3, 18.7.4, 18.7.5, 18.7.6, 18.7.7, 18.7.8,  
 18.7.9, 18.7.10, 19.7.1, 19.7.2, 19.7.3, 19.7.4, 19.7.5, 19.7.6, 19.7.7, 19.7.8, 19.7.9,  
 19.7.10, 20.7.1, 20.7.2, 20.7.3, 20.7.4, 20.7.5, 20.7.6, 20.7.7, 20.7.8, 20.7.9, 20.7.10,  
 21.7.1, 21.7.2, 21.7.3, 21.7.4, 21.7.5, 21.7.6, 21.7.7, 21.7.8, 21.7.9, 21.7.10, 22.7.1,  
 35 22.7.2, 22.7.3, 22.7.4, 22.7.5, 22.7.6, 22.7.7, 22.7.8, 22.7.9, 22.7.10, 1.8.1, 1.8.2, 1.8.3,

1.8.4, 1.8.5, 1.8.6, 1.8.7, 1.8.8, 1.8.9, 1.8.10, 2.8.1, 2.8.2, 2.8.3, 2.8.4, 2.8.5, 2.8.6, 2.8.7, 2.8.8, 2.8.9, 2.8.10, 3.8.1, 3.8.2, 3.8.3, 3.8.4, 3.8.5, 3.8.6, 3.8.7, 3.8.8, 3.8.9, 3.8.10, 4.8.1, 4.8.2, 4.8.3, 4.8.4, 4.8.5, 4.8.6, 4.8.7, 4.8.8, 4.8.9, 4.8.10, 5.8.1, 5.8.2, 5.8.3, 5.8.4, 5.8.5, 5.8.6, 5.8.7, 5.8.8, 5.8.9, 5.8.10, 6.8.1, 6.8.2, 6.8.3, 6.8.4, 6.8.5, 6.8.6, 6.8.7, 6.8.8, 6.8.9, 6.8.10, 7.8.1, 7.8.2, 7.8.3, 7.8.4, 7.8.5, 7.8.6, 7.8.7, 7.8.8, 7.8.9, 7.8.10, 8.8.1, 8.8.2, 8.8.3, 8.8.4, 8.8.5, 8.8.6, 8.8.7, 8.8.8, 8.8.9, 8.8.10, 9.8.1, 9.8.2, 9.8.3, 9.8.4, 9.8.5, 9.8.6, 9.8.7, 9.8.8, 9.8.9, 9.8.10, 10.8.1, 10.8.2, 10.8.3, 10.8.4, 10.8.5, 10.8.6, 10.8.7, 10.8.8, 10.8.9, 10.8.10, 11.8.1, 11.8.2, 11.8.3, 11.8.4, 11.8.5, 11.8.6, 11.8.7, 11.8.8, 11.8.9, 11.8.10, 12.8.1, 12.8.2, 12.8.3, 12.8.4, 12.8.5, 12.8.6, 12.8.7, 12.8.8, 12.8.9, 12.8.10, 13.8.1, 13.8.2, 13.8.3, 13.8.4, 13.8.5, 13.8.6, 13.8.7, 13.8.8, 13.8.9, 13.8.10, 14.8.1, 14.8.2, 14.8.3, 14.8.4, 14.8.5, 14.8.6, 14.8.7, 14.8.8, 14.8.9, 14.8.10, 15.8.1, 15.8.2, 15.8.3, 15.8.4, 15.8.5, 15.8.6, 15.8.7, 15.8.8, 15.8.9, 15.8.10, 16.8.1, 16.8.2, 16.8.3, 16.8.4, 16.8.5, 16.8.6, 16.8.7, 16.8.8, 16.8.9, 16.8.10, 17.8.1, 17.8.2, 17.8.3, 17.8.4, 17.8.5, 17.8.6, 17.8.7, 17.8.8, 17.8.9, 17.8.10, 18.8.1, 18.8.2, 18.8.3, 18.8.4, 18.8.5, 18.8.6, 18.8.7, 18.8.8, 18.8.9, 18.8.10, 19.8.1, 19.8.2, 19.8.3, 19.8.4, 19.8.5, 19.8.6, 19.8.7, 19.8.8, 19.8.9, 19.8.10, 20.8.1, 20.8.2, 20.8.3, 20.8.4, 20.8.5, 20.8.6, 20.8.7, 20.8.8, 20.8.9, 20.8.10, 21.8.1, 21.8.2, 21.8.3, 21.8.4, 21.8.5, 21.8.6, 21.8.7, 21.8.8, 21.8.9, 21.8.10, 22.8.1, 22.8.2, 22.8.3, 22.8.4, 22.8.5, 22.8.6, 22.8.7, 22.8.8, 22.8.9 and 22.8.10.

Identification of Active Precursors. It is desirable to select the amino acid residue or sequence of the invention compounds having one or more peptide bonds, such as formula VII compounds, based on the substrate specificity of esterases and/or carboxypeptidases expected to be found within cells where precursor hydrolysis is desired. To the extent that the specificity of these enzymes is unknown, one will screen a plurality of nucleotide analogs or esters until the desired substrate specificity is found. This will be apparent from assay either of the generation of free phosphonate or of antimicrobial activity. One selects compounds that are (i) not hydrolyzed or hydrolyzed comparatively slowly in the upper gut, (ii) gut and cell permeable and (iii) hydrolyzed in the cell cytoplasm and/or systemic circulation. Screens with cells from particular tissues are used to identify precursors that are released in organs susceptible to a target viral or microbial infection, e.g. in the case of liver, precursor drugs capable of hydrolysis in the liver. Other infections, e.g. CMV or HIV, are treated with a precursor that is hydrolyzed at substantially

the same rate and to substantially the same degree in all tissues, with no one tissue preferentially hydrolyzing the precursor nucleosides.

The assays used can be those known in the art including intestinal lumen stability, cell permeation, liver homogenate stability and plasma stability assays. These assays are used to determine the bioavailability characteristics of particular active precursors according to routinely used methods.

Therapeutic Indications. The hydrolysis products of the invention compounds have activity against viruses, malignant cells and/or parasitic protozoans. For example, 9-(3-hydroxy-2-phosphonylmethoxypropyl (HPMP) and (2-phosphonylmethoxy)ethyl (PME) analogs of purine (adenine (A), guanine (G), 2,6-diaminopurine (DAP), 2-monoaminopurine (MAP), hypoxanthine (Hx) and pyrimidine (cytosine (C), uracil (U), thymine (T) were evaluated for antiviral properties. (S)-HPMPA, (S)-cyclic HPMPA, (S)-HPMPC, (S)-HPMPG, (S)-HPMPDAP, PMEDAP, PMEG and PMEAP were active against herpes simplex virus, type 1 and 2 (HSV-1 and -2). (S)-HPMPA and (S)-cyclic HPMPA were active against varicella zoster virus (VZV). (S)-HPMPC was active against human cytomegalovirus (HCMV). (S)-HPMPA and (S)-cyclic HPMPA were shown to be active against adenovirus and vaccinia virus. PMEAP, PMEDAP, and PMEMAP are active against human immunodeficiency virus (HIV).

Acyclic nucleotide analogs having a common PME side chain covalently linked to a purine or pyrimidine heterocyclic base were prepared and tested for in vivo antiviral activity against retroviruses and herpes viruses. The adenine analog, PMEAP, was active in vitro against HIV and Rauscher murine leukemia virus (R-MuLV), and was more potent in vivo than 3'-azido-3'-deoxythymidine (AZT) in the treatment of R-MuLV in mice. PMEAP also had a significant antiviral effect in vivo against murine cytomegalovirus (MCMV), and in vitro activity against HCMV. The guanine analog, PMEG, was active in vitro against herpes viruses. In vivo, PMEG was >50-fold more potent than acyclovir against HSV 1 infection in mice.

(S)-HPMPA has potent and selective activity against a broad spectrum of DNA viruses, including HSV-1 and 2, VZV, thymidine kinase-deficient (TK<sup>-</sup>) mutants of herpes simplex virus, HCMV, phocid herpesvirus type 1

(seal herpesvirus, SeHV), simian herpesvirus type 1 (SHV-1), or pseudorabies virus or Aujeszky's disease virus), bovid herpesvirus type 1 (infectious bovine rhinotracheitis virus, BHV-1), equid herpesvirus type 1 (equine abortion virus, EHV-1), African swine fever (ASF) virus, vaccinia virus; and  
5 human adenoviruses, and retroviruses such as murine sarcoma virus (MSV). It is also reported that, in mice and rabbits in vivo, the compound is effective against both local and systemic infections with herpes simplex virus type 1, including herpetic keratitis caused by a TK<sup>-</sup> mutant which is resistant to the classical antiherpes drugs (DeClercq, E., et al, Antiviral Res (1987) 8:261-272;  
10 DeClercq, E., et al, Nature (1986) 323:464-467; Gil-Fernandez, C., et al, Antiviral Res (1987) 7:151-160; Baba, M., et al, Antimicrob Agents Chemother (1987) 31:337-339).

Phosphonylmethoxyalkylpurine analogs have also been evaluated for their antitumor activity in murine tumor models. HPMPPA, PMEA, and  
15 PMEG were found to be active against intraperitoneal P388 leukemia. PMEG was also found to be active against B16 melanoma.

As indicated above, the compounds of the invention are useful for treatment of microbial infections, for treatment of tumors or for other indications described below. Microbial infections include infection by viruses,  
20 parasites, yeasts and fungi. Exemplary viral infections that may be treated include infections mediated by DNA or RNA viruses including herpesviruses (CMV, HSV 1, HSV 2, EBV, varicella zoster virus, bovid herpesvirus type 1, equid herpesvirus type 1), papillomaviruses (HPV types 1-55), flaviviruses (including African swine fever virus and Japanese  
25 encephalitis virus), togaviruses (including Venezuelan equine encephalomyelitis virus), influenza viruses (types A-C), retroviruses (HIV 1, HIV 2, HTLV I, HTLV II, SIV, HBV, FeLV, FIV, MoMSV), adenoviruses (types 1-8), poxviruses (vaccinia virus), enteroviruses (polio virus type 1-3, hepatitis A virus), gastroenteritis viruses (Norwalk viruses, rotaviruses), hantaviruses  
30 (Hantaan virus), papovaviruses, rhinoviruses, parainfluenza virus types 1-4, rabies virus, and the like.

Some of the phosphonate compounds (such as PMEA) have a broad spectrum of antimicrobial activity and are thus unusual antiviral or antiparasitic agents. The activity of individual nucleotide analogs and  
35 nucleotide analog amidates is determined by routine assay of antiviral (or

other antimicrobial) activity using enzyme inhibition assays, tissue culture assays, animal model assays and/or other acceptable assays.

5 Nucleotide analogs (phosphonates such as HPMPC, PMEA, etc) are believed to exert their antimicrobial activity, at least in part, by a two step  
enzyme-mediated conversion to a diphosphate, followed by incorporation of  
the diphosphorylated nucleotide analog into nucleic acids. The incorporation  
of the diphosphates into nucleic acid is mediated by viral or other microbial  
DNA or RNA polymerases (bacterial, retroviral, etc). Thus, nucleotide  
analog (when diphosphorylated) are useful as chain terminators for  
10 dideoxynucleotide-type DNA sequencing protocols, provided that the nucleotide analog lacks a free hydroxyl group suitable for polymerase mediated chain elongation. These compounds will not have a hydroxyl group at R<sup>27</sup> in compounds of formulas IV and VI or are acyclic. Nucleotide analogs of formula XV, (HO)<sub>2</sub>P(O)-O-P(O)(OH)-O-(HO)P(O)-Z-B, can be  
15 prepared (Otvos, et al, Nucl Acids Res (1987) 15:1763-1777) and provided in a kit with other reagents (such as klenow polymerase or T4 polymerase, dNTPs, etc) needed for DNA sequencing. The invention nucleotide analogs and nucleotide analog amidates can also be (1) applied to tissue culture systems to eliminate or reduce viral spread or growth during the production of  
20 biopharmaceuticals or other products (such as proteins or vaccines), (2) used to eliminate or reduce viral spread or growth in clinical samples (such as blood), and (3) used to stop growth of tissue culture or bacterial cells (using toxic amounts of compound) without interfering with protein production.

Infections mediated by protozoan parasites can be treated using the  
25 compounds of the invention. Such infections can also be treated using the corresponding nucleotide analogs of the invention nucleotide analog amidates. The term protozoa is intended to include those members of the subphyla *Sarcomastigophora* and *Sporozoa* of the phylum *Protozoa*. More particularly, the term protozoa as used herein is intended to include those  
30 genera of parasitic protozoa which are important to man because they either cause disease in man or in his domestic animals. These genera are for the most part found classified in the superclass *Mastigophora* of the subphylum *Sarcomastigophora* and the class *Telosporea* of the subphylum *Sporozoa* in the classification according to Baker (1969). Illustrative genera of these  
35 parasitic protozoa include *Histomonas*, *Pneumocystis*, *Trypanosoma*, *Giardia*,

*Trichomonas*, *Eimeria*, *Isopora*, *Leishmania*, *Entamoeba*, *Toxoplasma* and *Plasmodium*. Parasitic protozoans include *Plasmodium falciparum*, *Plasmodium berghei*, *Plasmodium malariae*, *Plasmodium vivax*, *Leishmania braziliensis*, *Leishmania donovani*, *Trypanosoma cruzi*, *Trypanosoma brucei*,  
5 *Trypanosoma rhodesiense*, *Pneumocystis carinii*, *Entamoeba histolytica*, *Trichomonas vaginalis* and the like (de Vries, E., et al, Mol Biochem Parasitol (1991) 47:43-50). Nucleoside analog amidates of the invention and/or their corresponding nucleotide analogs can also be used to treat yeast or fungal infections caused by *Candida glabrata*, *Candida tropicalis*, *Candida albicans*,  
10 and other *Candida* species *Cryptococcus* species including *Cryptococcus neoformans*, *Blastomyces* species including *Blastomyces dermatidis*, *Torulopsis* species including *Torulopsis glabrata*, *Coccidioides* species including *Coccidioides immitis*, *Aspergillus* species and the like.

15       Pharmaceutical formulations. Compounds of the invention and their physiologically acceptable salts (hereafter collectively referred to as the active ingredients) may be administered by any route appropriate to the condition to be treated, suitable routes including oral, rectal, nasal, topical (including ocular, buccal and sublingual), vaginal and parenteral (including  
20 subcutaneous, intramuscular, intravenous, intradermal, intrathecal and epidural). The preferred route of administration may vary with for example the condition of the recipient.

While it is possible for the active ingredients to be administered alone it is preferably to present them as pharmaceutical formulations. The  
25 formulations, both for veterinary and for human use, of the present invention comprise at least one active ingredient, as above defined, together with one or more acceptable carriers therefor and optionally other therapeutic ingredients. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious  
30 to the recipient thereof.

The formulations include those suitable for oral, rectal, nasal, topical (including buccal and sublingual), vaginal or parenteral (including subcutaneous, intramuscular, intravenous, intradermal, intrathecal and epidural) administration. The formulations may conveniently be presented  
35 in unit dosage form and may be prepared by any of the methods well known

in the art of pharmacy. Such methods include the step of bringing into association the active ingredient with the carrier which constitutes one or more accessory ingredients. In general the formulations are prepared by uniformly and intimately bringing into association the active ingredient with  
5 liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient; as a powder or  
10 granules; as solution or a suspension in an aqueous liquid or a non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. The active ingredient may also be presented as a bolus, electuary or paste.

A tablet may be made by compression or moulding, optionally with  
15 one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, preservative, surface active or dispersing agent. Moulded tablets may be made by moulding in a suitable machine a mixture of the  
20 powdered compound moistened with an inert liquid diluent. The tablets may optionally be coated or scored and may be formulated so as to provide slow or controlled release of the active ingredient therein.

For infections of the eye or other external tissues e.g. mouth and skin, the formulations are preferably applied as a topical ointment or cream  
25 containing the active ingredient(s) in an amount of, for example, 0.075 to 20% w/w (including active ingredient(s) in a range between 0.1% and 20% in increments of 0.1% w/w such as 0.6% w/w, 0.7% w/w, etc), preferably 0.2 to 15% w/w and most preferably 0.5 to 10% w/w. When formulated in an ointment, the active ingredients may be employed with either a paraffinic or  
30 a water-miscible ointment base. Alternatively, the active ingredients may be formulated in a cream with an oil-in-water cream base.

If desired, the aqueous phase of the cream base may include, for example, at least 30% w/w of a polyhydric alcohol, i.e. an alcohol having two or more hydroxyl groups such as propylene glycol, butane 1,3-diol, mannitol,  
35 sorbitol, glycerol and polyethylene glycol (including PEG 400) and mixtures

thereof. The topical formulations may desirably include a compound which enhances absorption or penetration of the active ingredient through the skin or other affected areas. Examples of such dermal penetration enhancers include dimethyl sulphoxide and related analogs.

5       The oily phase of the emulsions of this invention may be constituted from known ingredients in a known manner. While the phase may comprise merely an emulsifier (otherwise known as an emulgent), it desirably comprises a mixture of at least one emulsifier with a fat or an oil or with both a fat and an oil. Preferably, a hydrophilic emulsifier is included  
10 together with a lipophilic emulsifier which acts as a stabilizer. It is also preferred to include both an oil and a fat. Together, the emulsifier(s) with or without stabilizer(s) make up the so-called emulsifying wax, and the wax together with the oil and fat make up the so-called emulsifying ointment base which forms the oily dispersed phase of the cream formulations.

15       Emulgents and emulsion stabilizers suitable for use in the formulation of the present invention include Tween<sup>®</sup> 60, Span<sup>®</sup> 80, cetostearyl alcohol, benzyl alcohol, myristyl alcohol, glyceryl mono-stearate and sodium lauryl sulfate.

20       The choice of suitable oils or fats for the formulation is based on achieving the desired cosmetic properties, since the solubility of the active compound in most oils likely to be used in pharmaceutical emulsion formulations is very low. Thus the cream should preferably be a non-greasy, non-staining and washable product with suitable consistency to avoid leakage from tubes or other containers. Straight or branched chain, mono- or dibasic  
25 alkyl esters such as di-isoadipate, isocetyl stearate, propylene glycol diester of coconut fatty acids, isopropyl myristate, decyl oleate, isopropyl palmitate, butyl stearate, 2-ethylhexyl palmitate or a blend of branched chain esters known as Crodamol CAP may be used, the last three being preferred esters. These may be used alone or in combination depending on the properties required.  
30 Alternatively, high melting point lipids such as white soft paraffin and/or liquid paraffin or other mineral oils can be used.

      Formulations suitable for topical administration to the eye also include eye drops wherein the active ingredient is dissolved or suspended in a suitable carrier, especially an aqueous solvent for the active ingredient. The  
35 active ingredient is preferably present in such formulations in a



concentration of 0.5 to 20%, advantageously 0.5 to 10% particularly about 1.5% w/w.

Formulations suitable for topical administration in the mouth include lozenges comprising the active ingredient in a flavored basis, usually sucrose  
5 and acacia or tragacanth; pastilles comprising the active ingredient in an inert basis such as gelatin and glycerin, or sucrose and acacia; and mouthwashes comprising the active ingredient in a suitable liquid carrier.

Formulations for rectal administration may be presented as a suppository with a suitable base comprising for example cocoa butter or a  
10 salicylate.

Formulations suitable for nasal administration wherein the carrier is a solid include a coarse powder having a particle size for example in the range 20 to 500 microns (including particle sizes in a range between 20 and 500 microns in increments of 5 microns such as 30 microns, 35 microns, etc),  
15 which is administered in the manner in which snuff is taken, i.e. by rapid inhalation through the nasal passage from a container of the powder held close up to the nose. Suitable formulations wherein the carrier is a liquid, for administration as for example a nasal spray or as nasal drops, include aqueous or oily solutions of the active ingredient. Formulations suitable for aerosol  
20 administration may be prepared according to conventional methods and may be delivered with other therapeutic agents such as pentamidine for treatment of pneumocystis pneumonia.

Formulations suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or spray formulations  
25 containing in addition to the active ingredient such carriers as are known in the art to be appropriate.

Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with  
30 the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier,  
35 for example water for injections, immediately prior to use. Extemporaneous

injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the kind previously described. Preferred unit dosage formulations are those containing a daily dose or unit daily sub-dose, as herein above recited, or an appropriate fraction thereof, of an active  
5 ingredient.

It should be understood that in addition to the ingredients particularly mentioned above the formulations of this invention may include other agents conventional in the art having regard to the type of formulation in question, for example those suitable for oral administration may include  
10 flavoring agents.

The present invention further provides veterinary compositions comprising at least one active ingredient as above defined together with a veterinary carrier therefor.

Veterinary carriers are materials useful for the purpose of  
15 administering the composition and may be solid, liquid or gaseous materials which are otherwise inert or acceptable in the veterinary art and are compatible with the active ingredient. These veterinary compositions may be administered orally, parenterally or by any other desired route.

Compounds of the invention can be used to provide controlled release  
20 pharmaceutical formulations containing as active ingredient one or more compounds of the invention ("controlled release formulations") in which the release of the active ingredient can be controlled and regulated to allow less frequency dosing or to improve the pharmacokinetic or toxicity profile of a given invention compound. Controlled release formulations adapted for  
25 oral administration in which discrete units comprising one or more compounds of the invention can be prepared according to conventional methods. Controlled release formulations may be employed for the treatment or prophylaxis of various microbial infections particularly human bacterial, human parasitic protozoan or human viral infections caused by  
30 microbial species including *Plasmodium*, *Pneumocystis*, herpesviruses (CMV, HSV 1, HSV 2, VZV, and the like), retroviruses, adenoviruses and the like. The controlled release formulations can be used to treat HIV infections and related conditions such as tuberculosis, malaria, pneumocystis pneumonia, CMV retinitis, AIDS, AIDS-related complex (ARC) and  
35 progressive generalized lymphadeopathy (PGL), and AIDS-related

neurological conditions such as multiple sclerosis, and tropical spastic paraparesis. Other human retroviral infections that may be treated with the controlled release formulations according to the invention include Human T-cell Lymphotropic virus (HTLV)-I and IV and HIV-2 infections.

- 5       The invention accordingly provides pharmaceutical formulations for use in the treatment or prophylaxis of the above-mentioned human or veterinary conditions and microbial infections.

10       Therapeutic Administration. For each of the above-indicated utilities and indications the amount required of an active ingredient (as above defined) will depend upon a number of factors including the severity of the condition to be treated and the identity of the recipient and will ultimately be at the discretion of the attendant physician or veterinarian. In general  
15       however, for each of these utilities and indications, a suitable, effective dose will be in the range 0.1 to 250 mg per kilogram bodyweight of recipient per dose (including active ingredient(s) in a range between 0.1 mg and 250 mg/Kg/dose in increments of 0.5 mg/Kg/dose such as 2.5 mg/Kg/dose, 3.0 mg/Kg/dose, 3.5 mg/Kg/dose, etc), preferably in the range 0.5 to 50 mg per  
20       kilogram body weight per dose and most preferably in the range 1 to 15 mg per kilogram body weight per dose; an optimum dose is about 3.0 mg per kilogram body weight per dose. (Unless otherwise indicated all weights of active ingredient are calculated as the parent compound of formula I: for salts thereof the figures would be increased proportionately). The desired dose is preferably presented as one dose or two sub-doses administered at appropriate  
25       intervals throughout a period of one to seven days. It is preferred to administer a dose once every 2, 3, 4, 5 or 6 days. The doses may be administered in unit dosage forms. The desired dose is may be presented as one, two, or three sub-doses administered at appropriate intervals throughout the one to seven day period. These sub-doses may be administered in unit  
30       dosage form, for example, containing 10 to 1000 mg, and or 100 to 500 mg of active ingredient per unit dosage form. The formulations should be desirably administered to achieve peak plasma concentrations of the active compound of from about 1 to about 100  $\mu$ M, preferably about 2 to 50  $\mu$ M, most preferably about 3 to about 30  $\mu$ M.

Compounds such as those of structures XXXI, XXXII and XXXIII (defined below) will generally (1) have a higher oral bioavailability than the corresponding uncyclized nucleotide analog (e.g., cHPMPC compared to HPMPC) and/or (2) will exhibit reduced toxicity when compared with the same  
5 dose of the corresponding uncyclized nucleotide analog, and/or (3) will have greater efficacy when compared with the same dose of the corresponding uncyclized nucleotide analog.

The compounds of the invention may be employed in combination with other therapeutic agents for the treatment or prophylaxis of the  
10 infections or conditions indicated above. Examples of such further therapeutic agents include agents that are effective for the treatment or prophylaxis of viral, parasitic or bacterial infections or associated conditions or for treatment of tumors or related conditions include 3'-azido-3'-  
15 deoxythymidine (zidovudine, AZT), 2'-deoxy-3'-thiacytidine (3TC), 2',3'-dideoxy-2',3'-didehydroadenosine (D4A), 2',3'-dideoxy-2',3'-didehydrothymidine (D4T), carbovir (carbocyclic 2',3'-dideoxy-2',3'-  
didehydroguanosine), 3'-azido-2',3'-dideoxyuridine, 5-fluorothymidine, (E)-5-(2-bromovinyl)-2'-deoxyuridine (BVDU), 2-chlorodeoxyadenosine, 2-  
20 deoxycoformycin, 5-fluorouracil, 5-fluorouridine, 5-fluoro-2'-deoxyuridine, 5-trifluoromethyl-2'-deoxyuridine, 6-azauridine, 5-fluoroorotic acid, methotrexate, triacetyluridine, 1-(2'-deoxy-2'-fluoro-1- $\beta$ -arabinosyl)-5-  
iodocytidine (FIAC), tetrahydro-imidazo(4,5,1-jk)-(1,4)-benzodiazepin-2(1H)-thione (TIBO), 2'-nor-cyclicGMP, 6-methoxypurine arabinoside (ara-M), 6-  
methoxypurine arabinoside 2'-O-valerate, cytosine arabinoside (ara-C), 2',3'-  
25 dideoxynucleosides such as 2',3'-dideoxycytidine (ddC), 2',3'-dideoxyadenosine (ddA) and 2',3'-dideoxyinosine (ddI), acyclic nucleosides such as acyclovir, penciclovir, famciclovir, ganciclovir, HPMPC, PMEA, PMEG, PMPA, PMPDAP, FPMMA, HPMPA, HPMPDAP, (2R, 5R)-9-  
[tetrahydro-5-(phosphonomethoxy)-2-furanyl]adenine, (2R, 5R)-1-[tetrahydro-  
30 5-(phosphonomethoxy)-2-furanyl]thymine, other antivirals including ribavirin (adenine arabinoside), 2-thio-6-azauridine, tubercidin, aurintricarboxylic acid, 3-deazaneoplanocin, neoplanocin, rimantidine, adamantine, and foscarnet (trisodium phosphonoformate), antibacterial agents including bactericidal fluoroquinolones (ciprofloxacin, pefloxacin and  
35 the like), aminoglycoside bactericidal antibiotics (streptomycin, gentamicin,

amicacin and the like)  $\beta$ -lactamase inhibitors (cephalosporins, penicillins and the like), other antibacterials including tetracycline, isoniazid, rifampin, cefoperazone, clathromycin and azithromycin, antiparasite or antifungal agents including pentamidine (1,5-bis(4'-aminophenoxy)pentane), 9-  
5 deazainosine, sulfamethoxazole, sulfadiazine, quinapyramine, quinine, fluconazole, ketoconazole, itraconazole, Amphotericin B, 5-fluorocytosine, clotrimazole, hexadecylphosphocholine and nystatin, renal excretion inhibitors such as probenecid, nucleoside transport inhibitors such as  
10 dipyridamole, dilazep and nitrobenzylthioinosine, immunomodulators such as FK506, cyclosporin A, thymosin  $\alpha$ -1, cytokines including TNF and TGF- $\beta$ , interferons including IFN- $\alpha$ , IFN- $\beta$  and IFN- $\gamma$ , interleukins including interleukin I, II, III, IV, V, VI, VII, VIII, X, XII, XIII macrophage/granulocyte colony stimulating factors including GM-CSF, G-CSF, M-CSF, cytokine  
15 antagonists including anti-TNF antibodies, anti-interleukin antibodies, soluble interleukin receptors, protein kinase C inhibitors and the like.

Immunogens and Antibodies. The compounds of this invention, or the biologically active substances produced from these compounds by hydrolysis in vivo, are used as immunogens to prepare antibodies capable of  
20 binding specifically to the compounds or their hydrolysis products. The immunogenic compositions therefore are useful as intermediates in the preparation of antibodies for use in diagnostic or quality control assays for the compounds or their hydrolysis products. The antibodies are useful for measuring the presence, absence or amounts of the compounds by any  
25 convenient homogenous or heterogenous procedure such as fluorescence polarization immunoassay, fluorescence immunoassay (using fluorescent labels such as fluorescein and the like), radioimmunoassay, enzyme immunoassay (using enzyme indicators such as alkaline phosphatase, horseradish peroxidase, glucose oxidase, urease and the like) and  
30 nephelometric inhibition assay by described methods (WO 92/22639, incorporated herein by reference). Such assays usually require a tracer (such as a fluorescent or radiolabeled labeled invention compound), an antibody and the sample to be analyzed containing the compound.

The hydrolysis products of interest are the phosphonates resulting  
35 from the hydrolysis of the amidate or ester bond(s) of the precursor

compounds of this invention, for example HPMPC, 6-aza-HPMPC, cyclic HPMPC, PMEA, PMEG, PMPDAP, PMPA, D4TMPI, D4AMPI, cyclic HPMPA, FPMMPA, PMEDAP, PMEMAP, 7-deaza-8-aza-FPMMPA, 7-deaza-8-aza-HPMPA, cyclic 7-deaza-8-aza-HPMPA, 7-deaza-8-aza-PMPA, 8-aza-FPMMPA, 8-aza-HPMPA, cyclic 8-aza-HPMPA, 8-aza-PMPA, PMPG, PMPMAP, 1-deaza-HPMPA, cyclic 1-deaza-HPMPA, 1-deaza-PMPA, 1-deaza-PMPG, 1-deaza-PMPMAP, 1-deaza-PMPDAP, 3-deaza-HPMPA, cyclic 3-deaza-HPMPA or 3-deaza-PMPA. Thus, the antibodies of this invention will be capable of binding to the precursors without binding to the hydrolysis products, will be capable of binding to the hydrolysis products without binding to the precursors, or will be capable of binding specifically to both. The antibodies will not cross-react with naturally-occurring nucleotides or nucleosides.

The immunogens of this invention contain the precursor or hydrolytic products in association with an immunogenic substance such as a protein or peptide. Immunogenic substances include adjuvants such as Freund's adjuvant, immunogenic proteins such as viral, bacterial, yeast, plant and animal polypeptides, in particular keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin or soybean trypsin inhibitor, and immunogenic polysaccharides. Typically, the precursor or a compound having the structure of a precursor hydrolytic product is covalently conjugated to an immunogenic polypeptide or polysaccharide by the use of a polyfunctional (ordinarily bifunctional) cross-linking agent. Methods for the manufacture of haptens are conventional *per se*, and any of the methods used heretofore for conjugating haptens to immunogenic polypeptides or the like are suitably employed here as well, taking into account the functional groups on the precursors or hydrolytic products which are available for cross-linking.

Typically the polypeptide is conjugated to a site on the heterocyclic base functionality of the compound or hydrolysis product rather than to a site on the alkyl or substituted-alkyl phosphonate moiety. In general, the site will be an amino group located on the purine or pyrimidine moiety of the nucleoside phosphonate, at the 5 position of pyrimidines (such as cytosine or uracil), at the 1 position of purines (such as adenosine or guanine) or, for compounds having a cyclic structure corresponding to a sugar or sugar analog and having a free hydroxyl group, through the hydroxyl group (usually at the

3' or 2' positions). Alternatively, the precursor compound is cross-linked through the phosphonate, typically by amidation or esterification of the phosphonate by the polypeptide itself or by a cross-linking functionality covalently bonded to the polypeptide. Thus, the groups L<sup>1</sup> or L<sup>2</sup> in structures (L<sup>1</sup>)(L<sup>2</sup>)-P(O)-Z-B can be immunogenic proteins (having more than 50 amino acid residues, usually less than 1000 residues) or peptides (about 5 to 50 amino acid residues).

The conjugates are prepared in conventional fashion. For example, N-hydroxysuccinimide, succinic anhydride or alkN=C=Nalk are useful in preparing the conjugates of this invention. The conjugates contain a precursor, its hydrolysis product, or both. Ordinarily, the conjugates will comprise the hydrolysis product, i.e., the biologically active drug. The conjugates are separated from starting materials and byproducts using chromatography or the like, and then are sterile filtered and vialled for storage.

Animals are typically immunized against the immunogenic conjugates or derivatives by combining 1 mg or 1 µg of conjugate (for rabbits or mice, respectively) with 3 volumes of Freund's complete adjuvant and injecting the solution intradermally at multiple sites. One month later the animals are boosted with 1/5 to 1/10 the original amount of conjugate in Freund's complete adjuvant (or other suitable adjuvant) by subcutaneous injection at multiple sites. 7 to 14 days later animals are bled and the serum is assayed for the desired antibody titer. Animals are boosted until the titer plateaus. Preferably, the animal is boosted with the conjugate in which the precursor or product is linked to a different protein, through a different cross-linking agent or both. Optionally, aggregating agents such as alum are used to enhance the immune response.

After immunization, monoclonal antibodies are prepared by recovering immune lymphoid cells (typically spleen cells or lymphocytes from lymph node tissue) from immunized animals and immortalizing the cells in conventional fashion, e.g., by fusion with myeloma cells or by Epstein-Barr virus transformation and screening for clones expressing the desired antibody. The hybridoma technique described originally by Kohler and Milstein, Eur. J. Immunol. (1976) 6:511 has been widely applied to

produce hybrid cell lines that secrete high levels of monoclonal antibodies against many specific antigens.

It is possible to fuse cells of one species with another. However, it is preferably that the source of the immunized antibody producing cells and the myeloma be from the same species.

The hybrid cell lines are maintained in culture in vitro. The cell lines of this invention are selected or maintained in a hypoxanthine-aminopterin thymidine (HAT) medium. However, the established hybridoma cell line can be maintained on a variety of nutritionally adequate media. The secreted antibody is recovered from culture by conventional methods such as precipitation, ion exchange chromatography, affinity chromatography, or the like. The antibodies described herein are also recovered from hybridoma cell cultures by conventional methods for purification of IgG or IgM as the case may be that heretofore have been used to purify immunoglobulins from pooled plasma, e.g., ethanol or polyethylene glycol precipitation procedures. The purified antibodies are sterile filtered, and optionally are conjugated to a detectable marker such as an enzyme or spin label for use in diagnostic assays of test samples.

The antibodies of this invention are obtained from any animal species, but ordinarily are murine or rat. Once a monoclonal antibody having the desired specificity and affinity is obtained, other conventional modifications of the antibodies are within the scope of this invention. For example, the complementarity determining regions of an animal antibody, together with as much of the framework domain as is needed, are substituted into an antibody of another animal species or class to produce a cross-class or cross-species chimeric antibody. Fragments or other amino acid sequence variants of monoclonal antibodies also are encompassed within the meaning of antibody as that term is used herein, for example, Fab, Fab' or (Fab')<sub>2</sub> fragments, single chain antibodies, bi or polyspecific antibodies, and the like.

The antibodies of this invention are from any suitable class or isotype, e.g. IgG, IgM, IgA, IgD or IgE. They may or may not participate in complement binding or ADCC.

Typically, hybridomas which are capable of binding to the immunogen are screened for the ability to bind to the hapten itself in typical test samples (plasma, serum and the like) with the requisite degree of affinity. The desired



affinity will depend upon the use intended for the antibody, but should be adequate to function in a conventional competitive-type ELISA or radioimmunoassays, or in conventional EMIT immunoassays.


5 The antibodies of this invention are used in such assays together with a labeled from of the precursor or its hydrolytic product. Alternatively, the antibody is labeled. Suitable labels are well-known and include radioisotopes, enzymes, stable free radicals, fluorophors, chemiluminescent moieties and other detectable groups heretofore employed to prepare covalent conjugates for use in assays. Methods for linking the labels to ligand amino groups, or  
10 amino acid side chains or termini of polypeptides, are known and are suitable for use herein. Other suitable linking methods will be apparent to the ordinary artisan.

The antibodies and labeled ligands herein optionally are assembled into kits for use in therapeutic drug monitoring or evaluation, or for process  
15 quality control, and used in the conventional manner.

Diagnostic applications. Novel compounds described herein are useful as intermediates in the preparation of detectable labels for oligonucleotide probes. The compounds are hydrolyzed to the diacid, diphosphorylated and then incorporated into an oligonucleotide by conventional enzymatic or  
20 chemical means. The incorporated heterocyclic base from the invention will generally be capable of participating in heterocyclic base pairing and thus will not interfere substantially with the binding of the oligonucleotide to its complementary sequence (E. DeClerq (1993) 3:85-96); should it interfere with  
25 oligonucleotide binding to its complementary sequence, the nucleotide analog is incorporated as the final 3' terminal residue, an innocuous position and a conventional site for oligonucleotide labeling. The nucleotide analog compound in the oligonucleotide is detected by any means, such as NMR, immune, fluorescence or radiolabel detection.

30

Bis amidate synthesis. Synthesis of bis-phosphoroamidate nucleotide analogs of Formula Id,



- where L<sup>1</sup> and L<sup>2</sup> are the same and are an amino acid, dipeptide, tripeptide or oligopeptide (4, 5 or 6 amino acid residues) are prepared by conversion of a nucleotide analog (such as PMEA, HPMPC, HPMPA, PMEG, FMPMA, PMPDAP, 9-[2,3-dideoxy-2,3-didehydro-4-phosphonmethoxy-β-D-erythrofuransyl]adenine (D4AMPI; reg no. 132178-53-1), 1-[2,3-dideoxy-2,3-didehydro-4-phosphonmethoxy-β-D-erythrofuransyl]thymine (D4TMPI; reg no. 132178-49-5) and the like) directly to the corresponding bis-phosphoroamidate compound. L<sup>1</sup> is a protein, an amino acid, dipeptide, tripeptide or oligopeptide (4 to 6 amino acid residues) which is esterified at free α-carboxyl group(s) by R<sup>4</sup>. Suitable R<sup>4</sup> groups include methyl, ethyl, propyl, isopropyl, butyl, t-butyl, phenyl, benzyl, N-ethylmorpholino, pivaloyloxymethyl and the like. The amino acids can comprise an aliphatic or aromatic side group (such as ala, phe, pro, leu, ile, met, trp and the like) or a dipeptide comprising amino acids having aliphatic or aromatic side groups (such as gly-gly, ala-ala, gly-ala, ala-gly, phe-gly, gly-phe, ala-phe, phe-ala, leu-ala, ala-leu and the like), or is a tripeptide comprising amino acids having aliphatic or aromatic side groups or is an oligopeptide comprising amino acids having aliphatic or aromatic side groups.
- The procedure is suitable for all of the nucleotide analogs described herein. The synthesis is accomplished by suspension of the nucleotide analog and approximately 2 equivalents of the L<sup>1</sup> species in a solvent such as dry pyridine or DMF (dimethylformamide) optionally containing a non-nucleophilic organic base such as triethylamine (about 3 to 10 equivalents).
- The dehydration step is accomplished by modification of a described reaction (Mukaiyama, T. et al, J Am Chem Soc (1972) 94:8528-8532) by adding a 1:1 mixture of triphenylphosphine (reg. no. 603-35-0; Aldrich) and 2,2'-dipyridyl disulfide (2 to 4 equivalents; reg. no. 2127-03-9; Aldrich) in pyridine to the nucleotide analog/amino acid mixture and (a) stirring at room temperature for about 4 to 16 hours or (b) heating to 60° C to 100° C (including any temperature in one degree C increments between 60° and 100° C such as 70°, 80° or 90° C) for about 4 to 16 hours. The resulting reaction mixture is then

concentrated and the final bis-amidate product is recovered and purified by conventional methods.

5 An alternative reaction suitable for synthesizing most amidate compounds is converting a nucleotide analog phosphonate to the corresponding chloridate by reaction with thionyl chloride in solvent (DMF) as described in EP 481 214. An amino acid, dipeptide or other molecule bearing a free amine is then reacted with the chloridate to yield the corresponding bis-amidate.


10 Synthesis of compounds of Formula Id having amino acids that contain amino, guanidino or carboxyl groups (such as lys, arg, his, asn, gln, lys-lys, arg-arg, lys-arg and the like) is accomplished by the same method, but using protected amine or carboxyl groups. After synthesis of the protected bis-amidate compound, the protecting groups are removed by conventional methods. Suitable protecting groups are well known and include acid labile  
15 groups such as p-tosyl, BOC (t-butoxycarbonyl) and FMOC (fluorene methoxycarbonyl) for protecting amine groups. Groups such as t-butyl, methyl, ethyl, benzyl and the like can be used to protect carboxyl groups. These groups can be removed under acid, base or hydrogenolysis conditions or can be removed with an esterase according to conventional methods.

20 Synthesis of compounds of Formula Id having amino acids such as tyr, cys, ser and thr is accomplished by optionally protecting hydroxyl or thiol groups using protecting groups known in the art. For example, the hydroxyl group of ser, thr or tyr can be protected using benzyl, ethyl and the like and the thiol group of cys can be protected using trityl, p-methylbenzyl and the  
25 like. The choice of a protecting group will depend on the stability of the bis-amidate toward conditions used to remove a particular protecting group. Appropriate protecting groups can be selected or determined by the skilled artisan using routine methods.

30 Dipeptide or tripeptide species can be selected on the basis of known transport properties and/or susceptibility to peptidases that can affect transport to intestinal mucosal or other cell types. Dipeptides and tripeptides lacking an  $\alpha$ -amino group are transport substrates for the peptide transporter found in brush border membrane of intestinal mucosal cells (Bai, J.P.F., Pharm Res (1992) 9:969-978). Transport competent peptides can thus be used  
35 to enhance bioavailability of bis amidate compounds. Di- or tripeptides

having one or more amino acids in the D configuration are also compatible with peptide transport and can be utilized in bis amidate compounds. Amino acids in the D configuration can be used to reduce the susceptibility of a di- or tripeptide to hydrolysis by proteases common to the brush border such as aminopeptidase N (EC 3.4.11.2). In addition, di- or tripeptides with amino acid residues can be selected on the basis of their relative resistance to hydrolysis by proteases found in the lumen of the intestine. For example, tripeptides or oligopeptides lacking asp and/or glu are poor substrates for aminopeptidase A (EC 3.4.11.7) and di- or tripeptides lacking amino acid residues on the N-terminal side of hydrophobic amino acids (leu, tyr, phe, val, trp) are poor substrates for endopeptidase 24.11 (EC 3.4.24.11) while peptides lacking a pro residue at the penultimate position at a free carboxyl terminus are poor substrates for carboxypeptidase P (EC 3.4.17). Similar considerations can also be applied to the selection of peptides that are either relatively resistant or relatively susceptible to hydrolysis by cytosolic, renal, hepatic, serum or other peptidases.

Synthesis of N-alkylamine amidates (where  $-NHR^{40}$  is linked to the phosphorus atom and  $R^{40}$  is  $C_{1-20}$  alkyl, including  $C_{4-16}$  alkyl) is accomplished essentially as described (Saito Chem. Pharm. Bull. (1991) 39:3207). Thus, compounds such as, for example, of structure  $(R^{40}HN)(L^1)P(O)-Z-B^2$  or



wherein  $B^2$  is B or  $B^1$ , are synthesized in this manner.

Amidate-ester synthesis. Synthesis of mixed amidate-ester nucleotide analog amidates of Formula Id where  $L^1$  is an amino acid ester and  $L^2$  is a group of the formula OR, SR or  $OR^{31}$  is accomplished by conversion of a nucleotide analog (such as PMEAs, HPMPCs, HPMPAs, PMEGs, FPMAs, PMPDAPs, D4AMPIs, D4TMPIs and the like) di- or bis-ester to a corresponding mixed ester-phosphoramidate compound. A bis ester is converted to a mono ester by treatment with a base such as ammonia to remove one ester group. The resulting mono ester is then converted to a mixed amidate-ester as described for synthesis of bis amidate compounds.

Bis ester synthesis. Bis esters of the formula  $(RO)_2P(O)-Z-B$  are generally synthesized as described in EP 481 214 or as described in Mukaiyama, T. et al, J Am Chem Soc (1972) 94:8528-8532. Dialkyl phosphonate esters are synthesized via conversion of a dichlorophosphonate (chloridate) such as  $(Cl)_2P(O)-Z-B$  (Quast, H. et al, Synthesis (1974) 7:489-490; Quast, H. et al, Synthesis (1974) 7:490; Moedritzer, K. et al, Synth Reac Inorg Met - Org Chem (1974) 5:417-27; Moedritzer, K., Chem Abs 82:86340; Stowell, M.H.B., et al Tet Lett (1990) 31:3261-3262) to a corresponding dialkylester (or dialkylamide) by reaction with alcohols (or amines). Monoalkylesters (or mono alkylamides) are obtained by hydrolysis of the disubstituted phosphonate in base (NaOH, KOH and the like). Disubstituted diacyloxyalkyl phosphonates are obtained by reaction of the unsubstituted phosphonate with a substituted chloromethyl ester  $(R-C(O)-O-CH(R)-Cl)$ . A corresponding monosubstituted acyloxyalkyl phosphonate is obtained by hydrolysis in acid or base.

For synthesis of Z substructures having a free hydroxyl group, such as  $(RO)_2P(O)-CH_2-O-CH(CH_2OH)-CH_2-$ , the hydroxyl is, in some cases, protected by a protecting group such as benzyl, acetyl, trityl, dimethoxytrityl and the like.

Bis esters having aryl, substituted aryl, alkyl-aryl or substituted alkyl-aryl (such as phenyl, alkoxyphenyl, benzyl, alkoxybenzyl) are also synthesized as described by reaction of  $(OH)_2P(O)-B-Z$  with thionyl chloride and a catalytic amount of DMF in a solvent such as acetonitrile. The resulting dichloridate,  $P(O)(Cl)_2-Z-B$  is then reacted with about 4, 5 or 6 equivalents of the sodium or potassium alkoxide or a sodium or potassium aryloxide obtained from reaction with sodium hydride or potassium hydride and the alcohol (such as phenol, benzyl alcohol and the like) in a solvent such as THF or acetonitrile at a reduced temperature (below about  $-70^\circ C$ , preferably about  $-76^\circ C$  to  $-78^\circ C$ ).

cHPMPC and the cyclic analogues of other cHPMPs are prepared by a number of methods from the free hydroxy phosphonic acid. These methods include treatment with DCC in DMF, reaction with Vilsmeier's reagent  $(ClCH=N(CH_3)_2Cl)$ , or methods of phosphate activation known per se. In one embodiment of this invention for the preparation of a cHPMP from the corresponding phosphonate nucleotide analog, the phosphonate is (a) treated with  $ClCH=N(CH_3)_2Cl$  to yield the phosphonylchloridate and (b) optionally

the phosphonylchloride is reacted with a nucleophile (preferably at low temperature, e.g. lower than about  $-20^{\circ}\text{C}$ ) such as an alcohol or amine to produce one of the intermediates described above. In a further step the product of steps (a) or (b) are subject to hydrolysis or protonolysis (typically acid protonolysis) respectively to yield the cHSNA (treatment of the product of step (a)) or its intermediate (treatment of the product of step (b)).


Vilsmeier's reagent is advantageously produced *in situ* by combining  $\text{SOCl}_2$ ,  $\text{PCl}_5$ ,  $\text{POCl}_3$ ,  $\text{COCl}_2$  or the like with DMF. Advantageously, the product of step (a) is not purified or separated from the reaction mixture before being reacted with the nucleophile, a distinct economic advantage for this synthetic route.

The compounds of structure (Ia) and (Va) are readily made from their uncyclized counterparts by the same methods, e.g. treatment with DCC in DMF.


Substituted and unsubstituted alkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl and other  $\text{L}^1$  esters and amidates of cHPMPs typically are made by reacting the appropriate HPMP compound with  $\text{SOCl}_2/\text{DMF}$  to yield the activated phosphonylchloride (see Scheme 1), followed by treatment with the corresponding nucleophile (e.g. alkoxide, phenolate, amine, etc.) to yield the protected intermediate formamidine which is subsequently hydrolyzed to the target compound. Alternatively, esters can also be prepared as depicted in Scheme 2. The N-,O- protected intermediate phosphonate diester is obtained from the three building blocks by known methods. The N- and O- protecting groups are subsequently removed followed by treatment of the phosphonate diester **3** with NaH leading to cyclization yielding target compound **4**. A third method for the synthesis of cHSNA esters entails alkylation of the cHSNA using common alkylating agents  $\text{D}^1\text{L}$  (where L is a leaving group) such as alkyl halides, tosylates, diazoalkanes and the like (see Scheme 3). This method is particularly useful for preparing acyloxyalkyl esters by treatment of the cHSNA with the corresponding acyloxyalkylhalide. In an exemplary method for the preparation of acyloxyalkyl esters of cHPMPs, as shown in more detail in Example 12, DCC and  $\text{R}^{45}\text{C}(\text{O})\text{OCH}_2\text{Cl}$  are reacted with the cyclic compound; but in contradistinction with prior methods the stoichiometric proportion of DCC:  $\text{R}^{45}\text{C}(\text{O})\text{OCH}_2\text{Cl}$ , cyclic HPMP is 1-2:1-2:1. Use of such low proportions of reactants lessens side reactions with any exocyclic amino group of B and thereby greatly improves yields.  $\text{R}^{45}$  is H or is

C<sub>3</sub>-C<sub>12</sub> alkyl which is unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N and halogen, C<sub>3</sub>-C<sub>6</sub> aryl which is unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N and halogen or C<sub>3</sub>-C<sub>9</sub> aryl-alkyl which is unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N and halogen.


Each of the following schemes exemplify HPMPC as the nucleotide analog. However, any B is employed in place of cytosine, provided that any exocyclic oxo or amino groups are protected as required. Also, step 3 of scheme 1 will be omitted when B contains no exocyclic amine.








Scheme 2



Scheme 3


A third method for the synthesis of cyclic HPMP esters entails alkylation of the cyclic HPMP ester as shown in Scheme 3 using common alkylating agents  $R^{35}Lv$  (where Lv is a leaving group) such as alkyl halides, tosylates, diazoalkanes and the like. This method is particularly useful for preparing acyloxyalkyl esters by treatment of the cyclic HPMP (cHPMP) with the corresponding acyloxyalkylhalide.

Compounds where Z is of structure V and  $R^{25}$  and  $R^{29}$  is oxygen are synthesized by addition-elimination reaction using a compound of structure



50, previously described for B = adenine (EP 398 231) with iodine (about 2 equivalents) in organic solvent (such as acetonitrile or methylene chloride) at about 15-24° C and a compound having the structure  $(R^{35}O)_2P(O)-CH_2-OH$ , wherein  $R^{35}$  is R or  $R^{31}$  (defined below), to yield the 3-


iodophosphonate diester of structure 51,




, which is then eliminated to yield the corresponding structure V compound by reaction with about 5 equivalents of a base such as sodium methoxide or DBU in anhydrous organic solvent such as methanol or tetrahydrofuran at room temperature for about 2-12 hours.

The following schemes show synthesis of intermediates having fluorine or iodine at the 3' position that are converted to structure V compounds by

- elimination with a base. N-Fluorodibenzenesulfonamide is available commercially (Aldrich). Structure V compounds where B and the phosphonate ester substituent at the 4' position are either both up or down (i.e., substituents at the 1' and 4' positions are cis with respect to each other) are obtained by using the corresponding structure 50 reactant as follows




- 10 When the reaction with iodine is conducted at high temperature (about 50-80° C, usually about 60-70° C), a scalemic intermediate results as follows




- 15 The intermediate is then converted to the corresponding structure V compound and the various cis and trans isomers can be separated using standard methods such as HPLC, RPLC or crystallization.

- Exemplary esters are of the formula,  $(R^{31}O)_2P(O)-Z-B$ ,  $(RO)(R^{31}O)P(O)-Z-B$  or  $(RO)_2P(O)-Z-B$ , wherein  $R^{31}$  is independently 2,3-dihydro-6-hydroxyindene, sesamol, catechol monoester,  $-CH_2-C(O)-N(R^7)_2$  wherein each  $R^7$  is the same or different,  $-CH_2-S(O)(R^7)$ ,  $-CH_2-S(O)_2(R^7)$ ,  $-CH_2-$  CH(OC(O)CH<sub>2</sub>R<sup>7</sup>)-CH<sub>2</sub>(OC(O)CH<sub>2</sub>R<sup>7</sup>), cholesteryl, a 5 or 6 carbon monosaccharide, disaccharide or oligosaccharide (3 to 9 monosaccharide residues), enolpyruvate (HOOC-C(=CH<sub>2</sub>)O), glycerol,  $\alpha$ -D- $\beta$ -diglycerides (wherein the fatty acids composing glyceride lipids generally are naturally occurring saturated or unsaturated C<sub>6-26</sub>, C<sub>6-18</sub> or C<sub>6-10</sub> fatty acids such as linoleic, lauric, myristic, palmitic, stearic, oleic, palmitoleic, linolenic and the like fatty acids), trimethoxybenzyl, triethoxybenzyl, 2-alkyl pyridinyl (C<sub>1-4</sub> alkyl),




C<sub>3</sub>-C<sub>6</sub> aryl (including phenyl, 2- and 3-pyrrolyl, 2- and 3-thienyl, 2- and 4-imidazolyl, 2-, 4- and 5-oxazolyl, 3- and 4-isoxazolyl, 2-, 4- and 5-thiazolyl, 3-, 4- and 5-isothiazolyl, 3- and 4-pyrazolyl, 2-, 3- and 4-pyridinyl and 2-, 4- and 5-pyrimidinyl) substituted by 3, 4 or 5 halogen atoms or 1 or 2 atoms or groups selected from halogen, C<sub>1</sub>-C<sub>12</sub> alkoxy (including methoxy, ethoxy, 2,3-, 2,4-, 2,5-, 2,6-, 3,4- and 3,5-dimethoxy and 2,3-, 2,4-, 2,5-, 2,6-, 3,4- and 3,5-diethoxy substituted phenyl), cyano, nitro, OH, C<sub>1</sub>-C<sub>12</sub> haloalkyl (1 to 6 halogen atoms), C<sub>1</sub>-C<sub>12</sub> alkyl (including methyl and ethyl), C<sub>2</sub>-C<sub>12</sub> alkenyl or C<sub>2</sub>-C<sub>12</sub> alkynyl; or  
 5 R<sup>31</sup> is C<sub>1</sub>-C<sub>4</sub> alkylene-C<sub>3</sub>-C<sub>6</sub> aryl (including benzyl, -CH<sub>2</sub>-pyrrolyl, -CH<sub>2</sub>-thienyl, -CH<sub>2</sub>-imidazolyl, -CH<sub>2</sub>-oxazolyl, -CH<sub>2</sub>-isoxazolyl, -CH<sub>2</sub>-thiazolyl, -CH<sub>2</sub>-isothiazolyl, -CH<sub>2</sub>-pyrazolyl, -CH<sub>2</sub>-pyridinyl and -CH<sub>2</sub>-pyrimidinyl) substituted in the aryl moiety by 3 to 5 halogen atoms or 1 to 2 atoms or groups selected from halogen, C<sub>1</sub>-C<sub>12</sub> alkoxy (including methoxy and ethoxy),  
 10 cyano, nitro, OH, C<sub>1</sub>-C<sub>12</sub> haloalkyl (1 to 6 halogen atoms; including -CH<sub>2</sub>-CCl<sub>3</sub>), C<sub>1</sub>-C<sub>12</sub> alkyl (including methyl and ethyl), C<sub>2</sub>-C<sub>12</sub> alkenyl or C<sub>2</sub>-C<sub>12</sub> alkynyl. Methods for linking cholesteryl, saccharide and other moieties to reactive groups have been described (Hadfield Adv. Pharmacol. Chemother. (1984) 20:21; Gouyette Tet. Lett. (1989) 30:6019; Ksander J. Med. Chem. (1994) 37:1823).  
 20


The compounds are used as intermediates in the synthesis of mixed amidate-ester nucleotide analog amidates, or in some cases, as drugs *per se*. Additional exemplary ester compounds have the formulas (R<sup>31</sup>O)<sub>2</sub>P(O)-Z<sup>1</sup>-B or (RO)(R<sup>31</sup>O)P(O)-Z<sup>1</sup>-B, where Z<sup>1</sup> is defined to mean the substructure in the  
 25 following representative structures; (R<sup>31</sup>O)<sub>2</sub>-P(O)-CH<sub>2</sub>-O-CH<sub>2</sub>-CH<sub>2</sub>-B, (R<sup>31</sup>O)<sub>2</sub>-P(O)-CH<sub>2</sub>-O-C#H(CH<sub>2</sub>OH)-CH<sub>2</sub>-B, (R<sup>31</sup>O)<sub>2</sub>-P(O)-CH<sub>2</sub>-O-C#H(CH<sub>3</sub>)-CH<sub>2</sub>-B, (R<sup>31</sup>O)<sub>2</sub>-P(O)-CH<sub>2</sub>-O-C#H(CH<sub>2</sub>F)-CH<sub>2</sub>-B, (R<sup>31</sup>O)<sub>2</sub>-P(O)-CH<sub>2</sub>-O-C#H(CH=CH<sub>2</sub>)-CH<sub>2</sub>-B, (R<sup>31</sup>O)<sub>2</sub>-P(O)-CH<sub>2</sub>-O-C#H(CH<sub>2</sub>N<sub>3</sub>)-CH<sub>2</sub>-B,



where C<sup>#</sup>, R<sup>25</sup> - R<sup>29</sup>, R<sup>31</sup> and B have the meanings previously defined with the proviso that PMEAs bis(4-nitrobenzyl ester) and PMEAs bis(4-trifluoromethyl ester) are excluded and for structure XXIX, R<sup>29</sup> and R<sup>25</sup> are both O. Additional ester and nucleotide compounds are of the formula



where substituents linked to the carbon atom designated # are in the *R*, *S* or *RS* configuration and R<sup>31</sup> and B are as previously defined. Nucleotides and esters of the formulas



wherein #, B, R<sup>25</sup>, R<sup>26</sup>, R<sup>27</sup>, R<sup>28</sup>, R<sup>29</sup>, R<sup>31</sup>, R<sup>33</sup> and R<sup>34</sup> are as defined, R<sup>35</sup> is defined as R or R<sup>31</sup> and for structure XXX, when R<sup>29</sup> is CH<sub>2</sub> or O and R<sup>25</sup> is CH<sub>2</sub> or O, R<sup>35</sup> is not H or C<sub>1</sub>-C<sub>6</sub> alkyl, are new. Compounds having R<sup>35</sup>

include species where both  $R^{35}$  are both H and their salts including pharmaceutically acceptable salts.

Exemplary  $R^{31}$  include 2-, 3- and 4-alkoxyphenyl ( $C_1$ - $C_{12}$  alkyl including 2-, 3- and 4-methoxyphenyl and 2-, 3- and 4-ethoxyphenyl), 2-, 3- and 4-carboethoxyphenyl, 2- and 3-carboethoxy-4-hydroxyphenyl, 2- and 3-ethoxy-4-hydroxyphenyl, 2- and 3-ethoxy-5-hydroxyphenyl, 2- and 3-ethoxy-6-hydroxyphenyl, 2-, 3- and 4-O-acetylphenyl, 2-, 3- and 4-dimethylaminophenyl, 2-, 3- and 4-methylmercaptophenyl, 2-, 3- and 4-halophenyl (including 2-, 3- and 4-fluorophenyl and 2-, 3- and 4-chlorophenyl), 2,3-, 2,4-, 2,5-, 2,6-, 3,4- and 3,5-dimethylphenyl, 2,3-, 2,4-, 2,5-, 2,6-, 3,4- and 3,5-biscarboxyethylphenyl, 2,3-, 2,4-, 2,5-, 2,6-, 3,4- and 3,5-dimethoxyphenyl, 2,3-, 2,4-, 2,5-, 2,6-, 3,4- and 3,5-dihalophenyl (including 2,4-difluorophenyl and 3,5-difluorophenyl), 2-, 3- and 4-haloalkylphenyl (1 to 5 halogen atoms,  $C_1$ - $C_{12}$  alkyl including 4-trifluoromethylphenyl), 2-, 3- and 4-cyanophenyl, and 2-, 3- and 4-nitrophenyl, 2-, 3- and 4-haloalkylbenzyl (1 to 5 halogen atoms,  $C_1$ - $C_{12}$  alkyl including 4-trifluoromethylbenzyl),  $\alpha$ -D-galactose,  $\alpha$ -D-glucose,  $\alpha$ -D-fructose. The bis esters of formula  $(OR^{31})(OR^{31})P(O)-Z-B$  and  $(OR)(OR^{31})P(O)-Z-B$  are novel and are useful as intermediates in the synthesis of the mixed amidate-ester nucleotide analog amidates of the invention. These compounds can also be used directly as antimicrobial agents per se. Table 5 lists a group of exemplary bis esters of compounds having the structure  $(OR^{35})_2P(O)-Z-B$  which includes novel compounds of structure  $(OR^{31})_2P(O)-Z-B$ .

TABLE 5

	$OR^{35*}$	$-P(O)-Z-B^{**}$
30	1 $-O-C_6H_4F$	1 $-P(O)-CH_2-O-CH_2-CH_2-B$
	2 $-O-C_6H_3F_2$	2 $-P(O)-CH_2-O-C^{\#}H(CH_2-OR^4)-CH_2-B$
	3 $-O-C_6H_4-OCH_3$	3 $-P(O)-CH_2-O-C^{\#}H(CH_3)-CH_2-B$
	4 $-O-C_6H_3-(OCH_3)_2$	4 $-P(O)-CH_2-O-C^{\#}H(CH_2F)-CH_2-B$
	5 $-O-C_6H_4-OC_2H_5$	5 $-P(O)-CH_2-O-C^{\#}H(CH=CH_2)-CH_2-B$
35	6 $-O-C_6H_3-(OC_2H_5)_2$	6 $-P(O)-CH_2-O-C^{\#}H(CH_2N_3)-CH_2-B$

- |    |          |  |   |    |
|----|----------|--|---|----|
|    | 7        | -O-CH <sub>2</sub> -C <sub>6</sub> H <sub>4</sub> F                                    | 7 | ** |
|    | 8        | -O-C <sub>6</sub> H <sub>4</sub> -(C(O)-O-C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub> | 8 | ** |
|    | 9        | -O-C <sub>6</sub> H <sub>4</sub> -C(O)-O-C <sub>2</sub> H <sub>5</sub>                 |   |    |
|    | 10       | -O-C <sub>6</sub> H <sub>3</sub> -(O-C(O)-CH <sub>3</sub> ) <sub>2</sub>               |   |    |
| 5  | 11       | -O-C <sub>6</sub> H <sub>3</sub> -C(O)-O-C <sub>3</sub> H <sub>7</sub>                 |   |    |
|    | 12       | -O-CH <sub>2</sub> -C <sub>6</sub> H <sub>4</sub> -O-CO-CH <sub>3</sub>                |   |    |
|    | 13       | -O-C <sub>5</sub> H <sub>4</sub> N   |   |    |
|    | 14       | -O-C <sub>6</sub> H <sub>3</sub> -(OC <sub>2</sub> H <sub>5</sub> )(OH)                |   |    |
|    | 15       | -O-C <sub>6</sub> H <sub>5</sub>   |   |    |
| 10 | 16       | -O-CH <sub>2</sub> -O-C(O)-C(CH <sub>3</sub> ) <sub>3</sub>                            |   |    |
|    | <b>B</b> |  |   |    |
|    | 1        | adenin-9-yl  |   |    |
|    | 2        | guanin-9-yl  |   |    |
|    | 3        | cytosin-1-yl   |   |    |
| 15 | 4        | 2, 6-diaminopurin-9-yl   |   |    |
|    | 5        | 2-aminopurin-9-yl  |   |    |
|    | 6        | thymidin-1-yl  |   |    |
|    | 7        | 5-fluorocytosin-1-yl   |   |    |

20 \* Monosubstituted phenyl and benzyl compounds (i.e., R<sup>35</sup> numbers 1, 3, 5, etc) include 2-, 3- and 4-substituted compounds and disubstituted phenyl compounds (i.e., R<sup>35</sup> numbers 2, 4, 6, etc) include 2,3-, 2,4-, 2,5-, 2,6-, 3,4- and 3,5-substituted compounds.

25 \*\* The structure =P(O)- indicates that two bonds are occupied by OR<sup>35</sup>; Z structure 7 is of formula IV where R<sup>25</sup> and R<sup>29</sup> are O, R<sup>26</sup> is S, R<sup>27</sup> is absent and R<sup>28</sup> is H and includes the (+) and (-) enantiomers; structure 8 is of formula V where R<sup>25</sup> and R<sup>29</sup> are O.

30 Compounds listed in Table 5 are designated herein by numbers assigned to (OR<sup>35</sup>)<sub>2</sub> (where each R<sup>35</sup> is the same), Z and B according to the following convention, R<sup>35</sup>.Z.B. Exemplary compounds include 1.1.1, 2.1.1, 3.1.1, 4.1.1, 5.1.1, 6.1.1, 7.1.1, 8.1.1, 9.1.1, 10.1.1, 11.1.1, 12.1.1, 13.1.1, 14.1.1, 15.1.1, 16.1.1, 1.2.1, 2.2.1, 3.2.1, 4.2.1, 5.2.1, 6.2.1, 7.2.1, 8.2.1, 9.2.1, 10.2.1, 11.2.1, 12.2.1, 13.2.1, 14.2.1, 15.2.1, 16.2.1, 1.3.1, 2.3.1, 3.3.1, 4.3.1, 5.3.1, 6.3.1, 7.3.1, 8.3.1, 9.3.1, 35 10.3.1, 11.3.1, 12.3.1, 13.3.1, 14.3.1, 15.3.1, 16.3.1, 1.4.1, 2.4.1, 3.4.1, 4.4.1, 5.4.1, 6.4.1,

7.4.1, 8.4.1, 9.4.1, 10.4.1, 11.4.1, 12.4.1, 13.4.1, 14.4.1, 15.4.1, 16.4.1, 1.5.1, 2.5.1, 3.5.1,  
 4.5.1, 5.5.1, 6.5.1, 7.5.1, 8.5.1, 9.5.1, 10.5.1, 11.5.1, 12.5.1, 13.5.1, 14.5.1, 15.5.1, 16.5.1,  
 1.6.1, 2.6.1, 3.6.1, 4.6.1, 5.6.1, 6.6.1, 7.6.1, 8.6.1, 9.6.1, 10.6.1, 11.6.1, 12.6.1, 13.6.1,  
 14.6.1, 15.6.1, 16.6.1, 1.7.1, 2.7.1, 3.7.1, 4.7.1, 5.7.1, 6.7.1, 7.7.1, 8.7.1, 9.7.1, 10.7.1,  
 5 11.7.1, 12.7.1, 13.7.1, 14.7.1, 15.7.1, 16.7.1, 1.8.1, 2.8.1, 3.8.1, 4.8.1, 5.8.1, 6.8.1, 7.8.1,  
 8.8.1, 9.8.1, 10.8.1, 11.8.1, 12.8.1, 13.8.1, 14.8.1, 15.8.1, 16.8.1, 1.1.2, 2.1.2, 3.1.2, 4.1.2,  
 5.1.2, 6.1.2, 7.1.2, 8.1.2, 9.1.2, 10.1.2, 11.1.2, 12.1.2, 13.1.2, 14.1.2, 15.1.2, 16.1.2, 1.2.2,  
 2.2.2, 3.2.2, 4.2.2, 5.2.2, 6.2.2, 7.2.2, 8.2.2, 9.2.2, 10.2.2, 11.2.2, 12.2.2, 13.2.2, 14.2.2,  
 15.2.2, 16.2.2, 1.3.2, 2.3.2, 3.3.2, 4.3.2, 5.3.2, 6.3.2, 7.3.2, 8.3.2, 9.3.2, 10.3.2, 11.3.2,  
 10 12.3.2, 13.3.2, 14.3.2, 15.3.2, 16.3.2, 1.4.2, 2.4.2, 3.4.2, 4.4.2, 5.4.2, 6.4.2, 7.4.2, 8.4.2,  
 9.4.2, 10.4.2, 11.4.2, 12.4.2, 13.4.2, 14.4.2, 15.4.2, 16.4.2, 1.5.2, 2.5.2, 3.5.2, 4.5.2, 5.5.2,  
 6.5.2, 7.5.2, 8.5.2, 9.5.2, 10.5.2, 11.5.2, 12.5.2, 13.5.2, 14.5.2, 15.5.2, 16.5.2, 1.6.2, 2.6.2,  
 3.6.2, 4.6.2, 5.6.2, 6.6.2, 7.6.2, 8.6.2, 9.6.2, 10.6.2, 11.6.2, 12.6.2, 13.6.2, 14.6.2, 15.6.2,  
 16.6.2, 1.7.2, 2.7.2, 3.7.2, 4.7.2, 5.7.2, 6.7.2, 7.7.2, 8.7.2, 9.7.2, 10.7.2, 11.7.2, 12.7.2,  
 15 13.7.2, 14.7.2, 15.7.2, 16.7.2, 1.8.2, 2.8.2, 3.8.2, 4.8.2, 5.8.2, 6.8.2, 7.8.2, 8.8.2, 9.8.2,  
 10.8.2, 11.8.2, 12.8.2, 13.8.2, 14.8.2, 15.8.2, 16.8.2, 1.1.3, 2.1.3, 3.1.3, 4.1.3, 5.1.3, 6.1.3,  
 7.1.3, 8.1.3, 9.1.3, 10.1.3, 11.1.3, 12.1.3, 13.1.3, 14.1.3, 15.1.3, 16.1.3, 1.2.3, 2.2.3, 3.2.3,  
 4.2.3, 5.2.3, 6.2.3, 7.2.3, 8.2.3, 9.2.3, 10.2.3, 11.2.3, 12.2.3, 13.2.3, 14.2.3, 15.2.3, 16.2.3,  
 1.3.3, 2.3.3, 3.3.3, 4.3.3, 5.3.3, 6.3.3, 7.3.3, 8.3.3, 9.3.3, 10.3.3, 11.3.3, 12.3.3, 13.3.3,  
 20 14.3.3, 15.3.3, 16.3.3, 1.4.3, 2.4.3, 3.4.3, 4.4.3, 5.4.3, 6.4.3, 7.4.3, 8.4.3, 9.4.3, 10.4.3,  
 11.4.3, 12.4.3, 13.4.3, 14.4.3, 15.4.3, 16.4.3, 1.5.3, 2.5.3, 3.5.3, 4.5.3, 5.5.3, 6.5.3, 7.5.3,  
 8.5.3, 9.5.3, 10.5.3, 11.5.3, 12.5.3, 13.5.3, 14.5.3, 15.5.3, 16.5.3, 1.6.3, 2.6.3, 3.6.3, 4.6.3,  
 5.6.3, 6.6.3, 7.6.3, 8.6.3, 9.6.3, 10.6.3, 11.6.3, 12.6.3, 13.6.3, 14.6.3, 15.6.3, 16.6.3, 1.7.3,  
 2.7.3, 3.7.3, 4.7.3, 5.7.3, 6.7.3, 7.7.3, 8.7.3, 9.7.3, 10.7.3, 11.7.3, 12.7.3, 13.7.3, 14.7.3,  
 25 15.7.3, 16.7.3, 1.8.3, 2.8.3, 3.8.3, 4.8.3, 5.8.3, 6.8.3, 7.8.3, 8.8.3, 9.8.3, 10.8.3, 11.8.3,  
 12.8.3, 13.8.3, 14.8.3, 15.8.3, 16.8.3, 1.1.4, 2.1.4, 3.1.4, 4.1.4, 5.1.4, 6.1.4, 7.1.4, 8.1.4,  
 9.1.4, 10.1.4, 11.1.4, 12.1.4, 13.1.4, 14.1.4, 15.1.4, 16.1.4, 1.2.4, 2.2.4, 3.2.4, 4.2.4, 5.2.4,  
 6.2.4, 7.2.4, 8.2.4, 9.2.4, 10.2.4, 11.2.4, 12.2.4, 13.2.4, 14.2.4, 15.2.4, 16.2.4, 1.3.4, 2.3.4,  
 3.3.4, 4.3.4, 5.3.4, 6.3.4, 7.3.4, 8.3.4, 9.3.4, 10.3.4, 11.3.4, 12.3.4, 13.3.4, 14.3.4, 15.3.4,  
 30 16.3.4, 1.4.4, 2.4.4, 3.4.4, 4.4.4, 5.4.4, 6.4.4, 7.4.4, 8.4.4, 9.4.4, 10.4.4, 11.4.4, 12.4.4,  
 13.4.4, 14.4.4, 15.4.4, 16.4.4, 1.5.4, 2.5.4, 3.5.4, 4.5.4, 5.5.4, 6.5.4, 7.5.4, 8.5.4, 9.5.4,  
 10.5.4, 11.5.4, 12.5.4, 13.5.4, 14.5.4, 15.5.4, 16.5.4, 1.6.4, 2.6.4, 3.6.4, 4.6.4, 5.6.4, 6.6.4,  
 7.6.4, 8.6.4, 9.6.4, 10.6.4, 11.6.4, 12.6.4, 13.6.4, 14.6.4, 15.6.4, 16.6.4, 1.7.4, 2.7.4, 3.7.4,  
 4.7.4, 5.7.4, 6.7.4, 7.7.4, 8.7.4, 9.7.4, 10.7.4, 11.7.4, 12.7.4, 13.7.4, 14.7.4, 15.7.4, 16.7.4,  
 35 1.8.4, 2.8.4, 3.8.4, 4.8.4, 5.8.4, 6.8.4, 7.8.4, 8.8.4, 9.8.4, 10.8.4, 11.8.4, 12.8.4, 13.8.4,



14.8.4, 15.8.4, 16.8.4, 1.1.5, 2.1.5, 3.1.5, 4.1.5, 5.1.5, 6.1.5, 7.1.5, 8.1.5, 9.1.5, 10.1.5, 11.1.5, 12.1.5, 13.1.5, 14.1.5, 15.1.5, 16.1.5, 1.2.5, 2.2.5, 3.2.5, 4.2.5, 5.2.5, 6.2.5, 7.2.5, 8.2.5, 9.2.5, 10.2.5, 11.2.5, 12.2.5, 13.2.5, 14.2.5, 15.2.5, 16.2.5, 1.3.5, 2.3.5, 3.3.5, 4.3.5, 5.3.5, 6.3.5, 7.3.5, 8.3.5, 9.3.5, 10.3.5, 11.3.5, 12.3.5, 13.3.5, 14.3.5, 15.3.5, 16.3.5, 1.4.5, 2.4.5, 3.4.5, 4.4.5, 5.4.5, 6.4.5, 7.4.5, 8.4.5, 9.4.5, 10.4.5, 11.4.5, 12.4.5, 13.4.5, 14.4.5, 15.4.5, 16.4.5, 1.5.5, 2.5.5, 3.5.5, 4.5.5, 5.5.5, 6.5.5, 7.5.5, 8.5.5, 9.5.5, 10.5.5, 11.5.5, 12.5.5, 13.5.5, 14.5.5, 15.5.5, 16.5.5, 1.6.5, 2.6.5, 3.6.5, 4.6.5, 5.6.5, 6.6.5, 7.6.5, 8.6.5, 9.6.5, 10.6.5, 11.6.5, 12.6.5, 13.6.5, 14.6.5, 15.6.5, 16.6.5, 1.7.5, 2.7.5, 3.7.5, 4.7.5, 5.7.5, 6.7.5, 7.7.5, 8.7.5, 9.7.5, 10.7.5, 11.7.5, 12.7.5, 13.7.5, 14.7.5, 15.7.5, 16.7.5, 1.8.5, 2.8.5, 3.8.5, 4.8.5, 5.8.5, 6.8.5, 7.8.5, 8.8.5, 9.8.5, 10.8.5, 11.8.5, 12.8.5, 13.8.5, 14.8.5, 15.8.5, 16.8.5, 1.1.6, 2.1.6, 3.1.6, 4.1.6, 5.1.6, 6.1.6, 7.1.6, 8.1.6, 9.1.6, 10.1.6, 11.1.6, 12.1.6, 13.1.6, 14.1.6, 15.1.6, 16.1.6, 1.2.6, 2.2.6, 3.2.6, 4.2.6, 5.2.6, 6.2.6, 7.2.6, 8.2.6, 9.2.6, 10.2.6, 11.2.6, 12.2.6, 13.2.6, 14.2.6, 15.2.6, 16.2.6, 1.3.6, 2.3.6, 3.3.6, 4.3.6, 5.3.6, 6.3.6, 7.3.6, 8.3.6, 9.3.6, 10.3.6, 11.3.6, 12.3.6, 13.3.6, 14.3.6, 15.3.6, 16.3.6, 1.4.6, 2.4.6, 3.4.6, 4.4.6, 5.4.6, 6.4.6, 7.4.6, 8.4.6, 9.4.6, 10.4.6, 11.4.6, 12.4.6, 13.4.6, 14.4.6, 15.4.6, 16.4.6, 1.5.6, 2.5.6, 3.5.6, 4.5.6, 5.5.6, 6.5.6, 7.5.6, 8.5.6, 9.5.6, 10.5.6, 11.5.6, 12.5.6, 13.5.6, 14.5.6, 15.5.6, 16.5.6, 1.6.6, 2.6.6, 3.6.6, 4.6.6, 5.6.6, 6.6.6, 7.6.6, 8.6.6, 9.6.6, 10.6.6, 11.6.6, 12.6.6, 13.6.6, 14.6.6, 15.6.6, 16.6.6, 1.7.6, 2.7.6, 3.7.6, 4.7.6, 5.7.6, 6.7.6, 7.7.6, 8.7.6, 9.7.6, 10.7.6, 11.7.6, 12.7.6, 13.7.6, 14.7.6, 15.7.6, 16.7.6, 1.8.6, 2.8.6, 3.8.6, 4.8.6, 5.8.6, 6.8.6, 7.8.6, 8.8.6, 9.8.6, 10.8.6, 11.8.6, 12.8.6, 13.8.6, 14.8.6, 15.8.6, 16.8.6, 1.1.7, 2.1.7, 3.1.7, 4.1.7, 5.1.7, 6.1.7, 7.1.7, 8.1.7, 9.1.7, 10.1.7, 11.1.7, 12.1.7, 13.1.7, 14.1.7, 15.1.7, 16.1.7, 1.2.7, 2.2.7, 3.2.7, 4.2.7, 5.2.7, 6.2.7, 7.2.7, 8.2.7, 9.2.7, 10.2.7, 11.2.7, 12.2.7, 13.2.7, 14.2.7, 15.2.7, 16.2.7, 1.3.7, 2.3.7, 3.3.7, 4.3.7, 5.3.7, 6.3.7, 7.3.7, 8.3.7, 9.3.7, 10.3.7, 11.3.7, 12.3.7, 13.3.7, 14.3.7, 15.3.7, 16.3.7, 1.4.7, 2.4.7, 3.4.7, 4.4.7, 5.4.7, 6.4.7, 7.4.7, 8.4.7, 9.4.7, 10.4.7, 11.4.7, 12.4.7, 13.4.7, 14.4.7, 15.4.7, 16.4.7, 1.5.7, 2.5.7, 3.5.7, 4.5.7, 5.5.7, 6.5.7, 7.5.7, 8.5.7, 9.5.7, 10.5.7, 11.5.7, 12.5.7, 13.5.7, 14.5.7, 15.5.7, 16.5.7, 1.6.7, 2.6.7, 3.6.7, 4.6.7, 5.6.7, 6.6.7, 7.6.7, 8.6.7, 9.6.7, 10.6.7, 11.6.7, 12.6.7, 13.6.7, 14.6.7, 15.6.7, 16.6.7, 1.7.7, 2.7.7, 3.7.7, 4.7.7, 5.7.7, 6.7.7, 7.7.7, 8.7.7, 9.7.7, 10.7.7, 11.7.7, 12.7.7, 13.7.7, 14.7.7, 15.7.7, 16.7.7, 1.8.7, 2.8.7, 3.8.7, 4.8.7, 5.8.7, 6.8.7, 7.8.7, 8.8.7, 9.8.7, 10.8.7, 11.8.7, 12.8.7, 13.8.7, 14.8.7, 15.8.7 and 16.8.7.

Exemplary bis esters include bis(pivaloyloxymethyl)PMEA (i.e. bis(pivaloyloxymethyl)-9-(2-phosphonylmethoxyethyl)adenine), bis(pivaloyloxymethyl)HPMPC, bis(pivaloyloxymethyl)D4AMPI, bis(pivaloyloxymethyl)D4TMPI, bis(N-ethylmorpholino)PMEA, bis(N-ethylmorpholino)HPMPC, bis(N-ethylmorpholino)PMPDAP, bis(N-

ethylmorpholino)HPMPA, bis(N-ethylmorpholino)PMEG, bis(N-ethylmorpholino)D4AMPI, bis(N-ethylmorpholino)D4TMPI, bis(phenyl)PMEA, bis(phenyl)HPMPC, bis(phenyl)HPMPA, bis(phenyl)D4AMPI, bis(phenyl)D4TMPI, bis(t-butyl)PMEA, bis(t-butyl)D4AMPI, bis(t-butyl)D4TMPI, bis(t-butyl)HPMPC, bis(2-ethoxyphenyl)PMEA, bis(2-ethoxyphenyl)HPMPC, , bis(4-fluorophenyl)PMEA, bis(4-fluorophenyl)HPMPC, bis(3,5-dimethoxyphenyl)PMEA, bis(3,5-dimethoxyphenyl)HPMPC and the like. L<sup>1</sup> is an amino acid which is, in general, esterified at free  $\alpha$ -carboxyl group(s) by R<sup>4</sup>, or is a dipeptide, tripeptide or oligopeptide which is optionally esterified at the free  $\alpha$ -carboxyl group by R<sup>4</sup>. L<sup>2</sup> is an ester or thioester group. Suitable L<sup>2</sup> esters (and the corresponding thioesters) include methyl ester, ethyl ester, propyl ester, isopropyl ester, butyl ester, t-butyl ester, phenyl ester, benzyl ester, N-ethylmorpholino ester (-O-CH<sub>2</sub>-CH<sub>2</sub>-N[(CH<sub>2</sub>)<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>O]), pivaloyloxymethyl ester (-O-CH<sub>2</sub>-O-C(O)-C(CH<sub>3</sub>)<sub>3</sub>) and the like. The suitability of the presence or absence of any particular L<sup>2</sup> or R<sup>4</sup> group is determined by stability and/or bioavailability assays (e.g., stability assay in aqueous conditions such as low pH/intestinal lumen conditions or assay in the presence of cellular extracts containing esterases or by bioavailability assay using animal models) known in the art. These assays are routinely performed by the skilled artisan.

The bis ester is then converted to a monoester by chemical hydrolysis in base or acid according to the bis ester used. For example, treatment with NaOH (0.5 to 2 N) or NH<sub>4</sub>OH in a solvent such as THF (tetrahydrofuran), dioxane or an alcohol for 1 to 24 hours at 22° to 90° is suitable for most esters. The choice of solvent will depend on the characteristics of the bis ester used. The stability of the ester groups of phosphonate bis esters and phosphonate bis thioesters toward hydrolysis is unequal and provides a means for obtaining the monoester. Selection of hydrolysis conditions is determined by routine testing. Alkaline hydrolysis yields the phosphonate monoester and a corresponding alcohol or phenol. L<sup>1</sup> is then linked to the monoester or monothioester using reagents and conditions (i.e., a 1:1 mixture of triphenylphosphine (PPh<sub>3</sub>) and 2,2'-dipyridyl disulfide in a suitable solvent such as pyridine or DMF) essentially as described for synthesis of bis amidates.

Nucleoside bis esters of formulas VI, VII and VIII compounds are shown in Figures 4 - 7. 3',4'-Unsaturated nucleosides that are used as a starting material was previously described (Zemlicka, et al J Am Chem Soc (1970) 92:4744-4745). 4'-Modified nucleosides have also been described (Yang, et al Tet Lett (1992) 33: 41-44; Yang, et al Tet Lett (1992) 33: 37-40; Prisbe, et al Nucleosides and Nucleotides as Antitumor and Antiviral Agents (1993) Plemun Press, New York, Chu, C.K. et al eds., p. 101-113). The phosphonate ester is condensed with the unsaturated nucleoside using an oxidizing agent such as MCPBA (m-chloroperoxybenzoic acid), IBr or N-iodosuccinimide (NIS). The choice of a particular oxidizing agent will be guided by considerations such as the type of heterocyclic base or sugar substituent that is present. For example, IBr may not be generally compatible with a substituent such as azide (at R<sup>27</sup> or R<sup>33</sup>) or 1-propynyl (at B). In these cases, NIS or MCPBA is used. A further example is reduction of the 2',3'-double bond using H<sub>2</sub>/Pd/C, which is generally not compatible with an alkynyl group that can be present at B. In this case, the alkynyl group would be added to an appropriate heterocyclic base (a purine such as 7-deaza-7-iodoadenine or 7-deaza-7-iodoguanine, etc or a pyrimidine such as 5-iodocytosine, 5-iodouracil, uracil, etc) that is later converted to the alkynyl derivative (7-deaza-7-(1-propynyl)adenine, 5-(1-propynyl)uracil, etc) using an alkyne such as propyne and palladium (08/050,698; PCT/US92/10115; Hobbs et al, J Org Chem (1989) 54:3420-3422). For Figures 3 - 7, R<sup>33</sup> is H, OH, TBSO, halogen, cyano, CH<sub>2</sub>N<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy (including OCH<sub>3</sub>), CH<sub>2</sub>OH or azido; R<sup>34</sup> is H, OH halogen (fluorine is preferred), azide, O-alkyl (C<sub>1</sub>-C<sub>6</sub> including O-methyl and O-ethyl), S-alkyl (C<sub>1</sub>-C<sub>6</sub> including S-methyl and S-ethyl) and O-alkenyl (including O-allyl); R is as defined above, except that for the structure (RO)<sub>2</sub>P(O)-CH<sub>2</sub>-OH, R is not hydrogen, and R includes C<sub>1</sub>-C<sub>20</sub> alkoxyacyl groups including methoxyacyl (pivaloyloxymethyl, adamantoyl oxymethyl and the like) and ethoxyacyl (pivaloyloxyethyl and the like) moieties; TBSO is t-butyldimethylsilyl ether. The phosphonates and monoesters shown in Figures 4 - 7 are converted to bis amidates or mixed amidate ester compounds using reagents and conditions (e.g., a 1:1 mixture of triphenylphosphine (PPh<sub>3</sub>) and 2,2'-dipyridyl disulfide in a suitable solvent such as pyridine or DMF) essentially as described above.

35

Mixed bis amidate synthesis. Synthesis of compounds of formula Id where  $L^1$  and  $L^2$  are both amino acids or where  $L^1$  is an amino acid and  $L^2$  is an amine ( $NH_2$ ,  $NHR^6$ ,  $N(R^6)_2$ ) but are not both the same is accomplished by direct conversion as described above for bis amidates followed by separation of the final products. Another method to synthesize mixed bis amidates is amidation of an appropriate phosphonate monoester to give a compound of formula Id, followed by removal of the ester group under conditions that do not remove the first amide. Synthesis of phosphonate monoester compounds has been described (EP 481 214). This compound is then converted to a mixed bis amide by condensation with a second amino acid to yield the final product as described (i.e., using a 1:1 mixture of triphenylphosphine and 2,2'-dipyridyl disulfide).


Mono amidate synthesis. Synthesis of compounds of formula Ib where  $L^1$  is an amino acid and  $X^1$  is O (oxygen) is accomplished essentially as described for bis amidate synthesis using a cyclic nucleotide analog such as cHPMPC (cyclic HPMPC), cHPMPA, cHPMPDAP, cHPMPG and the like. Cyclic HPMP series compounds (cHPMPC, etc) are prepared by direct dehydration of the corresponding HPMP nucleotide analog using DCC (dicyclohexylcarbodiimide) or using 4-morpholino-N,N'-dicyclohexylcarboxamide as described (Ho et al *Mol Pharmacol* (1992) 41:197-202). The cyclic phosphonate is condensed with an optionally protected amino acid ester in the presence of a 1:1 mixture of triphenylphosphine and 2,2'-dipyridyl disulfide in a suitable solvent such as pyridine or DMF.

Synthesis of formula Ib compounds where  $X^1$  is S is accomplished as shown in Figure 1. Conversion of the six-membered heterocycle to an amidate is accomplished in essentially the same manner as described (i.e., using triphenylphosphine and 2,2'-dipyridyl disulfide).

Synthesis of formula IV compounds where  $R^{26}$  is S and  $R^{25}$  and  $R^{29}$  are O is accomplished as shown in Figure 3. The starting material is synthesized by reaction of thiolacetic acid (Aldrich Cat. No. T3,080-5), bromoacetaldehyde diethyl acetal (Aldrich Cat. No. 12,398-6) and potassium tert-butoxide (Aldrich Cat. No. 15,667-1) in DMF. Synthesis of **1** where  $R^{34}$  is H is accomplished using neat  $(EtO)_3CH$ . Synthesis of **1** where  $R^{34}$  is  $CH_2CN$  or  $CF_3$  is accomplished using  $(EtO)_3CH_2CN$  or  $(EtO)_3CF_3$  in methylene chloride with a

catalytic acid (such as p-toluenesulfonic acid). Conversion of the thiaorthoester **1** to the phosphonate **2** is accomplished using an acid such as tosic acid or perchloric acid in catalytic amounts. The resulting bis ester is then converted to a bis amidate in essentially the same manner as described (i.e., using triphenylphosphine and 2,2'-dipyridyl disulfide). Mixed ester-amidate compounds are obtained by removing a single ester from the bis ester using base (NaOH, NH<sub>4</sub>OH, etc) as described. The phosphonate **3** is obtained by treatment with a base such as TMSBr or TMSI in a solvent (such as methylene chloride, DMF or acetonitrile) in the presence of lutidine (where R is alkyl, aryl or substituted aryl, acyloxyalkyl such as isopropyl, phenyl, 2-ethoxyphenyl) or by treatment with Pd/C/H<sub>2</sub> (where R is alkaryl or substituted alkaryl such as benzyl and the like). R<sup>34</sup> in Figure 3 is H, CF<sub>3</sub> or CH<sub>2</sub>CN.

Protected heterocyclic base compounds. The present invention includes nucleotide analogs that comprise a protected heterocyclic base. These compounds are useful as synthetic intermediates and/or, as therapeutic agents *per se*. Protected heterocyclic base compounds structures, their isomers, tautomers and the salts of such compounds having the formula (R<sup>35</sup>O)<sub>2</sub>P(O)-Z-B<sup>1</sup>, (L<sup>1A</sup>O)(L<sup>2A</sup>O)P(O)-Z-B<sup>1</sup>, (HO)<sub>2</sub>P(O)-Z-B<sup>1</sup>




where L<sup>1A</sup> is L<sup>1</sup>, R or NHR<sup>40</sup>, wherein R<sup>40</sup> is C<sub>1</sub>-C<sub>20</sub> alkyl; L<sup>2A</sup> is L<sup>2</sup>, R<sup>35</sup> or NHR<sup>40</sup>; B<sup>1</sup> is a protected heterocyclic base having the formula Xa, XIa, XIb, XIIa or XIIIa previously defined.

Suitable exemplary Z include compounds of formulas IV, V, VI, VII, VIII, -CH<sub>2</sub>-O-CH<sub>2</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-O-C#H(CH<sub>3</sub>)-CH<sub>2</sub>- and -CH<sub>2</sub>-O-C#H(CH<sub>2</sub>OH)-CH<sub>2</sub>- having a heterocyclic base with an exocyclic amine can be converted to nucleotide analog amidates or esters comprising a protected heterocyclic base either by reacting the nucleotide analog amidate or ester with R<sup>36</sup>C(O)Cl or (CH<sub>3</sub>O)<sub>2</sub>CHR<sup>38</sup>. Protected heterocyclic bases include species having protecting groups at exocyclic amine groups such as the N<sup>4</sup>-amine of cytosine, the N<sup>6</sup>-

amine of adenine and the N<sup>2</sup>-amine of guanine. The phosphonate moiety of compounds containing B<sup>1</sup> may be present as an ester, an amidate or as the free acid.

5 Bases having NHR<sup>40</sup> at an exocyclic amine are synthesized to obtain a protected pyrimidine or purine essentially as described (Gilliam Anal. Biochem. (1986) 157:199; Gallo-Rodriguez J. Med. Chem. (1994) 37:636; Maillard J. Pharm. Sci. (1994) 83:46).

The exemplary reaction schemes used to synthesize protected heterocyclic base compounds shown below utilize cHPMPC as an example.  
 10 Analogous reactions will generate compounds comprising other Z moieties such as -CH<sub>2</sub>-O-CH<sub>2</sub>-CH<sub>2</sub>- or -CH<sub>2</sub>-O-CH(CH<sub>3</sub>)-CH<sub>2</sub>- linked to B<sup>1</sup>. Phosphonate alkyl and aryl esters of compounds comprising B<sup>1</sup> are prepared, using HPMPC and cHPMPC as an example, according to the following procedures





wherein R<sup>36</sup> is as defined above. Either procedure is readily adapted to synthesizing compounds containing protected heterocyclic bases other than cytosine, e.g., adenine, guanine, 2,6-diaminopurine or 2-aminopurine. Exemplary R<sup>35</sup> and/or R<sup>36</sup>, which can be the same or different, include

phenyl, substituted phenyl,  $-C_{10}H_{15}$  (where  $C_{10}H_{15}$  is adamantoyl),  $-CH_2-C_6H_5$ ,  $-C_6H_5$ ,  $-C(CH_3)_3$ ,  $-CH(CH_3)_2$ ,  $-CH_2CH_3$ , methyl, ethyl, butyl, t-butyl, heptyl, nonanyl, undecanyl, lauryl, steryl, undecenyl and the like. The amide linkage is conveniently formed by reaction of the acyl chloride with the exocyclic amine linked to the base. When  $R^1$  is linked to the free phosphonate the resulting ester will comprise a single isomer or a scalemic mixture at the phosphorus atom. Low temperature reaction conditions (lower than about  $-20^\circ$ , e.g., about  $-20^\circ$  to about  $-40^\circ$  C or about  $-40^\circ$  to about  $-80^\circ$  C) tend to favor single isomer products, while reaction at higher temperatures (above about  $-20^\circ$ , e.g.  $-20^\circ$  to  $40^\circ$  C) generally results in a scalemic mix. When a scalemic mixture is obtained, the isomers can be conveniently separated by, for example, HPLC, although the mixture can be used, for example, as a synthetic intermediate or as an active antimicrobial agent, without resolution. Synthesis of the phenyl ester of cHPMPC at  $-78^\circ$  C by reaction of the chloridate and phenoxide yielded a scalemic mixture consisting of about  $\geq 90\%$  of the product as one isomer (isomer #1) at the phosphorus atom while the remaining  $\sim \leq 10\%$  was present as the other isomer (isomer #2). The scalemic mixture was converted to isomer #2 ( $\geq 90\%$ ) by incubation at room temperature for about 10 minutes (about 10 to 30 minutes is generally suitable) with a catalytic amount of sodium phenoxide in DMF. This method can be used to convert one isomer of cHPMP-B or cHPMP-B<sup>1</sup> (such as cHPMPC or cHPMPA) aryloxy or alkoxy ester to the other isomer with catalytic amounts of the corresponding aryloxide ion or alkoxide ion.

The cHPMPC pivaloyloxymethyl ester synthesis yields a scalemic mixture at the phosphorus atom. The mixture was separated by HPLC into the two isomers which were then exposed to an rat intestinal homogenate or to a rat intestinal wash. One of the isomers was converted to cHPMPC after incubation in the homogenate while the other isomer was converted to HPMPC pivaloyloxymethyl monoester. Both isomers were converted to HPMPC pivaloyloxymethyl monoester after incubation in the intestinal wash. These results suggested that (1) in at least some cases, enzyme activity can have a differential effect on the metabolic fate of a cHPMPC ester depending on which phosphorus isomer is present and (2) chemical activity (i.e., the acidity of the intestinal wash) can affect the metabolic fate of a given compound in a manner that differs from enzyme activity.


A method to obtain heterocyclic bases comprising the  $C(O)R^{36}$  protecting group is accomplished as follows using the acyl chloride ( $R^{36}C(O)Cl$ ) using HPMPC and cHPMPC as an example





wherein Tr is the hydroxyl protecting group trityl. The detritylation step is accomplished by acid treatment, such as 80% acetic acid at about 10° to 60° C for 1-2 hours. The R<sup>35</sup> moiety is removed using a Lewis acid such as TMSBr to yield the free phosphonate.


- 5        Phosphonate compounds comprising B<sup>1</sup> and a C<sub>2</sub>-C<sub>20</sub> 1-acyloxy-1-alkyl or a C<sub>4</sub>-C<sub>20</sub> 1-acyloxy-1-alkyl-1-aryl ester group are prepared as follows



- 10        wherein R<sup>37</sup> is C<sub>1</sub>-C<sub>20</sub> alkyl which is unsubstituted or substituted by substituents independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>1</sub>-C<sub>6</sub> haloalkyl (1 to 3 halogen atoms), cyano, nitro, OH, O, NH and halogen (including ethyl, propyl, isopropyl, t-butyl, isobutyl and adamantoyl), or C<sub>3</sub>-C<sub>10</sub> aryl which is unsubstituted or substituted by
- 15        substituents independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>1</sub>-C<sub>6</sub> haloalkyl (1 to 3 halogen atoms), cyano, nitro, OH, O, N and halogen (including phenyl, and 3- or 4-pyridyl).

The amine protecting group =CR<sup>41</sup>N(R<sup>38</sup>)<sub>2</sub> is incorporated into an exocyclic amine to yield protected heterocyclic base compounds as follows

20



Exemplary R<sup>38</sup> alkyl groups include methyl, ethyl, propyl, isopropyl, cyclopropyl, butyl, isobutyl and cyclobutyl. In general, both R<sup>38</sup> alkyl groups will be the same. The reaction can be carried out in dry DMF at room temperature (about 20-30° C) as previously described (Kerr et al *J. Pharm. Sci.* (1994) 83:582; Kerr et al *J. Med. Chem.* (1992) 35:1996), or DMF can be substituted with CH<sub>3</sub>CN and 4 Å molecular sieves. Exemplary compounds include species where R is hydrogen, alkyl (including ethyl, propyl, isopropyl), aryl (including phenyl) or acyloxymethyl. Protected heterocyclic bases where R<sup>41</sup> is hydrogen are stable under neutral anhydrous conditions and are generally labile under acidic aqueous conditions. When R<sup>41</sup> is methyl, the protecting group is more stable to aqueous acidic or basic conditions.


Compounds containing a protected heterocyclic base and 1 or 2 amino acids, dipeptides or oligopeptides attached to the phosphorus atom via an amidate linkage are obtained as described for synthesis of bis-amidate or amidate-ester compounds.

Table 5A lists R<sup>35</sup> ester and L<sup>1</sup> amidate moieties that can be incorporated into the phosphorus atom of both cyclic Z moieties (such as cHPMPC comprising a protected heterocyclic base or cHPMPC) or linear Z moieties (such as HPMPC comprising a protected heterocyclic base or PMEA comprising a protected heterocyclic base or PMEA). Esters of structures 1-5, 8-10 and 16, 17, 19-22 are synthesized by reacting a nucleotide analog (such as cHPMPC) the corresponding halide (chloride or acyl chloride and the like) and N,N-dicyclohexyl-N-morpholine carboxamidine (or another base such as DBU, triethylamine, CsCO<sub>3</sub>, N,N-dimethylaniline and the like) in DMF (or

other solvent such as acetonitrile or N-methylpyrrolidone). Esters of structures 5-7, 11, 12, 21, and 23-26 are synthesized by reaction of the alcohol or alkoxide salt (or the corresponding amines in the case of compounds such as 13, 14 and 15) with a nucleotide analog monochlorophosphonate or dichlorophosphonate (such as cHPMPC monochlorophosphonate or PMEA dichlorophosphonate) or another activated phosphonate.

TABLE 5A

	1. $-\text{CH}_2-\text{C}(\text{O})-\text{N}(\text{R}^7)_2^*$	10. $-\text{CH}_2-\text{O}-\text{C}(\text{O})-\text{C}(\text{CH}_3)_3$
10	2. $-\text{CH}_2-\text{S}(\text{O})(\text{R}^7)$	11. $-\text{CH}_2-\text{CCl}_3$
	3. $-\text{CH}_2-\text{S}(\text{O})_2(\text{R}^7)$	12. $-\text{C}_6\text{H}_5$
	4. $-\text{CH}_2-\text{O}-\text{C}(\text{O})-\text{CH}_2-\text{C}_6\text{H}_5$	13. $-\text{NH}-\text{CH}_2-\text{C}(\text{O})\text{O}-\text{CH}_2\text{CH}_3$
	5. 3-cholesteryl	14. $-\text{N}(\text{CH}_3)-\text{CH}_2-\text{C}(\text{O})\text{O}-\text{CH}_2\text{CH}_3$
	6. 3-pyridyl	15. $-\text{NHR}^{40}$
15	7. N-ethylmorpholino	16. $-\text{CH}_2-\text{O}-\text{C}(\text{O})-\text{C}_{10}\text{H}_{15}$
	8. $-\text{CH}_2-\text{O}-\text{C}(\text{O})-\text{C}_6\text{H}_5$	17. $-\text{CH}_2-\text{O}-\text{C}(\text{O})-\text{CH}(\text{CH}_3)_2$
	9. $-\text{CH}_2-\text{O}-\text{C}(\text{O})-\text{CH}_2\text{CH}_3$	18. $-\text{CH}_2-\text{C}\equiv\text{H}(\text{OC}(\text{O})\text{CH}_2\text{R}^7)-\text{CH}_2-$ $(\text{OC}(\text{O})\text{CH}_2\text{R}^7)^*$



25 \* - Each  $\text{R}^7$  is the same or different (includes methyl, ethyl, propyl, isopropyl and t-butyl).

All citations are hereby expressly incorporated by reference. The following examples are illustrative and do not limit the scope of this invention.

5        Example 1: Synthesis of phosphonate amidate compounds. The compounds of structural formula Id shown are in Table 6 (bis(glycyl benzyl ester)PMEA (compound Ex 4), bis(alanyl benzyl ester)PMEA (Ex 1), bis(phenylalanyl benzyl ester)PMEA (Ex 5), etc. Compounds Ex 1 - Ex 12 were synthesized by the following procedure. PMEA (Z-B = -CH<sub>2</sub>-O-CH<sub>2</sub>-CH<sub>2</sub>-B, where B is adenin-9-yl) (0.3 g; 1.1 mmol) and amino acid ester•HCl (2.2 mmol; Sigma) were suspended in dry pyridine (6 mL) containing triethylamine (0.3 mL; 22.2 mmol), followed by addition to a mixture of freshly prepared triphenylphosphine (3.3 mmol) and 2,2'-dipyridyl disulfide (3.3 mmol) in pyridine (3 mL). The mixture was stirred at room temperature overnight, concentrated and partitioned between methylene chloride and water. The organic solution was dried over MgSO<sub>4</sub>, concentrated and purified by flash column chromatography on silica gel.

Ex 14 was synthesized using freshly prepared triphenylphosphine (6.0 mmol) and 2,2'-dipyridyl disulfide (6.0 mmol) in pyridine (20 mL) at room temperature to which PMEA (2.0 mmol) was added. The suspension was stirred for 10 min. and ethyl sarcosine HCl (N-methylglycine HCl ethyl ester; 1.2 g, 8.0 mmol) was added. The suspension was warmed to 90°C and stirred for 24 hours. Crude product was concentrated by rotary evaporation and purified by silica flash chromatography (mobile phase 1% methanol gradient to 20% methanol/80% methylene chloride).

Compound Ex 13 was synthesized in a similar manner using PMEA and phenylalanine N-ethylmorpholino ester.

TABLE 6

Compound	L1
Ex 1	-NH-CH(CH <sub>3</sub> )-C(O)OCH <sub>2</sub> C <sub>6</sub> H <sub>5</sub>
Ex 2	-NH-CH(CH <sub>2</sub> C <sub>6</sub> H <sub>5</sub> )-C(O)OCH <sub>2</sub> C <sub>6</sub> H <sub>5</sub>
Ex 3	-NH-CH(CH <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub> )-C(O)OCH <sub>2</sub> C <sub>6</sub> H <sub>5</sub>
Ex 4	-NH-CH <sub>2</sub> -C(O)OCH <sub>2</sub> C <sub>6</sub> H <sub>5</sub>
Ex 5	-NH-CH(CH <sub>3</sub> )-C(O)OC <sub>2</sub> H <sub>5</sub>

Ex 6	-NH-CH(CH <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub> )-C(O)OC <sub>2</sub> H <sub>5</sub>
Ex 7	-NH-CH <sub>2</sub> -C(O)OC <sub>2</sub> H <sub>5</sub>
Ex 8	-NH-CH(CH <sub>2</sub> C <sub>6</sub> H <sub>5</sub> )-C(O)OC(CH <sub>3</sub> ) <sub>3</sub>
Ex 9	-NH-CH(CH <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub> )-C(O)OC(CH <sub>3</sub> ) <sub>3</sub>
5 Ex 10	-NH-CH(CH <sub>3</sub> )-C(O)OC(CH <sub>3</sub> ) <sub>3</sub>
Ex 11	-NH-CH <sub>2</sub> -C(O)OC(CH <sub>3</sub> ) <sub>3</sub>
Ex 12	-NH-CH(CH <sub>2</sub> C <sub>6</sub> H <sub>5</sub> )-C(O)OC <sub>2</sub> H <sub>5</sub>
Ex 13	-NH-CH(CH <sub>2</sub> C <sub>6</sub> H <sub>5</sub> )C(O)O-(CH <sub>2</sub> ) <sub>2</sub> -N[(CH <sub>2</sub> ) <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> ]O
Ex 14	-N(CH <sub>3</sub> )-CH <sub>2</sub> -C(O)OC <sub>2</sub> H <sub>5</sub>

10

Example 2: Antiviral activity. Compounds were individually tested for activity against HSV-1 and/or HSV-2. HSV-2 (strain 414-92) was tested using MA 104 cells in the following assay protocol. 96-Well plates were seeded with 1 x 10<sup>4</sup> MA 104 cells per well using 200  $\mu$ L minimal essential medium (MEM) containing 10% calf serum per well, and incubated overnight at 37°C. The compounds were dissolved in MEM Earle's Salts without serum. The medium was removed by aspiration and 100  $\mu$ L MEM Earle's Salts without serum was added to the wells. Serial 3-fold dilutions of the compounds were prepared by serial transfer of 50  $\mu$ L of medium from wells containing compound to wells lacking compound. The plates were incubated 15 minutes at 37°C followed by addition of 100 PFU/well of virus in MEM Earle's Salts with 2% fetal bovine serum. The plates were then incubated at 37°C for three days until approximately 90% of the cells in virus infected control wells containing no compound were killed. Following incubation, medium was aspirated and the wells were washed with sterile PBS. 100  $\mu$ L 0.5% crystal violet in 20% methanol was then added to the wells for 5 minutes, aspirated and the wells were washed two or three times with distilled water. 200  $\mu$ L of 0.01 N HCl was added to the wells and the absorbance of each well at 595 nm was determined. The results, shown in Table 6, were expressed as the IC<sub>50</sub>, the concentration ( $\mu$ M) that inhibits cell killing mediated by HSV-2 by 50%. IC<sub>50</sub> values varied from 2  $\mu$ M to >100  $\mu$ M compared to an IC<sub>50</sub> for PMEA of 21  $\mu$ M. Thus, some of the compounds were more active against HSV-1 than

PMEA.. The toxicity of the compounds were expressed as the CC<sub>50</sub>, the concentration that kills 50% of uninfected cells.

The compounds were also tested for activity against the KOS strain of HSV-1 in VERO cells. The results, shown in Table 7, were expressed as the EC<sub>50</sub>, the concentration (μM) that inhibits cell killing mediated by HSV-2 by 50%. EC<sub>50</sub> values varied from 2 μM to >200 μM compared to an EC<sub>50</sub> for PMEA of 138 μM. Thus, some of the compounds were more active against HSV-2 than PMEA.

Table 7

10

compound	HSV-1	HSV-2	
	EC <sub>50</sub>	IC <sub>50</sub>	CC <sub>50</sub>
Ex 7	>200	>100	>100
Ex 5	nt*	>100	>100
Ex 6	20	33	>100
Ex 12	nt	20	80
Ex 11	>200	>100	>100
Ex 10	>200	>100	>100
Ex 9	63	63	>100
Ex 8	3	9	20
Ex 4	nt	60	>100
Ex 1	nt	20	>100
Ex 3	nt	2	30
Ex 2	nt	4	20

\* nt - not tested

15 Example 3: PMEA, monophenyl ester, mono N-ethylmorpholino-phenylalanyl phosphoramidate. Bis(phenyl)PMEA is selectively hydrolyzed to the monophenyl ester of PMEA using NaOH in THF. The reaction mixture is neutralized with acid (1 N HCl), and the monophenyl PMEA is isolated by filtration. The anhydrous monophenyl PMEA and 2 equivalents of a freshly prepared 1:1 mixture of triphenylphosphine and 2,2'-dipyridyl disulfide in pyridine is condensed with 1 equivalent of phenylalanine N-ethyl-morpholino ester in triethylamine and pyridine to afford the title

20

compound. The title compound is recovered by evaporation of the solvents under reduced pressure and purified by silica gel chromatography.

5        Example 4: Antiviral activity of PME A esters. PME A and PME A esters  
were tested for inhibition of cytopathic effects by HSV II in MA 104 cells as  
described except that CPE was determined after incubation with virus by  
addition of 100 $\mu$ L XTT, 1 mg/mL in deficient DME containing 25  $\mu$ M PMF  
followed by measuring absorbance. The esters tested were bis(POM)PME A,  
bis(phenyl)PME A, monophenylPME A, bis(3-dimethylaminophenyl)PME A,  
10    bis(3-methoxyphenyl)PME A, bis(2-carboethoxyphenyl)PME A, bis(adamantoyl  
oxymethyl)PME A, bis(4-fluorophenyl)PME A and bis(2-ethoxyphenyl)PME A.  
All of the compounds tested were active, which indicated that the ester  
groups were removed, thereby allowing free PME A to inhibit virus  
replication and/or cytopathic effects. The IC<sub>50</sub> and CC<sub>50</sub> of PME A in the assay  
15    was 19.3  $\mu$ M and 2000  $\mu$ M respectively and the IC<sub>50</sub> and CC<sub>50</sub> of  
bis(POM)PME A in the assay was 0.5  $\mu$ M and >10  $\mu$ M respectively. IC<sub>50</sub> values  
for the mono and bis esters ranged from 1.1  $\mu$ M to 67.5  $\mu$ M and the CC<sub>50</sub>  
values ranged from 70  $\mu$ M to 500  $\mu$ M.

20        Example 5: Oral bioavailability of nucleotide analog amidates and  
PME A esters. Nucleotide analog amidates and nucleotide analogs are tested  
for their bioavailability when administered to cynomolgous (or rhesus)  
monkeys by oral, subcutaneous or intramuscular routes. Bioavailability is  
determined by measuring PME A levels in plasma or urine at different times  
25    after administering the drug using radiolabeled (<sup>3</sup>H, <sup>14</sup>C, etc) compound or,  
for compounds having adenine, essentially as described (Naesens, et al, Clin  
Chem (1992) 38:480-485; Russell, et al, J Chromatogr (Netherlands) (1991)  
572:321-326). Radiolabeled compounds are obtained commercially (Moravek  
Biochemicals, Brea, CA) or by standard procedures, such as catalytic hydrogen  
30    exchange for <sup>3</sup>H labeling. Compounds such as bis(2-ethoxyphenyl)PME A,  
bis(2-carboethoxyphenyl)PME A, bis(O-benzylphenylalanyl)PME A, bis(3,5-  
dimethoxyphenyl)PME A, bis(4-fluorophenyl)PME A, bis(adamantoyl  
oxymethyl)PME A, bis(phenyl)PME A, bis(3-methoxyphenyl)PME A are tested  
for oral bioavailability by administering about 10 - 30 mg/Kg (usually 15 to 25  
35    mg/Kg) containing about 20 - 50  $\mu$ Ci/Kg (usually about 40  $\mu$ Ci/Kg) of

radiolabeled compound, followed by withdrawing blood samples at several times after administration (exemplary time points are 0.1, 0.25, 0.5, 1.0, 1.5, 2.0, 2.5, 3.0, 4, 6, 12, 18, 24, 36, 48, 72, 96 hours after administration), obtaining plasma and determining the amount of radiolabeled compound present per  
5 volume (about 0.1 - 1.0 mL) of serum. Oral bioavailability of the tested compounds is 2 - 80% (or any value between 2% and 80% in 1% increments), preferably 10 - 80% and more preferably 15 to 80%. The oral bioavailability of bis(POM)PMEA by this type of assay is typically about 25% in monkeys and  
10 PMEA is about 2 - 4% (Balzarini et al, Animal Models in AIDS (1990) p. 131-138, Schellekens, H. et al (ed), Elsevier Science Publications, Amsterdam) while nucleotide analog amidates and nucleotide analogs (including mono- and diesters) can have oral bioavailabilities of about 5%, 10%, 15%, 30%, 40%, 50%, 60% or 80%.

Total radioactivity in plasma is determined by mixing about 200  $\mu$ L of  
15 plasma with a scintillation counting cocktail (such as 10 mL of Scinti-Safe plus LSC cocktail) and counting in a scintillation counter (usually for about 5 - 30 minutes). Detailed analysis of the radiochemical composition is accomplished using about 350  $\mu$ L of plasma, denaturing proteins in the serum (using about 700  $\mu$ L 0.1% trifluoroacetic acid in acetonitrile for example),  
20 drying the resulting sample under reduced pressure, suspending the sample in an appropriate buffer (for example using about 100  $\mu$ L of 2% acetonitrile in 25 mM potassium phosphate buffer with 10 mM tetrabutyl ammonium hydrogen phosphate (TBAHP), pH 6.0 for HPLC analysis), centrifuging the sample and analyzing the supernatant for individual radiolabeled species by  
25 reverse phase HPLC on commercially available columns (The Separation Group, Hesperia, CA; Vydac C18, 5  $\mu$ m, 250 x 4.6 mm column with an injection volume of about 50  $\mu$ L and a flow rate of about 1.0 mL/min. at about 35°C using buffer for 2 minutes followed by a linear gradient to about 65% acetonitrile in 25 mM potassium phosphate buffer with 10 mM TBAHP, pH  
30 6.0 over 13 about minutes). Radiolabel detection is accomplished using means such as commercially available radioactive flow detection systems or scintillation counting systems (Packard, Meridian, CT).

Fluorescence detection of PMEA in plasma is accomplished by measuring fluorescence emission (420 nm, with excitation at about 236 nm)  
35 with a detector (model F2000, Spectra Physics, San Jose, CA) from the HPLC



gradient essentially as described above (2 to 65% acetonitrile). Samples for analysis are prepared from plasma (200  $\mu$ L) by protein precipitation with TFA (400  $\mu$ L 0.1% in acetonitrile), drying and conversion of adenine to N<sup>6</sup>-ethenoadenine in 200  $\mu$ L of reaction buffer (0.34% chloroacetaldehyde, 100 mM sodium acetate, pH 4.5) for 40 minutes at 95°C followed by HPLC analysis using 50  $\mu$ L.

Example 6: Bis(adamantoyl oxymethyl)PMEA ester. DBU (1,8-diazabicyclo[5.4.0]undec-7-ene; 1.53 g, 10 mmol) was added to a suspension of PMEA (1.365 g, 5 mmol) in DMF (25 mL). Adamantoyl oxymethyl chloride (5.72 g, 25 mmol) in DMF (25 mL) was added to the reaction mixture which was then stirred for four days at room temperature and the volatiles were removed under vacuum. The crude product obtained after removal of the solvent was loaded onto a silica gel column and washed with 3% MeOH/CH<sub>2</sub>Cl<sub>2</sub> to remove nonpolar impurities. 1 g (30%) of bis(adamantoyl oxymethyl)PMEA ester was eluted in 8% MeOH/CH<sub>2</sub>Cl<sub>2</sub>. Adamantoyl oxymethyl chloride was obtained by conversion of 1-adamantanecarbonyl chloride (Aldrich No. 11,772-2) with (CH<sub>2</sub>O)<sub>n</sub>/ZnCl<sub>2</sub> and has been described (Bodor, et al J Med Chem (1980) 23 :474-480).

Example 7: Bis(phenyl)PMEA and bis(2-ethoxyphenyl)PMEA esters. PMEA (2.0 g, 7.3 mmol), acetonitrile (20 mL), thionyl chloride (20 mL) and N,N-dimethylformamide (2 drops) were added to a 250 mL single neck round bottom flask equipped with a magnetic stirrer, water cooled condenser and N<sub>2</sub> atmosphere. The flask was immersed in a 85°C oil bath and the resulting suspension was stirred for two hours. The resulting solution was then concentrated to dryness and acetonitrile (50 mL) was added to redissolve the crude chloridate.

To a separate 250 mL single neck round bottom flask equipped with a mechanical stirrer, and N<sub>2</sub> atmosphere, phenol (3.25 g, 35 mmol), tetrahydrofuran (80 mL) and sodium hydride (1.4 g, 34 mmol, 60% (w/w) dispersion in mineral oil) was charged. After stirring for 30 minutes, the solution was cooled to -78°C with a dry ice-acetone bath. The acetonitrile from the previous step was then added drop-wise at a rate that the internal temperature did not rise above -76°C. After the addition was complete, the

resulting suspension was poured into saturated aqueous  $\text{NaHCO}_3$  (100 mL) and extracted with methylene chloride (3 x 150 mL). The combined organic extracts were washed with  $\text{H}_2\text{O}$  (100 mL), brine (100 mL) and dried with anhydrous  $\text{Na}_2\text{SO}_4$ . Concentration by rotary evaporation afforded a yellow solid. Purification by recrystallization (ethyl acetate/hexanes) afforded pure bis(phenyl)PMEA (1.64 g, 53%). Bis(2-ethoxyphenyl)PMEA was made similarly using 2-ethoxyphenol in place of phenol in 36% yield.

Example 8: (R)-9-(2-Di-2 ethoxyphenylphosphonylmethoxypropyl) adenine. To a solution of 2-ethoxyphenol (45 mmol, 6.22 g) in pyridine (75 mL) was added (R)-9-(2-phosphonylmethoxypropyl adenine (PMPA, 15 mmol, 4.3 g), creating a white suspension. A separate solution of 2,2'-dipyridyl disulfide (45 mmol, 9.91 g) and triphenyl phosphine (45 mmol, 11.81 g) in pyridine (75 mL) was added at 22°C in a single portion to the white suspension. Then, triethylamine (30 mmol, 4.18 mL) was added in a single portion to the entire mixture, which was stirred at 75°C for 21 h (TLC:10% MeOH/EtoAc). The dark amber slurry was then coevaporated with toluene (100 mL). It was then dissolved in dichloromethane (200 mL) and extracted twice with water (200 mL). The organic phase was dried ( $\text{Na}_2\text{SO}_4$ ), filtered and concentrated (*in vacuo*) to a brown syrup (25.4 g). The syrup was purified by flash chromatography: 1-5% MeOH/EtoAc to elute impurities, then 6-12% MeOH/EtoAc (title compound elutes at 10-11%). The desired fractions were concentrated to afford 1.04 g of a brown solid. The solid was then recrystallized (EtoAc) to give the title compound (780 mg, 12% yield) as a tan solid. HNMR ( $\text{CDCl}_3$ )  $\delta$  1.25 (d,  $J=7.5\text{Hz}$ , 3H,  $\text{CH}_3$ ),  $\delta$  1.46 (m, 6H ( $\text{OCH}_2\text{CH}_3$ )<sub>2</sub>), 4H ( $\text{OCH}_2\text{CH}_3$ )<sub>2</sub>,  $\delta$  3.9 (m, 2H, O- $\text{CH}_2\text{P}$ ),  $\delta$  4.04 (m, 1H, H-2'),  $\delta$  4.09-4.39 (m, 2H, H-1'), 7.24 (m, 8H, ( $\text{C}_6\text{H}_4$ )<sub>2</sub>), 7.92 (s, 1H, ( $\text{C}_8\text{-H}$ )), 8.19 (s, 1H,  $\text{C}_2\text{-H}$ ).

Example 9: cHPMPU. cHPMPU was synthesized by adding thionyl chloride (60 mL, 0.812 mmol, 2.02 eq) dropwise to a suspension of disodium HPMPU (131 mg, 0.404 mmol) in *N,N*-dimethylformamide (1.25 mL) at ambient temperature. The resulting light-yellow solution was stirred for 20 min at ambient temperature and then concentrated to dryness (*in vacuo*, 45 °C).  $\text{H}_2\text{O}$  (2 mL) was added and the resulting solution was concentrated to dryness. Methanol (4 mL) was added and the resulting solution was


concentrated to dryness to afford the crude product as a light-yellow solid. Purification by silica flash chromatography (mobile phase: 30% methanol: 70% CH<sub>2</sub>Cl<sub>2</sub> gradient to 50% methanol: 50% CH<sub>2</sub>Cl<sub>2</sub>) afforded pure cHPMPU in 69% yield as a white amorphous solid. <sup>1</sup>H NMR (300 MHz, D<sub>2</sub>O) δ 7.62 d (1H, J = 7.1 Hz, CH=CH), 5.82 d (1H, J = 7.8 Hz, CH=CH), 4.30-3.71 m (7H, CH<sub>2</sub>CH(OCH<sub>2</sub>P)CH<sub>2</sub>OH), NH and OH not observed in D<sub>2</sub>O. <sup>13</sup>C NMR (75 MHz, D<sub>2</sub>O) δ 169.6 s (4-C), 155.1 s (2-C), 150.4 s (6-C), 104.2 s (5-C), 76.71 d (J<sub>P,C</sub> = 3.6 Hz, 2'-CH<sub>2</sub>), 72.30 d (J<sub>P,C</sub> = 6.2 Hz, 3'-CH<sub>2</sub>), 67.90 d (J<sub>P,C</sub> = 142.0 Hz, P-CH<sub>2</sub>), 50.71 s (1'-C). <sup>31</sup>P NMR (121 MHz, D<sub>2</sub>O) δ 9.23 s.

10

**Example 10: cHPMPC ethyl ester.** To a stirred solution of diethyl HPMPC (1.1g) in DMF, NaH (115 mg) was added. After 15 min, the reaction mixture was quenched with acetic acid (1 eq). The solvents were removed under reduced pressure. The crude mixture was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and water. The organic layer was washed with NaCl solution and the crude material obtained was purified on a silica gel column (elution with 5%-10% MeOH in CH<sub>2</sub>Cl<sub>2</sub>) to get cyclic ethyl HPMPC (950 mg) as a diastereomeric mixture (approximately 70%).

20

**Example 11: cHPMPC esters.**




To a stirred suspension of HPMPC (2.79 g) in DMF, thionylchloride (2.1 mL) was added dropwise under anhydrous conditions and the mixture was stirred for 1 hr. In another flask, sodium aryloxide (using the appropriate aryl substituent) was made using the corresponding phenol (8.9 g) and NaH (1.8 g)

in 1:1 DMF/THF (50 mL). This solution was cooled to  $-78^{\circ}\text{C}$  and the chloridate solution was added dropwise under anhydrous conditions. After 2 hrs, the reaction mixture was quenched with acetic acid (5 eq) and the solvents were evaporated under vacuum. The crude mixture was partitioned  
 5 between water and  $\text{CH}_2\text{Cl}_2$ . The organic layer was concentrated and the residue was purified on a silica gel column (elution with 5%-10% MeOH in  $\text{CH}_2\text{Cl}_2$ ) to get the cyclic aryl compound as a single diastereomer in approximately 60% yield. This method is suitable for all substituted or unsubstituted  $\text{R}^{31}$  groups, especially aryl, subject of course to conventional  
 10 protection of labile groups other than amino for which reaction is undesired (amino is protected by reaction with DMF and deprotected with acetic acid and alkanol treatment). This method offers the advantages of producing substantially stereochemically pure product, superior yield and ease of synthesis.

15

Example 12: cHPMPC esters.



To a stirred suspension of cyclic HPMPC (1 mmol) was added N,N'-dicyclohexyl-4-morpholinecarboxamidine (2 mmol) followed by the  
 20 corresponding acyloxymethyl chloride (1.5 mmol). The reaction was stirred for 3 days and the DMF was evaporated under reduced pressure. The crude was purified on a silica gel column (eluted with 5% methanol in methylene chloride) to get the pure cyclic HPMPC derivatives (approximately 30% yield).


The final product was obtained in higher yield by the same reaction  
 25 using cyclic HPMPC (1 mmol), N,N'-dicyclohexyl-4-morpholinecarboxamidine (1.1 mmol) followed by the corresponding acyloxymethyl chloride (1.2 mmol).  $\text{N}^4$ -benzoyl cHPMPC pivaloyloxymethyl ester was

synthesized in a similar manner using N<sup>4</sup>-benzoyl cHPMPC as the starting material.


Example 13: cHPMPC esters.

5 cHPMPC esters were synthesized using appropriate reactants essentially as described in Example 11 for ester moieties corresponding to structure numbers 6, 7, 11, 12, 13, 23, 24, 25 and 26 in Table 5A. cHPMPC esters were synthesized using appropriate reactants essentially as described in Example 12  
10 for ester moieties corresponding to structure numbers 8, 9, 10, 16 and 17 in Table 5A. Melting point data for cHPMPC esters of compound numbers 6, 8, 9, 11, 24, 25 and 26 was as follows: cHPMPC 3-pyridyl ester (#6) - 268-273° C (decomposes); cHPMPC N-ethylmorpholino ester (#7) - 241° C; cHPMPC -CH<sub>2</sub>-O-C(O)-C<sub>6</sub>H<sub>5</sub> ester (#8) - 198-201° C; cHPMPC #9 ortho ester - 176° C; cHPMPC #11 ester - 100-250° C (decomposes); cHPMPC phenyl ester (#12) -  
15 190° C; cHPMPC #24 ester - 218-225° C (waxy liquid); cHPMPC #25 ester - 171° C; cHPMPC #26 ester - 181° C.

Example 14: 9-[2,3-dideoxy-2,3-didehydro-4-phosphonomethoxy-β-D-erythrofuranosyl]adenine esters. Compounds where Z is of structure V and  
20 R<sup>25</sup> and R<sup>29</sup> is oxygen were synthesized by addition-elimination reaction using



, where B was adenine with iodine (2 equivalents) in acetonitrile and a compound having the structure (R<sup>35</sup>O)<sub>2</sub>P(O)-CH<sub>2</sub>-OH (where R<sup>35</sup> was isopropyl, phenyl or 2-ethoxyphenyl) to yield the 3-



25 iodophosphonate diester, which was then eliminated to yield the corresponding structure V compound by reaction with 5 equivalents of sodium methoxide or DBU in anhydrous organic solvent such as methanol or tetrahydrofuran at room temperature for 12 hours.

Corresponding compounds where R<sup>29</sup> is sulfur, are synthesized by the  
30 same method using (R<sup>35</sup>O)<sub>2</sub>P(O)-CH<sub>2</sub>-SH as a reactant. The compound of

structure  $(R^{35}O)_2P(O)-CH_2-OH$  where  $R^{35}$  is isopropyl has been described (Kluge Organic Synthesis (1986) 64:80-83).

Compounds of structure  $(R^{35}O)_2P(O)-CH_2-OH$  where  $R^{35}$  was phenyl or 2-ethoxyphenyl were obtained by reaction of 1 equivalent of  $PCl_3$  with 1 equivalent of t-butanol at 55° C to obtain  $(R^{35}O)_2P(O)H$  (U.S. Patent 3,329,742).  $(R^{35}O)_2P(O)H$  was then silylated using 1 equivalent of bis(trimethylsilyl)-trifluoroacetamide and the resulting  $(R^{35}O)_2P(OTMS)$  was dried under vacuum.  $(R^{35}O)_2P(OTMS)$  was then converted to  $(R^{35}O)_2P(O)-CH_2-OH$  by reaction in paraformaldehyde containing catalytic amounts of titanium isopropoxide (or another lewis acid such as titanium tetrachloride and the like can be used) for 12 hrs (12-16 hours) at 70° C (65 to 75° C). The 2-ethoxyphenyl product was isolated by crystallization. The bis-phenyl product was isolated by silica gel chromatography.

bis(2-ethoxyphenyl) D4AMPI ester:  $^1H$ -NMR (300 MHz,  $CDCl_3$ )  $\delta$  8.38 (s, 1H), 7.97 (s, 1H), 7.21-6.82 (m, 9H), 6.40 (d, 1H,  $J=5.7$  Hz), 6.30 (d, 1H,  $J=5.8$  Hz), 6.16 (s, 1H), 5.61 (s, 2H), 4.48 (dd, 1H,  $J=14$ , 8.8 Hz), 4.38 (dd, 1H,  $J=14$ , 6.5 Hz), 4.10-3.93 (m, 4H), 1.38 (t, 3H,  $J=7.1$  Hz), 1.35 (t, 3H,  $J=7.1$  Hz);  $^{31}P$ -NMR (121 MHz,  $CDCl_3$ )  $\delta$  14.6.

bis(phenyl) D4AMPI ester:  $^1H$ -NMR (300 MHz,  $CDCl_3$ )  $\delta$  8.38 (s, 1H), 7.93 (s, 1H), 7.34 - 7.10 (m, 10H), 7.03 (s, 1H), 6.42 (d, 1H,  $J=5.6$  Hz), 6.34 (d, 1H,  $J=5.6$  Hz), 5.98 (s, 1H), 5.83 (s, 2H), 4.32 (dd, 1H,  $J=14$ , 6.5 Hz), 4.19 (dd, 1H,  $J=14$ , 6.5 Hz);  $^{31}P$ -NMR (121 MHz,  $CDCl_3$ )  $\delta$  13.3.

$(C_6H_4(OC_2H_5)-O)_2P(O)-CH_2-OH$ :  $^1H$ -NMR (300 MHz,  $CDCl_3$ )  $\delta$  7.36-7.16 (m, 10H), 4.19 (dd, 2H,  $J=6.7$ , 5.9 Hz), OH not detected;  $^{31}P$ -NMR (121 MHz,  $CDCl_3$ )  $\delta$  17.0.

$(C_6H_5-O)_2P(O)-CH_2-OH$ :  $^1H$ -NMR (300 MHz,  $CDCl_3$ )  $\delta$  7.25-6.89 (m, 8H), 4.24 (d, 2H,  $J=5.01$  Hz), 4.18-4.08 (m, 4H), 1.46 (t, 6H,  $J=7.0$  Hz);  $^{31}P$ -NMR (121 MHz,  $CDCl_3$ )  $\delta$  19.9.

Example 14: N<sup>4</sup>-benzoyl cHPMPC. The title compound was synthesized using N<sup>4</sup>-benzoyl HPMPC diethyl ester tritylated at the hydroxyl group as a starting material. The starting material was detritylated using acetic acid and then converted to N<sup>4</sup>-benzoyl HPMPC using TMSBr. The resulting compound was converted to N<sup>4</sup>-benzoyl cHPMPC using DCC and morpholine in pyridine. The title compound was tested for activity against


HCMV in tissue culture (NHDF cell line) and was found to be active with an  $IC_{50}$  of 22  $\mu$ M compared with 0.4  $\mu$ M for HPMPC.

$^1H$ NMR (300 MHz,  $CDCl_3$ )  $\delta$  8.02 ( $H_6$ , 1H, d, 7.2 Hz), 7.97 (aromatic, 2H, d, 7.2 Hz), 7.62 (aromatic, 1H, t, 7.2 Hz), 7.5 (aromatic, 2H, t, 7.2 Hz), 7.26 ( $H_5$ , 1H, d, 7.2 Hz), 4.28 (1H, t, 14.7 Hz), 4.15 (1H, t, 10.8 Hz), 4.0 (m, 3H), 3.84(1H,m), 2.49 (1H, d, 14.1 Hz);  $^{31}P$  -NMR (121 MHz,  $CDCl_3$ )  $\delta$  10.07. Melting point 243-246° C.

The claims shall be construed to exclude any subject matter that, at the date of the invention, would not have been patentable under applicable statutory and judicial authority.

What is claimed is:

1. A compound of the formula I







5

or a physiologically acceptable salt thereof, wherein


$L^1$  and  $L^2$  are independently an amino acid or polypeptide residue bonded to the phosphorus atom of the nucleotide analog amidate by an amidate bond, or an oxyester, thioester, a substituted or unsubstituted amine, or hydroxy, provided that one or both of  $L^1$  and  $L^2$  is an amino acid or polypeptide residue and any carboxyl group that is linked by less than about 5 atoms to the amidate N is esterified or amidated and the dotted lines represent facultative bonds;

10  $Z$  is  $-\text{CHR}^7-\text{R}^{11}-(\text{CH}_2)_{m1}-\text{C}^\#(\text{R}^8)((\text{CH}_2)_{m2}(\text{R}^9))-(\text{CH}_2)_{m3}-\text{R}^{10}-(\text{CH}_2)_{m4}-$ ,  $-\text{Q}-\text{C}_6\text{H}_4-\text{CH}_2-$ ,  $-\text{CHR}^7-\text{O}-\text{CHR}^7-\text{O}-\text{CHR}^7-$ ,  $-\text{CHR}^7-(\text{CHR}^{13})_{m1}-\text{CHR}^{14}-\text{R}^{10}-$ ,



20






or VIII

- R<sup>7</sup> is H or C<sub>1</sub>-C<sub>4</sub> alkyl;  
 R<sup>8</sup> is H or C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>2</sub>-C<sub>4</sub> alkenyl, azidomethyl or azidoethyl;  
 R<sup>9</sup> is halogen (F, Cl, Br or I), H or OH;  
 5 R<sup>10</sup> is O, CH<sub>2</sub> or a chemical bond;  
 R<sup>11</sup> is O, S, CH<sub>2</sub>, CHF, CF<sub>2</sub>;  
 Q is -C(R<sup>12</sup>)<sub>2</sub>-CH<sub>2</sub>-, -C(R<sup>12</sup>)<sub>2</sub>-O-, -CR<sup>12</sup>=CR<sup>12</sup>-, or -C≡C-, wherein each R<sup>12</sup>  
 is independently H, or halogen;  
 R<sup>13</sup> is H, halogen, OH, CH<sub>3</sub>, CH<sub>2</sub>OH, or C<sub>3</sub>-C<sub>12</sub> acyloxymethyl;  
 10 R<sup>14</sup> is independently H, halogen, OH, CH<sub>3</sub>, CH<sub>2</sub>OH, C<sub>3</sub>-C<sub>12</sub>  
 acyloxymethyl, or C<sub>2</sub>-C<sub>12</sub> alkoxy;  
 R<sup>25</sup> is CH<sub>2</sub>, CHF or O;  
 R<sup>26</sup> is CH or S, provided that when R<sup>25</sup> is CH, R<sup>26</sup> is not S;  
 R<sup>27</sup> is H, OH, halogen, N<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy or when, R<sup>26</sup> is S,  
 15 R<sup>27</sup> is absent;  
 R<sup>27a</sup> is H, OH, halogen, N<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy;  
 R<sup>28</sup> is H, OH, halogen, N<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl or C<sub>1</sub>-C<sub>4</sub> alkoxy;  
 R<sup>29</sup> is O, S, CH<sub>2</sub>, CHF, CF<sub>2</sub>;  
 R<sup>32</sup> is O;  
 20 m<sub>1</sub> is an integer having a value from 0 to 4;  
 m<sub>2</sub> is an integer having a value from 0 to 4;  
 m<sub>3</sub> is an integer having a value from 0 to 4;  
 m<sub>4</sub> is an integer having a value from 0 to 4;  
 the carbon atom designated C<sup>#</sup> has linked substituents that are in the R,  
 25 S or RS configuration; and  
 B is a heterocyclic base.

30

2. The compound of claim 1 of the formula Ib



and stereoisomers and salts of such compounds wherein


5 X¹ is O or S;

L¹ is an amino acid, a polypeptide, a substituted or unsubstituted amine, an oxyester or a thio ester; and

the carbon atom designated # has linked substituents that are in the *R*, *S* or *RS* configuration, provided that L¹ is not a C₁-C₄ alkyl ester or, when  
 10 when B is cytosin-1-yl, then L¹ is not OCH₂C(O)NR⁵ᵃ₂, OCH₂C(O)OR⁵ᵃ, OCH₂OC(O)R⁵ᵃ, OCH(R⁵ᵃ)OC(O)R⁵ᵃ (*R*, *S* or *RS* stereochemistry), OCH₂C(R⁵ᵃ)₂CH₂OH, OCH₂OR⁵ᵃ, OR⁵ᵃ, NHR⁵ᵃ or NR⁵ᵃ₂ wherein R⁵ᵃ is C₁-C₂₀ alkyl, aryl or aryl-alkyl which may be substituted or unsubstituted by  
 15 substituents independently selected from the group consisting of hydroxy and halogen.

3. The compound of claim 2 wherein L¹ is NHR⁴⁰ wherein R⁴⁰ is C₁-₂₀ alkyl; and OR³¹ wherein R³¹ is

2,3-dihydro-6-hydroxyindene; sesamol; catechol monoester; -CH₂-C(O)-  
 20 N(R⁷)₂ wherein each R⁷ the same or different; -CH₂-S(O)(R⁷); -CH₂-S(O)₂(R⁷); -O-CH₂-CH(OC(O)CH₂R⁷)-CH₂(OC(O)CH₂R⁷); cholesteryl; a monosaccharide; a disaccharide; an oligosaccharide (3 to 9 monosaccharide residues), enolpyruvate; glycerol; an α-D-β-diglyceride; trimethoxybenzyl; triethoxybenzyl; 2-alkyl pyridinyl (C₁-₄ alkyl);







C₃-C₆ aryl substituted by 3, 4 or 5 halogen atoms or 1 or 2 atoms or groups selected from halogen, C₁-C₁₂ alkoxy, cyano, nitro, OH, C₁-C₁₂ haloalkyl, C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl or C₂-C₁₂ alkynyl; or

C<sub>1</sub>-C<sub>4</sub> alkylene-C<sub>3</sub>-C<sub>6</sub> aryl substituted in the aryl moiety by 3 to 5 halogen atoms or 1 to 2 atoms or groups selected from halogen, C<sub>1</sub>-C<sub>12</sub> alkoxy, cyano, nitro, OH, C<sub>1</sub>-C<sub>12</sub> haloalkyl, C<sub>1</sub>-C<sub>12</sub> alkyl, C<sub>2</sub>-C<sub>12</sub> alkenyl or C<sub>2</sub>-C<sub>12</sub> alkynyl.

5

4. The compound of claim 3 wherein B is



10

wherein R<sup>15</sup> is H, OH, F, Cl, Br, I, OR<sup>16</sup>, SH, SR<sup>16</sup>, NH<sub>2</sub>, or NHR<sup>17</sup>;

R<sup>16</sup> is C<sub>1</sub> - C<sub>6</sub> alkyl

R<sup>17</sup> is C<sub>1</sub> - C<sub>6</sub> alkyl;

15

R<sup>18</sup> is N, CF, CCl, CBr, Cl, CR<sup>19</sup> or CSR<sup>19</sup>, COR<sup>19</sup>;

R<sup>19</sup> is H, C<sub>1</sub> - C<sub>9</sub> alkyl, C<sub>2</sub> - C<sub>9</sub> alkenyl, C<sub>2</sub> - C<sub>9</sub> alkynyl or C<sub>7</sub> - C<sub>9</sub> aryl-alkyl unsubstituted or substituted by OH, O, N, F, Cl, Br or I;

R<sup>20</sup> is N or CH;

R<sup>21</sup> is N, CH, CCN, CCF<sub>3</sub>, CC≡CH or CC(O)NH<sub>2</sub>;

$R^{22}$  is H, OH,  $NH_2$ , SH,  $SCH_3$ ,  $SCH_2CH_3$ ,  $SCH_2CCH$ ,  $SCH_2CHCH_2$ ,  $SC_3H_7$ ,  $NH(CH_3)$ ,  $N(CH_3)_2$ ,  $NH(CH_2CH_3)$ ,  $N(CH_2CH_3)_2$ ,  $NH(CH_2CCH)$ ,  $NH(CH_2CHCH_2)$ ,  $NH(C_3H_7)$  or halogen;


$R^{23}$  is H, OH, F, Cl, Br, I,  $SCH_3$ ,  $SCH_2CH_3$ ,  $SCH_2CCH$ ,  $SCH_2CHCH_2$ ,  $SC_3H_7$ ,  $OR^{16}$ ,  $NH_2$ , or  $NHR^{17}$ ; and

$R^{24}$  is O, S or Se.

5. The compound of claim 4 wherein B is cytosin-1-yl, 6-azacytosin-1-yl, 5-fluorocytosin-1-yl, adenin-9-yl, guanin-9-yl or 2, 6-diaminopurin-9-yl.

6. The compound of claim 4 wherein  $R^{31}$  is 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 2-ethoxy-5-hydroxyphenyl, 2-ethoxy-4-hydroxyphenyl, 3,5-dimethoxyphenyl, 2,4-difluorophenyl, 2-(haloalkyl)-phenyl, 3-(haloalkyl)phenyl, 4-(haloalkyl)-phenyl, 2-cyanophenyl, 3-cyanophenyl, 4-cyanophenyl, 2-ethoxyphenyl, 3-ethoxyphenyl, 4-ethoxyphenyl, 2-carboethoxyphenyl, 3-carboethoxyphenyl, 4-carboethoxyphenyl, or 2-haloalkylbenzyl, 3-haloalkylbenzyl or 4-haloalkylbenzyl.

7. The compound of claim 2 of the formula IIa



wherein

25 n is 1, 2, 3, 4 or 5, wherein for  $n > 1$ , each  $-C(R^2)(R^3)$  is the same or different;

$n1$  is an integer;

$R^1$  is H or C<sub>1</sub>-C<sub>9</sub> alkyl which is unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N, COOR<sup>4</sup> and halogen, C<sub>3</sub>-C<sub>6</sub> aryl which is unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N, COOR<sup>4</sup> and halogen or C<sub>3</sub>-C<sub>9</sub> aryl-alkyl which is unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N, COOR<sup>4</sup> and halogen;

$R^2$  is H or C<sub>1</sub>-C<sub>9</sub> alkyl which is unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N, COOR<sup>4</sup> and halogen, C<sub>3</sub>-C<sub>6</sub> aryl which is unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N, COOR<sup>4</sup> and halogen or C<sub>3</sub>-C<sub>9</sub> aryl-alkyl which is unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N, COOR<sup>4</sup> and halogen;

$R^3$  is C(O)-OR<sup>4</sup>, amino, C<sub>1</sub> - C<sub>3</sub> alkylamino, C<sub>1</sub> - C<sub>3</sub> alkylidiamino, C<sub>1</sub> - C<sub>6</sub> alkenylamino, hydroxy, thiol, C<sub>1</sub> - C<sub>3</sub> alkoxy, C<sub>1</sub> - C<sub>3</sub> alkthiol, (CH<sub>2</sub>)<sub>n</sub>COOR<sup>4</sup>, C<sub>1</sub>-C<sub>6</sub> alkyl which is unsubstituted or substituted with OH, halogen, SH, NH<sub>2</sub>, phenyl, hydroxyphenyl or C<sub>7</sub> - C<sub>10</sub> alkoxyphenyl; C<sub>2</sub>-C<sub>6</sub> alkenyl which is unsubstituted or substituted with OH, halogen, SH, NH<sub>2</sub>, phenyl, hydroxyphenyl or C<sub>7</sub> - C<sub>10</sub> alkoxyphenyl; C<sub>6</sub>-C<sub>12</sub> aryl which is unsubstituted or substituted with OH, halogen, SH, NH<sub>2</sub>, phenyl, hydroxyphenyl or C<sub>7</sub> - C<sub>10</sub> alkoxyphenyl; and

$R^4$  is H provided that n<sub>1</sub> greater than 1, or is C<sub>3</sub>-C<sub>9</sub> alkyl which is substituted by substituents independently selected from the group consisting of OH, O, N and halogen, C<sub>3</sub>-C<sub>6</sub> aryl which is substituted by substituents independently selected from the group consisting of OH, O, N and halogen or C<sub>3</sub>-C<sub>9</sub> aryl-alkyl which is substituted by substituents independently selected from the group consisting of OH, O, N and halogen.

8. The compound of claim 7 wherein  
n and n<sub>1</sub> are 1;

$R^1$  is H, methyl, phenyl or benzyl;

$R^2$  is H;

$R^3$  is H, -CH<sub>3</sub>, -CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>-CH(CH<sub>3</sub>)<sub>2</sub>, -CHCH<sub>3</sub>-CH<sub>2</sub>-CH<sub>3</sub>, -CH<sub>2</sub>-C<sub>6</sub>H<sub>5</sub>, -CH<sub>2</sub>CH<sub>2</sub>-S-CH<sub>3</sub>, -CH<sub>2</sub>OH, -CH(OH)-CH<sub>3</sub>, -CH<sub>2</sub>-SH, -CH<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>OH, -CH<sub>2</sub>-

CO-NH<sub>2</sub>, -CH<sub>2</sub>-CH<sub>2</sub>-CO-NH<sub>2</sub>, -CH<sub>2</sub>-COOH, -CH<sub>2</sub>-CH<sub>2</sub>-COOH, -(CH<sub>2</sub>)<sub>4</sub>-NH<sub>2</sub>, -(CH<sub>2</sub>)<sub>3</sub>-NH-C(NH<sub>2</sub>)-NH<sub>2</sub>, 1-guanidinoprop-3-yl, benzyl, 4-hydroxybenzyl, imidazol-4-yl, indol-3-yl, methoxyphenyl or ethoxyphenyl; and


- 5 R<sup>4</sup> is methyl, ethyl, propyl, isopropyl, butyl, t-butyl, phenyl, benzyl, 1-pyridyl, 3-pyridyl, 1-pyrimidinyl, pivaloyloxymethyl, N-ethylmorpholino, N-2-propylmorpholino, methoxyethyl, 4-N-methylpiperidyl, 3-N-methylpiperidyl, 2-, 3-, or 4-N,N-dimethylaminophenyl, 2-, 3-, or 4-N,N-diethylaminophenyl or 1-ethylpiperazinyl.

- 10 9. The compound of claim 8 wherein B is cytosin-1-yl, 6-azacytosin-1-yl, adenin-9-yl, guanin-9-yl or 2, 6-diaminopurin-9-yl, and X<sup>1</sup> is O.

- 10 10. The compound of claim 1 of the formula Id



11. The compound of claim 10 wherein L<sup>1</sup> is of the formula III



wherein

- 20 n is 1, 2, 3, 4 or 5, wherein for n > 1, each -C(R<sup>2</sup>)(R<sup>3</sup>) is the same or different;

n1 is an integer;

- 25 R<sup>1</sup> is H or C<sub>1</sub>-C<sub>9</sub> alkyl which is unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N, COOR<sup>4</sup> and halogen, C<sub>3</sub>-C<sub>6</sub> aryl which is unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N,

COOR<sup>4</sup> and halogen or C<sub>3</sub>-C<sub>9</sub> aryl-alkyl which is unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N, COOR<sup>4</sup> and halogen;

5 R<sup>2</sup> is H or C<sub>1</sub>-C<sub>9</sub> alkyl which is unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N, COOR<sup>4</sup> and halogen, C<sub>3</sub>-C<sub>6</sub> aryl which is unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N, COOR<sup>4</sup> and halogen or C<sub>3</sub>-C<sub>9</sub> aryl-alkyl which is unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N, COOR<sup>4</sup> and halogen;

10 R<sup>3</sup> is C(O)-OR<sup>4</sup>, amino, C<sub>1</sub> - C<sub>3</sub> alkylamino, C<sub>1</sub> - C<sub>3</sub> alkylidiamino, C<sub>1</sub> - C<sub>6</sub> alkenylamino, hydroxy, thiol, C<sub>1</sub> - C<sub>3</sub> alkoxy, C<sub>1</sub> - C<sub>3</sub> alkthiol, (CH<sub>2</sub>)<sub>n</sub>COOR<sup>4</sup>, C<sub>1</sub>-C<sub>6</sub> alkyl which is unsubstituted or substituted with OH, halogen, SH, NH<sub>2</sub>, phenyl, hydroxyphenyl or C<sub>7</sub> - C<sub>10</sub> alkoxyphenyl; C<sub>2</sub>-C<sub>6</sub> alkenyl which is  
15 unsubstituted or substituted with OH, halogen, SH, NH<sub>2</sub>, phenyl, hydroxyphenyl or C<sub>7</sub> - C<sub>10</sub> alkoxyphenyl; C<sub>6</sub>-C<sub>12</sub> aryl which is unsubstituted or substituted with OH, halogen, SH, NH<sub>2</sub>, phenyl, hydroxyphenyl or C<sub>7</sub> - C<sub>10</sub> alkoxyphenyl; and

20 R<sup>4</sup> is H provided that n<sub>1</sub> greater than 1, or is C<sub>3</sub>-C<sub>9</sub> alkyl which is substituted by substituents independently selected from the group consisting of OH, O, N and halogen, C<sub>3</sub>-C<sub>6</sub> aryl which is substituted by substituents independently selected from the group consisting of OH, O, N and halogen or C<sub>3</sub>-C<sub>9</sub> aryl-alkyl which is substituted by substituents independently selected from the group consisting of OH, O, N and halogen;

25 L<sup>2</sup> is OR, SR or is the same as L<sup>1</sup> wherein,  
R is H,

30 C<sub>3</sub>-C<sub>24</sub> acyloxyalkyl,  
C<sub>6</sub>-C<sub>24</sub> acyloxyarylalkyl,  
C<sub>3</sub>-C<sub>24</sub> acyloxyalkoxyalkyl,  
C<sub>3</sub>-C<sub>24</sub> acyloxyhaloalkyl,  
C<sub>1</sub>-C<sub>20</sub> alkyl which is unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N and halogen (F, Cl, Br, I),

35 C<sub>3</sub>-C<sub>20</sub> aryl which is unsubstituted or substituted by substituents independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy,

C<sub>1</sub>-C<sub>6</sub> haloalkyl (1 to 3 halogen atoms), cyano, nitro, OH, O, N and halogen, or  
 C<sub>4</sub>-C<sub>20</sub> aryl-alkyl which is unsubstituted or substituted in the aryl  
 moiety by substituents independently selected from the group consisting of  
 C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>1</sub>-C<sub>6</sub> haloalkyl (1 to 3 halogen atoms), cyano, nitro,  
 5 OH, O, N and halogen.

12. The compound of claim 11 wherein  
 n and n1 are 1;

R is N-ethylmorpholino, pivaloyloxymethyl, phenyl, benzyl, isopropyl,  
 10 t-butyl, ethyl, isopropyl, butyl, adamantoyloxymethyl, 3-methoxyphenyl, 2-  
 carboethoxyphenyl, 4-fluorophenyl, 2,4-difluorophenyl, 3,5-dimethoxyphenyl,  
 2,4-dichlorophenyl, 2-ethoxyphenyl, 3-dimethylaminophenyl, 4-  
 trifluoromethylbenzyl, 2-ethylsalicyl, -O-CH<sub>2</sub>-O-C(O)-C<sub>10</sub>H<sub>15</sub>, -C<sub>6</sub>H<sub>4</sub>-CH<sub>2</sub>-  
 N(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>-CH<sub>2</sub>F, -CH<sub>2</sub>-CH<sub>2</sub>Cl, -CH<sub>2</sub>-CF<sub>3</sub>, -CH<sub>2</sub>-CCl<sub>3</sub>, R<sup>5</sup>, NHR<sup>6</sup> or N(R<sup>6</sup>)<sub>2</sub>  
 15 wherein,

R<sup>5</sup> is CH<sub>2</sub>C(O)N(R<sup>6</sup>)<sub>2</sub>, CH<sub>2</sub>C(O)OR<sup>6</sup>, CH<sub>2</sub>OC(O)R<sup>6</sup>,  
 CH(R<sup>6</sup>)OC(O)R<sup>6</sup>, CH<sub>2</sub>C(R<sup>6</sup>)<sub>2</sub>CH<sub>2</sub>OH, or CH<sub>2</sub>OR<sup>6</sup>, and

R<sup>6</sup> is C<sub>1</sub>-C<sub>20</sub> alkyl which is unsubstituted or substituted by  
 substituents independently selected from the group consisting of  
 20 OH, O, N and halogen (1 to 5 halogen atoms), C<sub>6</sub>-C<sub>20</sub> aryl which  
 is unsubstituted or substituted by substituents independently  
 selected from the group consisting of OH, O, N and halogen (1 to  
 5 halogen atoms) or C<sub>7</sub>-C<sub>20</sub> aryl-alkyl which is unsubstituted or  
 substituted by substituents independently selected from the  
 25 group consisting of OH, O, N and halogen (1 to 5 halogen atoms);

R<sup>1</sup> is H, methyl, ethyl, isopropyl, phenyl or benzyl;

R<sup>2</sup> is H;


R<sup>3</sup> is H, -CH<sub>3</sub>, -CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>-CH(CH<sub>3</sub>)<sub>2</sub>, -CHCH<sub>3</sub>-CH<sub>2</sub>-CH<sub>3</sub>, -CH<sub>2</sub>-  
 C<sub>6</sub>H<sub>5</sub>, -CH<sub>2</sub>CH<sub>2</sub>-S-CH<sub>3</sub>, -CH<sub>2</sub>OH, -CH(OH)-CH<sub>3</sub>, -CH<sub>2</sub>-SH, -CH<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>OH, -CH<sub>2</sub>-  
 30 CO-NH<sub>2</sub>, -CH<sub>2</sub>-CH<sub>2</sub>-CO-NH<sub>2</sub>, -CH<sub>2</sub>-COOH, -CH<sub>2</sub>-CH<sub>2</sub>-COOH, -(CH<sub>2</sub>)<sub>4</sub>-NH<sub>2</sub>,  
 -(CH<sub>2</sub>)<sub>3</sub>-NH-C(NH<sub>2</sub>)-NH<sub>2</sub>, 1-guanidinoprop-3-yl, benzyl, 4-hydroxybenzyl,  
 imidazol-4-yl, indol-3-yl, methoxyphenyl or ethoxyphenyl; and

R<sup>4</sup> is methyl, ethyl, propyl, isopropyl, butyl, t-butyl, phenyl, benzyl, 1-  
 pyridyl, 3-pyridyl, 1-pyrimidinyl, pivaloyloxymethyl, N-ethylmorpholino, N-  
 35 2-propylmorpholino, methoxyethyl, 4-N-methylpiperidyl, 3-N-




methylnpiperidyl, 2-, 3-, and 4-N,N-dimethylaminophenyl and 2-, 3-, and 4-N,N-diethylaminophenyl or 1-ethylpiperazinyl.

13. The compound of claim 12 wherein Z is  $-\text{CHR}^7-\text{R}^{11}-(\text{CH}_2)_{m1}-$   
 5  $\text{C}(\text{R}^8)((\text{CH}_2)_{m2}(\text{R}^9))-(\text{CH}_2)_{m3}-\text{R}^{10}-(\text{CH}_2)_{m4}-$



10 14. The compound of claim 13 wherein  
Z is  $-\text{CH}_2-\text{O}-\text{CH}_2-\text{CH}_2-$ ,  $-\text{CH}_2-\text{O}-\text{CH}_2-\text{CH}(\text{CH}_2\text{OH})-$ ,  $-\text{CH}_2-\text{O}-\text{CH}_2-$   
 $\text{CH}(\text{CH}_2\text{F})-$ ,  $-\text{CH}_2-\text{O}-\text{CH}_2-\text{CH}(\text{CH}_3)-$ ,  $-\text{CH}_2-\text{O}-\text{CH}_2-\text{CH}(\text{CH}=\text{CH}_2)-$  or  $-\text{CH}_2-\text{O}-$   
 $\text{CH}_2-\text{CH}(\text{CH}_2\text{N}_3)-$ , or is of formula IV or V



wherein


R<sup>25</sup> and R<sup>29</sup> are O;

R<sup>26</sup> is CH<sub>3</sub>;

20 R<sup>27</sup> and R<sup>28</sup> are H; and


B is adenin-9-yl, 1-deazaadenin-9-yl, 3-deazaadenin-9-yl, 7-deaza-8-azaadenin-9-yl, 8-azaadenin-9-yl, guanin-9-yl, 2, 6-diaminopurin-9-yl, 2-aminopurin-9-yl, thymine-1-yl, cytosine-1-yl, 5-fluorocytosine-1-yl, 6-azacytosine-1-yl, 5-methylcytosine-1-yl, 5-bromovinyluracil-1-yl, 5-fluorouracil-1-yl or 5-trifluoromethyluracil-1-yl.

15. The compound of claim 1 of the formula Ic




16. The compound of claim 15 of the formula  $\text{IIc}$

5



wherein  $L^2$  is OR, SR or



10

wherein

R is H,

C<sub>3</sub>-C<sub>24</sub> acyloxyalkyl,

C<sub>6</sub>-C<sub>24</sub> acyloxyarylalkyl,

- C<sub>3</sub>-C<sub>24</sub> acyloxyalkoxyalkyl,  
 C<sub>3</sub>-C<sub>24</sub> acyloxyhaloalkyl,  
 C<sub>1</sub>-C<sub>20</sub> alkyl which is unsubstituted or substituted by substituents  
 independently selected from the group consisting of OH, O, N and halogen (F,  
 5 Cl, Br, I),  
 C<sub>3</sub>-C<sub>20</sub> aryl which is unsubstituted or substituted by substituents  
 independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy,  
 C<sub>1</sub>-C<sub>6</sub> haloalkyl (1 to 3 halogen atoms), cyano, nitro, OH, O, N and halogen, or  
 C<sub>4</sub>-C<sub>20</sub> aryl-alkyl which is unsubstituted or substituted in the aryl  
 10 moiety by substituents independently selected from the group consisting of  
 C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>1</sub>-C<sub>6</sub> haloalkyl (1 to 3 halogen atoms), cyano, nitro,  
 OH, O, N and halogen; and  
 R<sup>1</sup> is O-C<sub>6</sub>H<sub>4</sub>-CH<sub>2</sub>-N(CH<sub>3</sub>)<sub>2</sub>, OR<sup>5</sup>, NHR<sup>6</sup> or N(R<sup>6</sup>)<sub>2</sub> wherein R<sup>5</sup> is  
 CH<sub>2</sub>C(O)N(R<sup>6</sup>)<sub>2</sub>, CH<sub>2</sub>C(O)OR<sup>6</sup>, CH<sub>2</sub>OC(O)R<sup>6</sup>, CH(R<sup>6</sup>)OC(O)R<sup>6</sup>,  
 15 CH<sub>2</sub>C(R<sup>6</sup>)<sub>2</sub>CH<sub>2</sub>OH, or CH<sub>2</sub>OR<sup>6</sup>, and wherein R<sup>6</sup> is C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>6</sub>-C<sub>20</sub> aryl or  
 C<sub>7</sub>-C<sub>20</sub> aryl-alkyl which is unsubstituted or substituted by substituents  
 independently selected from the group consisting of OH, O, N and halogen.

17. A compound of the formula (OR<sup>31</sup>)<sub>2</sub>P(O)-Z<sup>1</sup>-B or  
 20 (OR)(OR<sup>31</sup>)P(O)-Z<sup>1</sup>-B, wherein;

B is a heterocyclic base;


Z<sup>1</sup> is selected from the group consisting of -CH<sub>2</sub>-O-CH<sub>2</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-O-  
 C#H(CH<sub>2</sub>OH)-CH<sub>2</sub>-, -CH<sub>2</sub>-O-C#H(CH<sub>3</sub>)-CH<sub>2</sub>-, -CH<sub>2</sub>-O-C#H(CH<sub>2</sub>F)-CH<sub>2</sub>-, -CH<sub>2</sub>-  
 O-C#H(CH=CH<sub>2</sub>)-CH<sub>2</sub>- and -CH<sub>2</sub>-O-C#H(CH<sub>2</sub>N<sub>3</sub>)-CH<sub>2</sub>-;

- 25 R is H,

- C<sub>3</sub>-C<sub>24</sub> 1-acyloxy-1-alkyl,  
 C<sub>6</sub>-C<sub>24</sub> 1-acyloxy-1-aryl-1-alkyl,  
 C<sub>3</sub>-C<sub>24</sub> 1-acyloxy-2-alkoxy-1-alkyl,  
 C<sub>3</sub>-C<sub>24</sub> 1-acyloxy-2-halo-1-alkyl,  
 30 C<sub>1</sub>-C<sub>20</sub> alkyl which is unsubstituted or substituted by substituents  
 independently selected from the group consisting of OH, O, N and halogen (F,  
 Cl, Br, I),  
 C<sub>3</sub>-C<sub>20</sub> aryl which is unsubstituted or substituted by substituents  
 independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy,  
 35 C<sub>1</sub>-C<sub>6</sub> haloalkyl, cyano, nitro, OH, O, N and halogen, or

C<sub>4</sub>-C<sub>20</sub> aryl-alkyl which is unsubstituted or substituted in the aryl moiety by substituents independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>1</sub>-C<sub>6</sub> haloalkyl, cyano, nitro, OH, O, N and halogen;

- R<sup>31</sup> is 2,3-dihydro-6-hydroxyindene; sesamol; catechol monoester;  
 5 -CH<sub>2</sub>-C(O)-N(R<sup>7</sup>)<sub>2</sub> wherein each R<sup>7</sup> is hydrogen or C<sub>1-4</sub> alkyl and is the same or different; -CH<sub>2</sub>-S(O)(R<sup>7</sup>); -CH<sub>2</sub>-S(O)<sub>2</sub>(R<sup>7</sup>); -O-CH<sub>2</sub>-CH(OC(O)CH<sub>2</sub>R<sup>7</sup>)-CH<sub>2</sub>(OC(O)CH<sub>2</sub>R<sup>7</sup>); cholesteryl; a monosaccharide; a disaccharide; an oligosaccharide (3 to 9 monosaccharide residues), enolpyruvate; glycerol; an α-D-β-diglyceride; trimethoxybenzyl; triethoxybenzyl; 2-alkyl pyridinyl (C<sub>1-4</sub> alkyl);  
 10




C<sub>3</sub>-C<sub>6</sub> aryl substituted by 3, 4 or 5 halogen atoms or 1 or 2 atoms or groups selected from halogen, C<sub>1</sub>-C<sub>12</sub> alkoxy, cyano, nitro, OH, C<sub>1</sub>-C<sub>12</sub> haloalkyl, C<sub>1</sub>-C<sub>12</sub> alkyl, C<sub>2</sub>-C<sub>12</sub> alkenyl or C<sub>2</sub>-C<sub>12</sub> alkynyl; or

- 15 C<sub>1</sub>-C<sub>4</sub> alkylene-C<sub>3</sub>-C<sub>6</sub> aryl substituted in the aryl moiety by 3 to 5 halogen atoms or 1 to 2 atoms or groups selected from halogen, C<sub>1</sub>-C<sub>12</sub> alkoxy, cyano, nitro, OH, C<sub>1</sub>-C<sub>12</sub> haloalkyl, C<sub>1</sub>-C<sub>12</sub> alkyl, C<sub>2</sub>-C<sub>12</sub> alkenyl or C<sub>2</sub>-C<sub>12</sub> alkynyl, provided that when Z<sup>1</sup> is -CH<sub>2</sub>-O-CH<sub>2</sub>-CH<sub>2</sub>- and B is adenin-9-yl, both R<sup>31</sup> are not 4-nitrobenzyl or 4-trifluoromethyl-benzyl.  
 20

18. The compound of claim 17 wherein R<sup>31</sup> is 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 2-ethoxy-5-hydroxyphenyl, 2-ethoxy-4-hydroxyphenyl, 3,5-dimethoxyphenyl, 2,4-difluorophenyl, 2-(haloalkyl)-phenyl, 3-(haloalkyl)phenyl, 4-(haloalkyl)-phenyl, 2-cyanophenyl, 3-cyanophenyl, 4-cyanophenyl, 2-ethoxyphenyl, 2-carboethoxyphenyl, 3-carboethoxyphenyl, 4-carboethoxyphenyl, or 2-haloalkylbenzyl, 3-haloalkylbenzyl or 4-haloalkylbenzyl.  
 25

19. The compound of claim 18 wherein B is cytosin-1-yl, 6-azacytosin-1-yl, 5-fluorocytosin-1-yl, adenin-9-yl, guanin-9-yl or 2, 6-diaminopurin-9-yl.  
 30

20. A compound of the formula  $(L^1O)_2P(O)-Z-B^1$  or



5




wherein

substituents linked to the carbon atom designated # are in the *R*, *S* or *RS* configuration;

$L^1$  is independently an amino acid, a polypeptide, an oxyester, a  
10 thioester or a substituted or unsubstituted amine;

$B^1$  is a protected heterocyclic base; and

$Z-B^1$  is



15

wherein

$R^{27}$  is H, OH, halogen,  $N_3$ ,  $C_1-C_4$  alkyl,  $C_1-C_4$  alkoxy or when,  $R^{26}$   
is S,  $R^{27}$  is absent;

$R^{28}$  is H, OH, halogen,  $N_3$ ,  $C_1-C_4$  alkyl or  $C_1-C_4$  alkoxy;

20


$R^{29}$  is O, S,  $CH_2$ , CHF or  $CF_2$ ;

$R^{33}$  is H, OH, TBSO, halogen, cyano,  $CH_2N_3$ ,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  alkoxy,  $CH_2OH$  or azido; and

$R^{34}$  is H,  $CH_2CN$  or  $CF_3$ , with the proviso that, for structure XXX, when  $R^{25}$  is O or  $CH_2$  and  $R^{29}$   $CH_2$  or O,  $R^{35}$  is not H or  $C_1$ - $C_6$  alkyl.

5

21. The compound of claim 20 wherein  $B^1$  is



10

wherein

$R^{18}$  is N, CF, CCl, CBr, Cl,  $CR^{19}$  or  $CSR^{19}$ ,  $COR^{19}$ ;

$R^{20}$  is N or CH;

15

$R^{21}$  is N, CH, CCN,  $CCF_3$ ,  $CC\equiv CH$  or  $CC(O)NH_2$ ;

$R^{22A}$  is  $R^{39}$  or  $R^{22}$  provided that  $R^{22}$  is not  $NH_2$ ;

$R^{22}$  is H, OH,  $NH_2$ , SH,  $SCH_3$ ,  $SCH_2CH_3$ ,  $SCH_2CCH$ ,  $SCH_2CHCH_2$ ,  $SC_3H_7$ ,  $NH(CH_3)$ ,  $N(CH_3)_2$ ,  $NH(CH_2CH_3)$ ,  $N(CH_2CH_3)_2$ ,  $NH(CH_2CCH)$ ,  $NH(CH_2CHCH_2)$ ,  $NH(C_3H_7)$  or halogen (F, Cl, Br or I);

20

$R^{23A}$  is  $R^{39}$  or  $R^{23}$  provided that  $R^{23}$  is not  $NH_2$ ;

$R^{23}$  is H, OH, F, Cl, Br, I,  $SCH_3$ ,  $SCH_2CH_3$ ,  $SCH_2CCH$ ,  $SCH_2CHCH_2$ ,  $SC_3H_7$ ,  $OR^{16}$ ,  $NH_2$ , or  $NHR^{17}$ ;

$R^{24}$  is O, S or Se; and

$R^{39}$  is  $NHR^{40}$ ,  $NHC(O)R^{36}$  or  $NCR^{41}N(R^{38})_2$  wherein,

$R^{36}$  is  $C_1$ - $C_{19}$  alkyl,  $C_1$ - $C_{19}$  alkenyl,  $C_3$ - $C_{10}$  aryl, adamantoyl, alkylanyl, or  $C_3$ - $C_{10}$  aryl substituted with 1 or 2 atoms or groups selected from halogen, methyl, ethyl, methoxy, ethoxy, hydroxy and cyano;

5  $R^{38}$  is  $C_1$ - $C_{10}$  alkyl, or both  $R^{38}$  together are 1-morpholino, 1-piperidine or 1-pyrrolidine;

$R^{40}$  is  $C_1$ - $C_{20}$  alkyl; and

$R^{41}$  is hydrogen or  $CH_3$ .

10 22. The compound of claim 21 wherein  $L^1$  is R or  $R^{31}$  wherein R is

$C_3$ - $C_{24}$  1-acyloxy-1-alkyl,

$C_6$ - $C_{24}$  1-acyloxy-1-aryl-1-alkyl,

$C_3$ - $C_{24}$  1-acyloxy-2-alkoxy-1-alkyl,

15  $C_3$ - $C_{24}$  1-acyloxy-2-halo-1-alkyl,

$C_1$ - $C_{20}$  alkyl which is unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N and halogen (F, Cl, Br, I),

$C_3$ - $C_{20}$  aryl which is unsubstituted or substituted by substituents independently selected from the group consisting of  $C_1$ - $C_6$  alkyl,  $C_1$ - $C_6$  alkoxy,  $C_1$ - $C_6$  haloalkyl (1 to 3 halogen atoms), cyano, nitro, OH, O, N and halogen,


$C_4$ - $C_{20}$  aryl-alkyl which is unsubstituted or substituted in the aryl moiety by substituents independently selected from the group consisting of  $C_1$ - $C_6$  alkyl,  $C_1$ - $C_6$  alkoxy,  $C_1$ - $C_6$  haloalkyl (1 to 3 halogen atoms), cyano, nitro, OH, O, N and halogen,

$C_3$ - $C_6$  aryl substituted by 3 to 5 halogen atoms or 1 to 2 atoms or groups independently selected from the group consisting of halogen,  $C_1$ - $C_{12}$  alkoxy, cyano, nitro, hydroxy,  $C_1$ - $C_{12}$  haloalkyl,  $C_1$ - $C_{12}$  alkyl,  $C_2$ - $C_{12}$  alkenyl or  $C_2$ - $C_{12}$  alkynyl, or

30  $C_1$ - $C_4$  alkylene- $C_3$ - $C_6$  aryl substituted in the aryl moiety by 3 to 5 halogen atoms or 1 to 2 atoms or groups independently selected from the group consisting of halogen,  $C_1$ - $C_{12}$  alkoxy, cyano, nitro, hydroxy,  $C_1$ - $C_{12}$  haloalkyl,  $C_1$ - $C_{12}$  alkyl,  $C_2$ - $C_{12}$  alkenyl or  $C_2$ - $C_{12}$  alkynyl; and

$R^{31}$  is 2,3-dihydro-6-hydroxyindene; sesamol; catechol monoester;  
35  $-CH_2-C(O)-N(R^7)_2$  wherein each  $R^7$  the same or different;  $-CH_2-S(O)(R^7)$ ;  $-CH_2-$

$S(O)_2(R^7)$ ;  $-O-CH_2-CH(OC(O)CH_2R^7)-CH_2(OC(O)CH_2R^7)$ ; cholesteryl; a monosaccharide; a disaccharide; an oligosaccharide (3 to 9 monosaccharide residues), enolpyruvate; glycerol; an  $\alpha$ -D- $\beta$ -diglyceride; trimethoxybenzyl; triethoxybenzyl; 2-alkyl pyridinyl ( $C_{1-4}$  alkyl);



$C_3-C_6$  aryl substituted by 3, 4 or 5 halogen atoms or 1 or 2 atoms or groups selected from halogen,  $C_1-C_{12}$  alkoxy, cyano, nitro, OH,  $C_1-C_{12}$  haloalkyl,  $C_1-C_{12}$  alkyl,  $C_2-C_{12}$  alkenyl or  $C_2-C_{12}$  alkynyl; or

10  $C_1-C_4$  alkylene- $C_3-C_6$  aryl substituted in the aryl moiety by 3 to 5 halogen atoms or 1 to 2 atoms or groups selected from halogen,  $C_1-C_{12}$  alkoxy, cyano, nitro, OH,  $C_1-C_{12}$  haloalkyl,  $C_1-C_{12}$  alkyl,  $C_2-C_{12}$  alkenyl or  $C_2-C_{12}$  alkynyl.

15 23. The compound of claim 21 wherein  $L^1$  is ethylglycine or N-methylglycine.

24. A compound of the formula  $(OR^{35})(OR^{35})P(O)-Z-B$ , wherein; B is a heterocyclic base;

$R^{35}$  is independently R or  $R^{31}$ , wherein R is independently

20 H,

$C_3-C_{24}$  1-acyloxy-1-alkyl,

$C_6-C_{24}$  1-acyloxy-1-aryl-1-alkyl,

$C_3-C_{24}$  1-acyloxy-2-alkoxy-1-alkyl,

$C_3-C_{24}$  1-acyloxy-2-halo-1-alkyl,

25  $C_1-C_{20}$  alkyl which is unsubstituted or substituted by substituents independently selected from the group consisting of OH, O, N and halogen (F, Cl, Br, I),

$C_3-C_{20}$  aryl which is unsubstituted or substituted by substituents independently selected from the group consisting of  $C_1-C_6$  alkyl,  $C_1-C_6$  alkoxy,  $C_1-C_6$  haloalkyl (1 to 3 halogen atoms), cyano, nitro, OH, O, N and halogen,

30  $C_4-C_{20}$  aryl-alkyl which is unsubstituted or substituted in the aryl moiety by substituents independently selected from the group consisting of




C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>1</sub>-C<sub>6</sub> haloalkyl (1 to 3 halogen atoms), cyano, nitro, OH, O, N and halogen,

5 C<sub>3</sub>-C<sub>6</sub> aryl substituted by 3 to 5 halogen atoms or 1 to 2 atoms or groups independently selected from the group consisting of halogen, C<sub>1</sub>-C<sub>12</sub> alkoxy, cyano, nitro, hydroxy, C<sub>1</sub>-C<sub>12</sub> haloalkyl, C<sub>1</sub>-C<sub>12</sub> alkyl, C<sub>2</sub>-C<sub>12</sub> alkenyl or C<sub>2</sub>-C<sub>12</sub> alkynyl, or

10 C<sub>1</sub>-C<sub>4</sub> alkylene-C<sub>3</sub>-C<sub>6</sub> aryl substituted in the aryl moiety by 3 to 5 halogen atoms or 1 to 2 atoms or groups independently selected from the group consisting of halogen, C<sub>1</sub>-C<sub>12</sub> alkoxy, cyano, nitro, hydroxy, C<sub>1</sub>-C<sub>12</sub> haloalkyl, C<sub>1</sub>-C<sub>12</sub> alkyl, C<sub>2</sub>-C<sub>12</sub> alkenyl or C<sub>2</sub>-C<sub>12</sub> alkynyl;


R<sup>31</sup> is 2,3-dihydro-6-hydroxyindene; sesamol; catechol monoester; -CH<sub>2</sub>-C(O)-N(R<sup>7</sup>)<sub>2</sub> wherein each R<sup>7</sup> the same or different; -CH<sub>2</sub>-S(O)(R<sup>7</sup>); -CH<sub>2</sub>-S(O)<sub>2</sub>(R<sup>7</sup>); -O-CH<sub>2</sub>-CH(OC(O)CH<sub>2</sub>R<sup>7</sup>)-CH<sub>2</sub>(OC(O)CH<sub>2</sub>R<sup>7</sup>); cholesteryl; a  
15 monosaccharide; a disaccharide; an oligosaccharide (3 to 9 monosaccharide residues), enolpyruvate; glycerol; an α-D-β-diglyceride; trimethoxybenzyl; triethoxybenzyl; 2-alkyl pyridinyl (C<sub>1-4</sub> alkyl);



20 C<sub>3</sub>-C<sub>6</sub> aryl substituted by 3, 4 or 5 halogen atoms or 1 or 2 atoms or groups selected from halogen, C<sub>1</sub>-C<sub>12</sub> alkoxy, cyano, nitro, OH, C<sub>1</sub>-C<sub>12</sub> haloalkyl, C<sub>1</sub>-C<sub>12</sub> alkyl, C<sub>2</sub>-C<sub>12</sub> alkenyl or C<sub>2</sub>-C<sub>12</sub> alkynyl; or

C<sub>1</sub>-C<sub>4</sub> alkylene-C<sub>3</sub>-C<sub>6</sub> aryl substituted in the aryl moiety by 3 to 5 halogen atoms or 1 to 2 atoms or groups selected from halogen, C<sub>1</sub>-C<sub>12</sub> alkoxy, cyano, nitro, OH, C<sub>1</sub>-C<sub>12</sub> haloalkyl, C<sub>1</sub>-C<sub>12</sub> alkyl, C<sub>2</sub>-C<sub>12</sub> alkenyl or C<sub>2</sub>-C<sub>12</sub> alkynyl;

25 Z-B is selected from the group consisting of



5        wherein

substituents linked to the carbon atom designated # are in the *R*, *S* or *RS* configuration,

$R^{25}$  is  $\text{CH}_2$ ,  $\text{CHF}$  or  $\text{O}$ ;

$R^{26}$  is  $\text{CH}$  or  $\text{S}$ , provided that when  $R^{25}$  is  $\text{CH}$ ,  $R^{26}$  is not  $\text{S}$ ;

10         $R^{27}$  is  $\text{H}$ ,  $\text{OH}$ , halogen,  $\text{N}_3$ ,  $\text{C}_1\text{-C}_4$  alkyl,  $\text{C}_1\text{-C}_4$  alkoxy or when,  $R^{26}$  is  $\text{S}$ ,  $R^{27}$  is absent;

$R^{28}$  is  $\text{H}$ ,  $\text{OH}$ , halogen,  $\text{N}_3$ ,  $\text{C}_1\text{-C}_4$  alkyl or  $\text{C}_1\text{-C}_4$  alkoxy;

$R^{29}$  is  $\text{O}$ ,  $\text{S}$ ,  $\text{CH}_2$ ,  $\text{CHF}$  or  $\text{CF}_2$ ;

15         $R^{33}$  is  $\text{H}$ ,  $\text{OH}$ ,  $\text{TBSO}$ , halogen, cyano,  $\text{CH}_2\text{N}_3$ ,  $\text{C}_1\text{-C}_4$  alkyl,  $\text{C}_1\text{-C}_4$  alkoxy,  $\text{CH}_2\text{OH}$  or azido; and

$R^{34}$  is  $\text{H}$ ,  $\text{CH}_2\text{CN}$  or  $\text{CF}_3$ , with the proviso that, for structure XXX, when  $R^{25}$  is  $\text{O}$  or  $\text{CH}_2$  and  $R^{29}$   $\text{CH}_2$  or  $\text{O}$ ,  $R^{35}$  is not  $\text{H}$  or  $\text{C}_1\text{-C}_6$  alkyl.

25        The compound of claim 24 wherein  $R^{35}$  is independently  
 20        phenyl, benzyl, adamantoyl oxymethyl, pivaloyloxymethyl, 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 2-ethoxy-5-hydroxyphenyl, 2-ethoxy-4-hydroxyphenyl, 3,5-dimethoxyphenyl, 2,4-difluorophenyl, 2-(haloalkyl)-

phenyl, 3-(haloalkyl)phenyl, 4-(haloalkyl)-phenyl, 2-cyanophenyl, 3-cyanophenyl, 4-cyanophenyl, 2-ethoxyphenyl, 2-carboethoxyphenyl, 3-carboethoxyphenyl, 4-carboethoxyphenyl, or 2-haloalkylbenzyl, 3-haloalkylbenzyl or 4-haloalkylbenzyl.

5

26. The compound of claim 25 wherein B<sup>1</sup> is N<sup>4</sup>-benzoylcytosin-1-yl, N<sup>4</sup>-(6-aminohexyl)cytosin-1-yl, N<sup>4</sup>-(10-aminodecyl)cytosin-1-yl, N<sup>4</sup>-(14-aminolauryl)cytosin-1-yl.

10

27. The compound of claim 1 where L<sup>1</sup> or L<sup>2</sup> is an immunogenic peptide or protein.

28. An antibody capable of binding specifically to a compound of claim 27.


15

29. A method comprising orally administering to a subject an antivirally-effective dose of the compound of claim 2.

30. The method of claim 29 wherein the compound is enriched or resolved at the phosphate atom chiral center.

20

31. A method comprising orally administering to a subject an antivirally-effective dose of the compound of claim 20 having the structure



25


32. The method of claim 31 wherein the compound is enriched or resolved at the phosphate atom chiral center.

33. A compound of formula I, wherein the compound is labeled with a detectable moiety selected from the group of an enzyme, radioisotope, stable free radical, fluorophor, and a chemiluminescent group.

30

34. A compound of the formula  $(R^{31}O)_2P(O)-CH_2-OH$  or  $(R^{31}O)_2P(OSi(CH_3)_3)$  wherein


$R^{31}$  is trimethoxybenzyl; triethoxybenzyl; 2-alkyl pyridinyl ( $C_{1-4}$  alkyl);



wherein  $R^7$  is hydrogen or  $C_{1-4}$  alkyl; or

5  $C_3-C_6$  aryl substituted by 3, 4 or 5 halogen atoms or 1 or 2 atoms or groups selected from halogen,  $C_1-C_{12}$  alkoxy, cyano, nitro,  $C_1-C_{12}$  haloalkyl,  $C_1-C_{12}$  alkyl or  $C_2-12$  alkynyl, provided that for compound  $(R^{31}O)_2P(O)-CH_2-OH$ ,  $R^{31}$  is not phenyl.

10 35. A method to synthesize a compound of structure  $(R^{31}O)_2P(O)-CH_2-OH$  comprising silylating a compound of structure  $(R^{31}O)_2P(O)H$  with about 1 equivalent of bis(trimethylsilyl)trifluoroacetamide, drying the resulting compound and reacting the resulting compound with  
15 paraformaldehyde containing catalytic amounts of a lewis acid, wherein  $R^{31}$  is trimethoxybenzyl; triethoxybenzyl; 2-alkyl pyridinyl ( $C_{1-4}$  alkyl);




wherein  $R^7$  is hydrogen or  $C_{1-4}$  alkyl;


$C_3-C_6$  aryl substituted by 3, 4 or 5 halogen atoms or 1 or 2 atoms or groups selected from halogen,  $C_1-C_{12}$  alkoxy, cyano, nitro,  $C_1-C_{12}$  haloalkyl,  $C_1-C_{12}$  alkyl or  $C_2-12$  alkynyl.

20


36. A method to synthesize a compound of structure  
 $(R^{35}O)_2P(O)-CH_2-R^{29}$



by reacting a compound of structure



with iodine and  $(R^{31}O)_2P(O)-CH_2-OH$  at high temperature,  
wherein  $B^2$  is a heterocyclic base or a protected heterocyclic base;  $R^{31}$  is  
25 trimethoxybenzyl; triethoxybenzyl; 2-alkyl pyridinyl ( $C_{1-4}$  alkyl);



wherein  $R^7$  is hydrogen or  $C_{1-4}$  alkyl;

$C_3-C_6$  aryl substituted by 3, 4 or 5 halogen atoms or 1 or 2 atoms or groups selected from halogen,  $C_1-C_{12}$  alkoxy, cyano, nitro,  $C_1-C_{12}$  haloalkyl,  $C_1-C_{12}$  alkyl or  $C_2-12$  alkynyl; and  $R^{44}$  is iodine or fluorine.

5




Figure 1




Figure 2




Figure 3






Figure 4




Figure 5




Figure 6




Figure 7

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 94/10539

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07H19/10 C07H19/20 A61K31/70 C07F9/6561 C07F9/6509  
C07F9/6571 C07F9/6558 A61K31/685 C07F9/6564 C07F9/36  
C07F9/40 C07F9/6512 C07F9/58 C07F9/09 C07F9/44

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07H A61K C07F G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP,A,0 481 214 (BRISTOL-MYERS SQUIBB COMPANY) 22 April 1992 cited in the application ---	1
A	EP,A,0 398 231 (BRISTOL-MYERS SQUIBB COMPANY) 22 November 1990 cited in the application ---	1
X	NUCLEOSIDES AND NUCLEOTIDES, vol.12, no.3-4, 1993 pages 279 - 294 D.WOLFF-KUGEL ET AL. 'Studies Towards the Synthesis of the Saturated and Unsaturated Carbocyclic Methylene Phosphonate Analogs of Dideoxyadenosine.' see page 284, compound 15 and 7b --- -/--	24

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance  
 "E" earlier document but published on or after the international filing date  
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)  
 "O" document referring to an oral disclosure, use, exhibition or other means  
 "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention  
 "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone  
 "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.  
 "&" document member of the same patent family

Date of the actual completion of the international search

10 January 1995

Date of mailing of the international search report

18. 01. 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

Scott, J

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 G01N33/53

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	NUCLEOSIDES AND NUCLEOTIDES, vol.12, no.2, 1993 pages 157 - 162 J-L KRAUS 'New Phosphonate Analogues of 3'-thia-2',3'-dideoxycytidine (BCH-189).' See page 159, compounds 7a,7b,8a,8b,3a and 3b ---	24
X	TETRAHEDRON LETTERS., vol.33, no.37, 8 September 1992, OXFORD GB pages 5335 - 5338 G.JAHNE ET AL. 'Preparation of Carbocyclic Phosphonate Nucleosides.' See page 5336, structures d,e and f --- -/--	24

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

10 January 1995

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+ 31-70) 340-3016

Authorized officer

Scott, J

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 94/10539

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JOURNAL OF MEDICINAL CHEMISTRY, vol.35, no.17, 21 August 1992, WASHINGTON US pages 3192 - 3196 G.A.FREEMAN ET AL '3'-Azido-3',5'-dideoxyt hymidine-5'-methylphosphonic Acid Diphosphate: Synthesis and HIV-1 Reverse Transcriptase Inhibition.' See page 3192, scheme 1, compound 4 ---	24,25
X	DE,A,41 38 584 (HOECHST AG) 27 May 1993 See page 3, compounds (I) and (II) ---	24
X	WO,A,92 13869 (GILEAD SCIENCES, INC.) 20 August 1992 See Table 3, compounds 26-40, Table 6, compounds 68-82 -----	24,25

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 94/10539

**Box I** Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 24-26, 29-32  
because they relate to subject matter not required to be searched by this Authority, namely:  
Claims searched incompletely: 24-26 (see annex)  
Remark: Although claims 29-32 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II** Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.



FURTHER INFORMATION CONTINUED FROM PCT/ISA/210

LIMITED SEARCH ON BROAD MAIN CLAIM

1. In accordance with the last paragraph of Chapter III, point 3.6 of the PCT Search Guidelines, the subject-matter of intermediate claims ..24 and 25.. has not been exhaustively searched. The only subject-matter of claims 24 and 25 that has been exhaustively searched is that which is further defined in claim 26, directed to preferred compounds of claims 24 and 25.

2. Arguments for limiting the scope of the search can also be found in the following considerations:

3. Each of the claimed compounds comprised in the Markush formula of claim 24 (or the more narrowly defined claim 25) is a compound resulting from the variation of a plurality of variables. These variables embrace a vast number of radicals which are heterogeneous in structure - see for example the possible Z moieties of claim 24, or the R<sup>35</sup> values of claim 25. The only common structural element linking all the possible compounds of claim 1 together, is either a -PO<sub>3</sub>- moiety or a heterocyclic base. These structural elements are common to all nucleotides !

4. There are many compounds falling within the general scope of claim 24 (see for example, either of the Nucleosides and Nucleotides citations mentioned on the search report), and more particularly, there are even a number of compounds that fall within the scope of claim 24, when searched within the limitations of claims 25. This can be seen from WO-A-92/13869, which details (page 16, table 3, compounds 26-29) compounds where R25=O, R26=CH, R27=OH, R28=H, R29=CH<sub>2</sub>, both R35s=Phenyl and the base may be thymine (compound 29), N6-benzoyladenine (28), N4-benzoylcytosine (27) and N2-isobutyrylguanine (26). Compound 27 even falls within the scope of claim 26.

5. In this perspective, starting from the plethora of individual compounds and without making any assumptions which would have no basis in the original disclosure, it is not possible to unambiguously determine a (or a plurality of) generalised set(s) of distinct features which could be considered as special technical features of a solution or of a plurality of

FURTHER INFORMATION CONTINUED FROM PCT/ISA/210

alternative solutions to an accordingly formulated problem underlying the invention as a whole.

6. Consequently, claim 1 does not comply with Article 6 and Rule 6.3(a) PCT, which require that an invention should be clearly defined in terms of features supported by the description. These features should be identifiable by means of an appropriate technical statement of the problem and its solution.

7. The application does not comply with Rule 5.1(a)(ii) and (iii) in that the description does not indicate the common feature(s) of the compounds embraced by the broad definition of claim 1 in a manner sufficient for understanding the invention and carrying out the search.

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.  
PCT/US 94/10539

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0481214	22-04-92	CA-A- 2051239 JP-A- 4230694	15-03-92 19-08-92
EP-A-0398231	22-11-90	AU-B- 646594 AU-A- 2459292 AU-B- 630953 AU-A- 5501290 CA-A- 2015671 JP-A- 3005493	24-02-94 19-11-92 12-11-92 15-11-90 15-11-90 11-01-91
DE-A-4138584	27-05-93	NONE	
WO-A-9213869	20-08-92	AU-A- 1549792 EP-A- 0570514 JP-T- 6507883	07-09-92 24-11-93 08-09-94